

La hemoglobina de Rothschild es una mutación de la cadena  $\beta$  de la hemoglobina donde se sustituye el residuo de triptófano en posición  $\beta37(C3)$  por arginina. El triptófano en B37 (C3) es un punto de contacto en las interfaces  $\alpha1\beta2$  y  $\alpha2\beta1$  y la sustitución del triptófano por la arginina conlleva una desestabilización de las interfaces afectadas durante la oxigenación y la desoxigenación, esperando una desviación del oxígeno normal, pero dicha desviación solo se produce a nivel molecular y no tiene aparente manifestación clínica en los pacientes, que generalmente se mantienen asintomáticos<sup>4,5</sup>. Cabe destacar que la presión parcial a la cual la hemoglobina está medio saturada es p50, la HbA tiene una p50 de  $25 \pm 2$  mmHg, mientras que la hemoglobina de Rothschild [ $\beta 37(C3)$  RTP-->Arg] tiene una p50 mayor por su baja afinidad al oxígeno<sup>4–6</sup>.

La alteración de la saturación de oxígeno en nuestro paciente, determinada mediante pulsioxímetro, fue el dato inicial tras el cual se realiza este estudio. Tras identificar como causa de este problema la hemoglobinopatía y explicarle al paciente la baja implicación clínica de este hallazgo, el paciente minimizó sus problemas y síntomas, que habían sido sobredimensionados por la conciencia de las alteraciones detectadas y el miedo provocado ante la sospecha de patología médica. Por todo ello debemos considerar que, ante un paciente con saturaciones de oxígeno bajas sin causa evidente y baja concordancia clínica, uno de los motivos a tener en cuenta es este tipo de hemoglobinopatías, que, aunque infrecuentes, pueden aparecer en nuestro quehacer diario.

Además, para completar el estudio familiar, e identificar otros afectos, se usó la pulsioximetría, que permitió identificar como portadores a su hermana y a 2 sobrinos.

Podemos considerar que ante un paciente con saturaciones de oxígeno bajas sin causa aparente de patología que justifique la hipoxia<sup>4,5,7</sup>, debemos pensar que se trate de una alteración de la afinidad del oxígeno<sup>7–9</sup> por la hemoglobina, sabiendo que muchas de ellas cursan sin síntomas<sup>5</sup> y que con una medición de la saturación mediante un pulsioxímetro podremos detectar patologías infrecuentes<sup>10,11</sup> como es el caso que nos ha ocupado.

## Bibliografía

- Sinex JE. Pulse oximetry: Principles and limitations. Am J Emerg Med. 1999;17:59–67.

- Neff TA. Routine oximetry. A fifth vital sign? Chest. 1988;94:227.
- Kasper D, Hauser S, Jameson J, Fauci A, Longo D, Loscalzo J. Hemoglobinas con alteración de la afinidad al oxígeno. Principios de Medicina Interna. Harrison. 19.<sup>a</sup> ed. 2015; Cap 127, pp. 636.
- Alli N, Wessels P, Rampersad N, Clark B, Thein S. Detection of Hb Rothschild HBB: c.(112T>A or 112T>C), through high of suspicion on abnormal pulse oximetry. Hemoglobin. 2017;41:137–9.
- Gacon G, Belkhodja O, Wajcman H, Labie D, Najman A. Structural and functional studies of Hb Rothschild beta (C3) Trp replaced by Arg. A new variant of the alpha1beta2 contact. FEBS Lett. 1977;82:243–6.
- Sharma V, Newton G, Ranney H, Ahmed F, Harris J, Danish E. Hemoglobin Rothschild (beta 37(C3)Trp replaced by Arg): A high/low affinity hemoglobin mutant. J Mol Biol. 1980;144:267–80.
- Hanning C, Alexander-Williams J. Pulse oximetry: A practical review. BMJ. 1995;311:367–70.
- Hladik A, Lynshue K. Dyshemoglobinemas and pulse oximetry: A therapeutic challenge. J Pediatr Hematol Oncol. 2008;30:850–2.
- Burns C, Thet L, Woodson R, Schultz J, Hla K. Hemoglobinopathy case finding by pulse oximetry. Am J Hematol. 2003;74:142–3.
- Craik C, Vallette I, Beychok S, Waks M. Refolding defects in hemoglobin Rothschild. J Biol Chem. 1979;255:6219–23.
- Corkeron M, Pavly T. Pulse oximetry in mixed hemoglobinopathy. Anaesth Intensive Care. 1996;24:619–20.

Luis Manuel González García <sup>a</sup>, José María Hernández Pérez <sup>b,\*</sup>, María Tapia Torres <sup>c</sup> y Paloma Ropero Gradilla <sup>d</sup>

<sup>a</sup> Centro de Salud de Breña Baja, Área de Salud de La Isla de La Palma, Breña Baja, Santa Cruz de Tenerife, España

<sup>b</sup> Sección de Neumología, Hospital General de La Palma, Breña Alta, La Palma, Santa Cruz de Tenerife, España

<sup>c</sup> Servicio de Hematología, Hospital General de La Palma, Breña Alta, La Palma, Santa Cruz de Tenerife, España

<sup>d</sup> Servicio de Hematología y Hemoterapia, Hospital Clínico San Carlos, Madrid, España

\* Autor para correspondencia.

Correo electrónico: jmherper@hotmail.com (J.M. Hernández Pérez).

<https://doi.org/10.1016/j.arbres.2019.05.009>

0300-2896/

© 2019 SEPAR. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

## Echogenic Swirling Seen on Ultrasound and Outcome of Pleurodesis in Malignant Pleural Effusion

### Ecografía con signo de remolino y resultado de la pleurodesis en un derrame pleural maligno

Dear Editor:

Malignant pleural effusion (MPE) is encountered in fifteen percent of patients with cancer<sup>1</sup> and frequently causes breathlessness that recurs after aspiration which necessitates more definitive treatment such as indwelling pleural catheter insertion or pleurodesis.<sup>2</sup> Talc is the agent used for pleurodesis in the UK<sup>2</sup> due to its superior efficacy compared to other agents<sup>1</sup> with success rates reported from large trials ranging between 75 and 80%.<sup>3,4</sup> About one quarter of the patients receiving talc will require further procedures. Some parameters, such as pleural fluid pH and adenosine deaminase<sup>5,6</sup> have been correlated with pleurodesis outcome but these results have not been replicated in other studies and their overall clinical usefulness remains questionable.

Thoracic ultrasound (TUS) is an essential tool in the evaluation of pleural effusion.<sup>7</sup> Echogenic swirling (ES), a finding which describes



multiple floating echogenic particles inside an effusion (supplementary video), is seen on TUS examination of some patients with MPE and has been hypothesised to be related to enhanced intrapleural fibrinolytic activity.<sup>8</sup> Effusions of MPE patients who failed pleurodesis were shown to have higher levels of fibrinolytic enzymes<sup>9</sup> and MPE samples obtained following intapleural application of sclerosants tended to exhibit more sustained elevation in levels of fibrin degradation products in patients who failed the procedure.<sup>10,11</sup> This study aimed to assess whether an ultrasound parameter available prior to intervention (ES) is associated with reduced rates of pleurodesis success.

Data on patients who underwent thoracoscopic talc poudrage or talc slurry instillation via chest drain for the management of MPE at the authors' Pleural Unit between 2016 and 2017 were collected. Baseline clinical details and pleural fluid biochemistry results were retrieved. Recorded TUS clips of pre-procedure effusions were assessed for the presence of ES.

Pleurodesis failure was defined as the need for another therapeutic procedure within the first 3 months after pleurodesis.

Patients were excluded if there were no assessable TUS clips on the database, post-drainage X-ray showed any degree of trapped lung, or data on pleurodesis outcome was not available.

**Table 1**

Baseline characteristics and pleural fluid composition of the study cohort and according to pleurodesis outcome. Data for continuous variables is presented as mean + SD or median (IQR) and for discrete variables as frequency (per cent).

Characteristic	Study cohort (n = 44)	Success (n = 21)	Failure (n = 23)	P value
Age years	74 + 10	71 + 8	76 + 11	.063 (T-test)
Sex (male/female)	21/23	14/7	9/14	.061 ( $\chi^2$ )
Primary tumour				
Mesothelioma	18 (40.9%)	8 (38%)	10 (43.4%)	.41 ( $\chi^2$ )
Lung	7 (15.9%)	4 (19%)	3 (13%)	
Breast	10 (22.7%)	3 (14.2%)	7 (30.4%)	
Other	9 (20.4%)	6 (28.5%)	3 (13%)	
P.F. glucose mmol/L	4.6 + 1.8	4.2 + 1.3	5 + 2.2	.217 (T-test)
P.F. protein g/L	41.5 + 7.3	44 + 4.6	39 + 10	.03 (T-test) [mean dif 4.8 g/L, 95% CI 4–9.1 g/L]
P.F. LDH IU/L	267 (139–571)	268 (216–501)	249 (133–1092)	.98 (Mann–Whitney)
Pleurodesis method (Poudrage)	24 (54.5%)	15 (71.4%)	9 (39.1%)	.032 ( $\chi^2$ 4.61)
Systemic treatment	29 (65.9%)	19 (90.4%)	10 (43.4%)	.001 ( $\chi^2$ 10.79)
Echogenic swirling	26 (59.1%)	7 (33.3%)	19 (82.6%)	.001 ( $\chi^2$ 11.01)

IQR: interquartile range; SD: standard deviation.

Overall rate of pleurodesis success as well as according to pleural fluid echogenicity was presented. All factors that are hypothesised to affect pleurodesis were entered into a multiple logistic regression analysis to identify variables associated with pleurodesis failure.

Sixty nine patients underwent pleurodesis for MPE in the period of study. Forty four patients were included, with the remaining excluded due to the absence of assessable TUS clips (18 patients), trapped lung (2 patients), or death prior to follow up visit (5 patients). Table 1 shows the clinical and pleural fluid characteristics of the studied patients. Pleurodesis failure was noted in 23/44 (52.2%) of the study cohort. Mean pleural fluid protein was higher in patients who failed pleurodesis (mean dif 4.8 g/L, 95% CI 0.4–9.1 g/L). Regarding the relation between pleurodesis outcome and the presence of ES, ES was found in 7/21 (33.3%) with successful pleurodesis and in 19/23 (82.6%) with failed pleurodesis ( $\chi^2 = 11.01$ , 1df,  $p = 0.001$ ).

The levels of protein, LDH and glucose did not differ significantly between effusions exhibiting ES and those without this sign. Mean protein was  $41 \pm 8$  g/L in effusion with ES vs.  $41 \pm 6$  g/L in those without ES. Median (IQR) LDH was 251 (11–885) IU/L with ES vs. 268 (166–348) in effusions without ES and median glucose was 4.4 mmol/L (3.5–6.1) vs. 4.2 (3.3–5.7) in effusion without ES.

Regarding the primary cause of MPE, 12/18 (66.6%) cases with mesothelioma had ES on TUS examination, while ES was seen in 4/7 (57.1%) patients with lung cancer and in 6/10 (60%) patients with breast cancer ( $\chi^2 = 1.24$ , 2df,  $p = 0.74$ ).

Multiple logistic regression analysis on pleurodesis success as the dependent variable included the presence of ES together with institution of systemic cancer therapy, talc poudrage as the method of pleurodesis and pleural fluid protein level as the independent variables. The factors independently associated with pleurodesis success according to regression analysis were the presence of ES on US (odds ratio 0.016, 95% CI 0.001–0.310,  $p = 0.006$ ) and lack of systemic therapy (odds ratio 0.014, 95% CI 0.001–0.304,  $p = 0.007$ ).

This study suggests that ES seen on TUS examination of patients with MPE may be associated with lower rates of success of pleurodesis and this is independent of other parameters such as the presence of systemic therapy.

MPEs are characterised by enhanced inflammatory activity and fibrin deposition,<sup>12</sup> coupled with high levels of fibrinolytic enzymes and fibrin degradation products.<sup>12</sup> It is thought that the echogenic material seen floating on TUS examination of MPE could be the broken down fibrinous material from the aforementioned processes.<sup>8</sup>

The association of ES with lower success rates of pleurodesis could be linked to the previously noted high levels of fibrinolytic enzymes such as u-PA in MPE patients who failed pleurodesis.<sup>9</sup>

The remarkably lower rates of pleurodesis success in cases with ES might thus be explained by failure of pleural fibrosis and symphysis that characterises pleurodesis and which could be antagonised by high fibrinolytic activity in such effusions. Indeed higher levels of D-dimer (a marker of fibrinolysis) have been noted in MPE with failed pleurodesis in comparison to successful pleurodesis.<sup>10,11</sup> Notably, the chemical composition of effusions showing ES was not different from that of effusions not showing ES.

The presence of ES was judged in a binary fashion which could be an oversimplification of a variable that can manifest with various degrees of intensity. It is not possible using current ultrasound machines to objectively quantify the degree of echogenicity of an effusion. However, the concept of 'echogenicity index' has been used to quantify the degree of echogenicity of structures such as thyroid nodules<sup>13</sup> and renal parenchyma<sup>14</sup> using softwares that compare the echogenicity of a structure in question to adjacent 'reference' organs to gain clinically relevant information.

The decision to exclude patients who had trapped lung (which is related to the extent of visceral pleural involvement) from the analysis was intended to remove confounding effect of heavier pleural tumour burden on pleurodesis failure. The definition of pleurodesis failure was based on objective criteria that mirror the definition used in large clinical trials of pleurodesis<sup>3</sup> in order to ensure the robustness of the data obtained.

This study has several limitations; we found a higher than usual failure rate of pleurodesis (52%). The studied cohort has a high proportion of mesothelioma cases which is known to respond less favourably to pleurodesis<sup>15</sup> and reflects differences that are seen often between data derived from trials and that seen in the 'real-world'. Despite the fact that this study involved a relatively small number of patients, there was strong signal to support a link between pleural fluid echogenicity and pleurodesis outcome.

The proposed TUS-based observation of ES as a tool for outcome prediction of pleurodesis is a potentially useful non-invasive and easy to interpret tool, which is importantly available to the clinician prior to any decision on intervention. This may therefore help towards better customisation of management according to patients' characteristics. Our results require prospective validation in a larger patient cohort and correlation with other variables such as pleural fluid cytological positivity and the presence of pleural thickening. The role that a pro-fibrotic/fibrinolytic balance may

play in determining pleural fluid echogenicity and pleurodesis outcome and the usefulness of quantifying pleural fluid echogenicity are areas of potential future research.

## Acknowledgement

This work uses data provided by patients and collected by the NHS as part of their care and support, and is therefore considered an audit not requiring ethical approval.

## Appendix A. Supplementary Data

Supplementary data associated with this article can be found, in the online version, at doi:10.1016/j.arbres.2019.05.010.

## Bibliografía

- Clive AO, Jones HE, Bhatnagar R, Preston NJ, Maskell N. Interventions for the management of malignant pleural effusions: a network meta-analysis. In: The cochrane collaboration cochrane database of systematic reviews. Chichester, UK: John Wiley & Sons Ltd.; 2016.
  - Roberts ME, Neville E, Berriford RG, Antunes G, Ali NJ, on behalf of the BTS Pleural Disease Guideline Group. Management of a malignant pleural effusion: British Thoracic Society pleural disease guideline 2010. Thorax. 2010;40–65.
  - Rahman NM, Pepperell J, Rehal S, Saba T, Tang A, Ali N, et al. Effect of opioids vs NSAIDs and larger vs smaller chest tube size on pain control and pleurodesis efficacy among patients with malignant pleural effusion: the TIME1 randomized clinical trial. JAMA. 2015;314:263–2641.
  - Dresler CM, Olak J, Herndon JE, Richards WG, Scalzetti E, Fleishman SB, et al. Phase III intergroup study of talc pouddrage vs talc slurry sclerosis for malignant pleural effusion. Chest. 2005;127:15–909.
  - Heffner JE, Nietert PJ, Barbieri C. Pleural fluid pH as a predictor of pleurodesis failure. Chest. 2000;117:87–95.
  - Yildirim H, Metintas M, Ak G, Metintas S, Erginol S. Predictors of talc pleurodesis outcome in patients with malignant pleural effusions. Lung Cancer. 2008;62:44–139.
  - Wangüemert Pérez AL. Is thoracic ultrasonography necessary in the respiratory medicine outpatient clinic? Arch Bronconeumol. 2017.
  - Chian C-F, Su W-L, Soh L-H, Yan H-C, Perng W-C, Wu C-P. Echogenic swirling pattern as a predictor of malignant pleural effusions in patients with malignancies. Chest. 2004;126:34–129.
  - Ishikawa H, Satoh H, Hasegawa S, Yazawa T, Naito T, Yamashita YT, et al. Urokinase-type plasminogen activator in carcinomatous pleural fluid. Eur Respir J. 1997;10:71–1566.
  - Rodríguez-Panadero F, Segado A, Martín Juan J, Ayerbe R, Torres García I, Castillo J. Failure of talc pleurodesis is associated with increased pleural fibrinolysis. Am J Respir Crit Care Med. 1995;151:90–785.
  - Psathakis K, Calderón-Osuna E, Romero-Romero B, Martín-Juan J, Romero-Falcón A, Rodríguez-Panadero F. The neutrophilic and fibrinolytic response to talc can predict the outcome of pleurodesis. Eur Respir J. 2006;27:21–817.
  - Vaz MAC, Vargas FS, Marinho FCdA, D'Amico EA, Rocha TRF, Teixeira LR. Does the evaluation of coagulation factors contribute to etiological diagnosis of pleural effusions? Clinics (Sao Paulo). 2009;64:5–891.
  - Wu M-H, Chen C-N, Chen K-Y, Ho M-C, Tai H-C, Wang Y-H, et al. Quantitative analysis of echogenicity for patients with thyroid nodules. Sci Rep. 2016;6:35632.
  - Manley JA, O'Neill WC. How echogenic is echogenic? Quantitative acoustics of the renal cortex. Am J Kidney Dis. 2001;37:11–706.
  - Chen J, Li Z, Xu N, Zhang X, Wang Y, Lin D. Efficacy of medical thoracoscopic talc pleurodesis in malignant pleural effusion caused by different types of tumors and different pathological classifications of lung cancer. Int J Clin Exp Med. 2015;8:53–18945.
- Maged Hassan <sup>a,b,\*</sup>, Rachelle Asciak <sup>a</sup>, Rachel M. Mercer <sup>a</sup>, David J. McCracken <sup>a</sup>, Stamatoula Tsikrika <sup>c</sup>, Hany Shaarawy <sup>b</sup>, Anwar Elganady <sup>b</sup>, John M. Wrightson <sup>a</sup>, Najib M. Rahman <sup>a,d</sup>
- <sup>a</sup> Oxford Pleural Unit, Oxford University Hospitals NHS Foundation Trust, UK  
<sup>b</sup> Chest Diseases Department, Faculty of Medicine, Alexandria University, Egypt  
<sup>c</sup> Pulmonary Division, Department of Critical Care, Medical School, National and Kapodistrian University of Athens, Evangelismos General Hospital, Athens, Greece  
<sup>d</sup> Oxford NIHR Biomedical Research Centre, Oxford, UK
- \* Corresponding author.  
E-mail address: magedhmf@gmail.com (M. Hassan).

<https://doi.org/10.1016/j.arbres.2019.05.010>

## Neumomediastino y enfisema subcutáneo espontáneo: síndrome de Hamman



### Spontaneous Pneumomediastinum and Subcutaneous Emphysema: Hamman's Syndrome

Estimado Director:

El neumomediastino se define por la presencia de aire libre en el mediastino y suele acompañarse de enfisema subcutáneo. Se clasifica en espontáneo cuando no existe una causa evidente y en traumático cuando ocurre de manera secundaria a una intervención, perforación o traumatismo a dicho nivel. El neumomediastino espontáneo se debe a la rotura de los alvéolos terminales secundaria a una hiperinsuflación traqueal<sup>1</sup>. Una de las causas menos conocidas de neumomediastino espontáneo es el síndrome de Hamman (descrito por Louis Hamman en 1939), que hace referencia al neumomediastino espontáneo asociado a enfisema subcutáneo que se produce en el posparto<sup>2,3</sup>. Es una entidad poco frecuente, con tan solo 200 casos descritos en la literatura hasta 1994<sup>4</sup>. Exponemos el caso de una mujer de 29 años que tras un parto vaginal presentó neumomediastino y enfisema subcutáneo.

Mujer de 29 años primigesta de 40+3 semanas, sin antecedentes médicos, que acudió a urgencias por dinámica uterina irregular cada 15 min de 12 h de evolución. No presentaba sangrado vaginal ni rotura de bolsa amniótica. La paciente había realizado un correcto seguimiento del embarazo, sin incidencias.

En el inicio de la dinámica de parto, una ecografía abdominal y un registro cardiotocográfico fetal no mostraron alteraciones. La paciente tuvo un parto espontáneo eutóxico con presentación cefálica, siendo el recién nacido un varón de 3,2 kg con Apgar de 10 en el primer minuto. El parto duró 10 h y no se registraron complicaciones.

A las 18 h de finalizar el parto la paciente presentó odinofagia y sensación de crepitación en la región cervical, sin disnea. La presión arterial era de 126/73 mmHg, la frecuencia cardíaca de 80 lpm y la saturación de oxígeno del 98% respirando aire ambiente. La auscultación pulmonar era normal. Una radiografía de tórax mostró enfisema subcutáneo en ambas regiones claviculares y cervical bilateral, sin evidencia de neumotórax. Un análisis de sangre mostró hemoglobina de 11,8 g/dL, 113.000 plaquetas/mm<sup>3</sup>, 12.700 leucocitos/mm<sup>3</sup> (82,3% neutrófilos). La función renal, iones y función hepática eran normales. Una tomografía computarizada (TC) de tórax demostró un extenso enfisema subcutáneo y profundo cervical y extenso neumomediastino. Se descartó la presencia de mediastinitis, líquido libre o rotura esofágica. Se realizó el diagnóstico de neumomediastino espontáneo secundario a trabajo de parto y se decidió un manejo conservador. La evolución posterior fue favorable, encontrándose asintomática y con resolución completa de las lesiones a las 2 semanas de seguimiento.

El síndrome de Hamman, descrito en 1939, se define por la asociación de enfisema subcutáneo extenso y neumomediastino en el puerperio inmediato y es una rara complicación asociada al trabajo del parto, con una incidencia aproximada de uno de cada