

Scientific Letter

Phenotype Analysis of Chronic Bronchial Infection Patients on Inhaled Antibiotic Therapy: A Multicentre Retrospective Cohort Study in Spain (INBREATHING Study)



To the Director,

Chronic bronchial infection (CBI) involves persistent inflammation and impaired mucociliary clearance, increasing exacerbations and symptom burden across structural lung diseases.¹ These events reduced quality of life and increase mortality.^{2,3} Long-term inhaled antibiotics (IA) can reduce bacterial load, eradicate pathogens, decrease exacerbations, and improve symptoms and quality of life.^{4–6} However, most evidence comes from cystic fibrosis (CF), with limited data in non-CF bronchiectasis or other structural lung diseases.^{7–9} Heterogeneity in underlying conditions contributes to inconsistent bronchiectasis trial results, limiting patient selection and statistical power.¹⁰ This highlights the need to understand differential IA responses in CBI to optimize future trials.^{10,11} Phenotyping approaches have been used in chronic obstructive pulmonary disease (COPD), bronchiectasis, and asthma to identify clinical clusters with distinct prognoses and therapeutic responses.¹⁰ However, such analyses have not yet been applied to CBI populations treated with IA.

The INBREATHING study is a retrospective, multicentre cohort study conducted across 10 Spanish hospitals, including 402 adult patients with CBI treated with IA and followed in outpatient clinics between January 2018 and June 2024. CBI was diagnosed according to national guideline criteria.^{1,12} Baseline clinical, functional, and microbiological data were collected at IA initiation. One-year follow-up assessed changes in exacerbations, symptoms, and treatment tolerance. Pre-treatment clinical data were used to identify phenotypes within the cohort, including demographic variables, comorbidities, underlying respiratory diseases, lung function, and chronic therapies. To identify phenotypes and evaluate its reproducibility, the cohort was randomly split into training and test subgroups (1:1). Phenotypes were derived in the training set using k-prototypes clustering with multiple imputation¹³ and reproducibility was assessed in the test set. Then, each dataset was clustered by the k-prototypes algorithm. Finally, hierarchical cluster with average linkage was used to assigned patient in each phenotype. The number of phenotypes was determined using elbow method based on the summation of within-cluster distances for classes ranging from 0 to 10. Clinical differences between phenotypes were analyzed, and their association with one-year exacerbation risk after IA treatment was evaluated using multivariable logistic regression. Predictor selection was based on bivariate analyses and refined by backward stepwise selection using the Akaike Information Criterion. Model performance was assessed via

the area under the ROC curve, and results were visualized using forest plots.

The cohort included 402 patients (mean age of 71.6 ± 13.3 years; 55.2% male). Common comorbidities were history of cancer (18.4%), heart disease (15.2%), atrial fibrillation (14.7%), and kidney disease (10.4%). The most prevalent underlying respiratory diseases were bronchiectasis (77.8%) and COPD (39.6%). Among bronchiectasis patients, the most common radiological pattern was cylindrical (78.1%), followed by varicose (23.5%) and cystic (17.0%). Mean FEV₁ was 65.8 (25.1) percent predicted value, with 39.4% showing mild-to-moderate and 29.17% severe impairment. Inhaled corticosteroids were used by 65.9%, and 19.2% received alternate-day azithromycin. *Pseudomonas aeruginosa* was the most common pathogen (81.1%); colistin (72.9%) and tobramycin (14.7%) were the main IA treatments. Median IA duration was 12.0 months (IQR: 5.6–23.8). Further details are provided in Table 1.

Three reproducible phenotypes were identified using k-prototypes clustering with multiple imputation (Table 1 and Fig. 1 Panel A). Phenotype 1 ($n = 138$, 34.3%) was mainly composed of women (56.5%) with preserved lung function (mean FEV₁ $93.9 \pm 13.8\%$ predicted), predominantly bronchiectasis (81.9%) and the highest asthma prevalence (12.3%). Phenotype 2 ($n = 148$, 36.8%) showed moderate airflow obstruction (FEV₁ $62.6 \pm 25.1\%$), with frequent bronchiectasis (79.7%) and COPD (38.5%). Phenotype 3 ($n = 116$, 28.8%) had severe obstruction (FEV₁ $35.9 \pm 8.3\%$), a higher prevalence of COPD, more comorbidities, older age, and greater use of dual or triple inhaled therapy. There were no significant differences between phenotypes in terms of IA type or delivery device.

After one year of IA therapy, all phenotypes showed symptomatic improvement and a reduction in exacerbations (Table 1). However, phenotype 3 remained at highest risk, with a mean exacerbation rate of 1.31 per year and 63.8% of patients experiencing ≥ 1 event (Table 1 and Fig. 1 Panel B). This group also had the poorest symptomatic response and highest treatment intolerance, leading to a 33.0% discontinuation rate due to adverse effects (Table 1). In contrast, phenotypes 1 and 2 had lower exacerbation rates and fewer recurrent events. Notably, phenotype 1 was the only group with a significant decline in lung function (FEV₁ -5.96% , FVC -5.54%) (Table 1). Bacterial eradication rates did not differ significantly between phenotypes. A multivariable predictive model (Fig. 1 Panel C) identified prior exacerbations (OR 1.37; 95% CI: 1.19–1.59), varicose/cystic bronchiectasis on chest CT (OR 1.84; 95% CI: 1.08–3.16), number of comorbidities (OR 1.48; 95% CI: 1.11–2.01), age (OR 1.01; 95% CI: 0.99–1.03), and phenotype classification (OR 0.93; 95% CI: 0.53–1.64 for phenotype 2; OR 2.36; 95% CI: 1.27–4.42 for phenotype 3) as independent predictors of exacerbation risk, with moderate discriminative performance (AUC 0.74). A web-based application (<https://trrm.shinyapps.io/IAscore>) and a

Table 1

Clinical characteristics, exacerbation rates, longitudinal changes in pulmonary function parameters and other outcomes before and after Inhaled antibiotic treatment by phenotypes.

| Baseline characteristics | Global N = 402 | Phenotype 1 n = 138 | Phenotype 2 n = 148 | Phenotype 3 n = 116 | p value |
|---|---------------------|---------------------|---------------------|---------------------|---------|
| Demographic | | | | | |
| Age (years) | 71.6 (13.3) | 70.5 (13.4) | 71.8 (14.3) | 72.7 (11.6) | 0.386 |
| Sex (Female) | 180 (44.8%) | 78 (56.5%) | 68 (45.9%) | 34 (29.3%) | |
| Underlying lung disease | | | | | |
| COPD | 159 (39.6%) | 22 (15.9%) | 57 (38.5%) | 80 (69.0%) | <0.001 |
| GOLD 0 | 12 (7.59%) | 3 (13.6%) | 6 (10.5%) | 3 (3.80%) | |
| GOLD 1 | 16 (10.1%) | 13 (59.1%) | 3 (5.26%) | 0 (0.00%) | |
| GOLD 2 | 39 (24.7%) | 1 (4.55%) | 35 (61.4%) | 3 (3.80%) | |
| GOLD 3 | 35 (22.2%) | 0 (0.00%) | 6 (10.5%) | 29 (36.7%) | |
| GOLD 4 | 56 (35.4%) | 5 (22.7%) | 7 (12.3%) | 44 (55.7%) | |
| Bronchiectasis | 311 (77.4%) | 113 (81.9%) | 118 (79.7%) | 80 (69.0%) | 0.034 |
| Cylindrical | 243 (60.4%) | 86 (62.3%) | 94 (63.5%) | 63 (54.3%) | 0.271 |
| Cystic | 53 (13.2%) | 7 (5.07%) | 30 (20.3%) | 16 (13.8%) | 0.001 |
| Varicose | 73 (18.2%) | 23 (16.7%) | 30 (20.3%) | 20 (17.2%) | 0.699 |
| Asthma | 29 (7.21%) | 17 (12.3%) | 10 (6.76%) | 2 (1.72%) | 0.005 |
| Comorbidities | | | | | |
| Total number | 0.78 (0.90) | 0.59 (0.79) | 0.70 (0.80) | 1.11 (1.06) | <0.001 |
| Cancer | 74 (18.4%) | 24 (17.4%) | 21 (14.2%) | 29 (25.0%) | 0.074 |
| Heart failure | 61 (15.2%) | 13 (9.42%) | 21 (14.2%) | 27 (23.3%) | 0.008 |
| Kidney failure | 42 (10.4%) | 12 (8.70%) | 13 (8.78%) | 17 (14.7%) | 0.214 |
| Diabetes mellitus | 49 (12.2%) | 8 (5.80%) | 19 (12.8%) | 22 (19.0%) | 0.006 |
| Pulmonary function | | | | | |
| FEV ₁ (%) | 65.8 (25.1) | 93.9 (13.8) | 62.6 (8.69) | 35.9 (8.26) | <0.00 |
| FVC (%) | 79.3 (21.6) | 100 (14.1) | 76.8 (11.9) | 56.9 (13.3) | <0.001 |
| FEV ₁ to FVC ratio | 63.0 (14.3) | 73.3 (8.73) | 63.1 (11.2) | 50.0 (13.2) | <0.001 |
| DLCO (%) | 72.9 (21.9) | 85.6 (15.2) | 66.4 (22.2) | 58.2 (19.9) | <0.001 |
| Previous pharmacological treatment | | | | | |
| Inhaled treatment | | | | | NE |
| No | 75 (18.7%) | 40 (29.0%) | 24 (16.2%) | 11 (9.48%) | |
| Long-acting β ₂ -agonist | 64 (15.9%) | 39 (28.3%) | 17 (11.5%) | 8 (6.90%) | |
| Long-acting muscarinic antag. | 15 (3.73%) | 8 (5.80%) | 4 (2.70%) | 3 (2.59%) | |
| Both | 248 (61.7%) | 51 (37.0%) | 103 (69.6%) | 94 (81.0%) | |
| Inhaled corticosteroids | 265 (65.9%) | 73 (52.9%) | 101 (68.2%) | 91 (78.4%) | <0.001 |
| Hypertonic saline | 63 (15.8%) | 18 (13.1%) | 25 (17.1%) | 20 (17.2%) | 0.576 |
| Azithromycin (alternate days) | 75 (19.2%) | 20 (14.8%) | 23 (16.0%) | 32 (28.6%) | 0.011 |
| Nebulized antibiotic treatment | | | | | |
| Type | | | | | 0.972 |
| Colistin | 293 (72.9%) | 97 (70.3%) | 105 (70.9%) | 91 (78.4%) | |
| Tobramycine | 59 (14.7%) | 19 (13.8%) | 26 (17.6%) | 14 (12.1%) | |
| Gentamycine | 21 (5.22%) | 9 (6.52%) | 9 (6.08%) | 3 (2.59%) | |
| Amikacin | 21 (5.22%) | 9 (6.52%) | 6 (4.05%) | 6 (5.17%) | |
| Other | 8 (1.99%) | 4 (2.90%) | 2 (1.35%) | 2 (1.72%) | |
| Device | | | | | 0.241 |
| INeb | 45 (11.3%) | 12 (8.89%) | 19 (12.8%) | 14 (12.3%) | |
| Jet | 140 (35.3%) | 58 (43.0%) | 49 (33.1%) | 33 (28.9%) | |
| Vibrating mesh | 204 (51.4%) | 64 (47.4%) | 77 (52.0%) | 63 (55.3%) | |
| Dry powder | 8 (2.02%) | 1 (0.74%) | 3 (2.03%) | 4 (3.51%) | |
| Exacerbation rates | Global n = 352 | Phenotype 1 n = 133 | Phenotype 2 n = 142 | Phenotype 3 n = 115 | p value |
| Mild-to-moderate exacerbations | | | | | |
| One-year before inhaled antibiotics | 78.9% [74.5%;82.9%] | 78.9% [71.0%;85.5%] | 78.7% [71.0%;85.2%] | 79.1% [70.6%;86.1%] | 0.997 |
| Number | 1.94 [1.77;2.10] | 2.09 [1.79;2.39] | 1.82 [1.55;2.08] | 1.90 [1.60;2.21] | 0.382 |
| One-year after inhaled antibiotics | 36.9% [31.8%;42.2%] | 35.0% [26.5%;44.4%] | 32.6% [24.8%;41.2%] | 45.3% [35.0%;55.8%] | 0.129 |
| Number | 0.61 [0.50;0.71] | 0.49 [0.34;0.63] | 0.56 [0.39;0.73] | 0.82 [0.57;1.07] | 0.044 |
| Change | −1.49 [−1.67;−1.31] | −1.63 [−1.93;−1.33] | −1.45 [−1.74;−1.17] | −1.37 [−1.76;−0.98] | 0.521 |
| Severe exacerbations (hospitalization) | | | | | |
| One-year before inhaled antibiotics | 38.9% [34.0%;43.9%] | 34.1% [26.1%;42.8%] | 29.6% [22.2%;37.8%] | 56.2% [46.6%;65.6%] | <0.001 |
| Number | 0.61 [0.51;0.72] | 0.41 [0.30;0.52] | 0.46 [0.32;0.60] | 1.05 [0.79;1.32] | <0.001 |
| One-year after inhaled antibiotics | 12.1% [8.86%;16.0%] | 4.27% [1.40%;9.69%] | 5.88% [2.57%;11.3%] | 30.9% [21.7%;41.2%] | <0.001 |
| Number | 0.18 [0.13;0.24] | 0.06 [0.00;0.12] | 0.07 [0.02;0.11] | 0.51 [0.33;0.69] | <0.001 |
| Change | −0.38 [−0.49;−0.28] | −0.35 [−0.46;−0.24] | −0.35 [−0.48;−0.21] | −0.48 [−0.78;−0.18] | 0.551 |
| Total exacerbations | | | | | |
| One-year before inhaled antibiotics | 88.8% [85.2%;91.8%] | 87.0% [80.0%;92.3%] | 86.5% [79.8%;91.7%] | 93.8% [87.5%;97.5%] | 0.142 |
| Number | 2.53 [2.34;2.72] | 2.47 [2.15;2.80] | 2.26 [1.97;2.55] | 2.94 [2.54;3.33] | 0.018 |
| One-year after inhaled antibiotics | 43.4% [38.1%;48.8%] | 35.9% [27.2%;45.3%] | 35.6% [27.5%;44.2%] | 63.8% [53.3%;73.5%] | <0.001 |
| Number | 0.78 [0.65;0.91] | 0.55 [0.38;0.71] | 0.61 [0.44;0.79] | 1.31 [0.99;1.62] | <0.001 |
| Change | −1.71 [−1.91;−1.51] | −1.83 [−2.14;−1.51] | −1.66 [−1.96;−1.36] | −1.64 [−2.11;−1.17] | 0.720 |

Table 1 (Continued)

| Longitudinal changes in pulmonary function parameters | Global <i>n</i> = 352 | Phenotype 1 <i>n</i> = 120 | Phenotype 2 <i>n</i> = 136 | Phenotype 3 <i>n</i> = 98 | <i>p</i> value |
|---|-----------------------|----------------------------|----------------------------|----------------------------|----------------|
| <i>FEV₁</i> | | | | | |
| Baseline | 65.8 [63.1;68.4] | 93.9 [91.4;96.4] | 62.6 [61.2;64.1] | 35.9 [34.3;37.6] | <0.001 |
| One-year | 64.2 [61.0;67.4] | 86.9 [82.8;91.0] | 63.1 [59.9;66.4] | 38.7 [35.8;41.7] | <0.001 |
| Change | −1.28 [−3.21;0.66] | −5.96 [−9.90;−2.03] | 1.18 [−1.96;4.33] | 1.07 [−1.43;3.56] | 0.003 |
| <i>FVC</i> | | | | | |
| Baseline | 79.3 [77.0;81.5] | 100 [97.9;103] | 76.8 [74.8;78.8] | 56.9 [54.2;59.6] | <0.001 |
| One-year | 76.9 [74.0;79.7] | 93.6 [89.8;97.4] | 77.1 [73.8;80.3] | 57.0 [52.8;61.1] | <0.001 |
| Change | −1.16 [−3.15;0.83] | −5.54 [−9.11;−1.97] | 1.93 [−1.20;5.05] | 0.02 [−3.55;3.59] | 0.005 |
| <i>FEV₁/FVC</i> | | | | | |
| Baseline | 63.0 [61.5;64.5] | 73.3 [71.7;74.9] | 63.1 [61.2;65.0] | 50.0 [47.3;52.6] | <0.001 |
| One-year | 63.0 [61.2;64.9] | 72.1 [69.8;74.5] | 63.6 [60.9;66.2] | 51.8 [48.6;55.1] | <0.001 |
| Change | −0.52 [−1.96;0.92] | −0.29 [−3.50;2.91] | −0.43 [−2.08;1.21] | −0.91 [−3.60;1.79] | 0.943 |
| Other outcomes | Global <i>n</i> = 402 | Phenotype 1 <i>n</i> = 138 | Phenotype 2 <i>n</i> = 148 | Phenotype 3 <i>n</i> = 116 | <i>p</i> value |
| <i>Clinical improvement measures</i> | | | | | |
| Sputum amount | 252 (83.2%) | 94 (84.7%) | 97 (87.4%) | 61 (75.3%) | 0.075 |
| Sputum purulence | 242 (81.2%) | 93 (86.1%) | 93 (84.5%) | 56 (70.0%) | 0.011 |
| Dyspnea | 185 (61.7%) | 78 (70.3%) | 77 (70.0%) | 30 (38.0%) | <0.001 |
| <i>Adverse effects</i> | | | | | |
| None | 289 (74.3%) | 105 (81.4%) | 116 (78.9%) | 68 (60.2%) | <0.001 |
| Cough | 47 (12.1%) | 11 (8.53%) | 14 (9.52%) | 22 (19.5%) | 0.016 |
| Aphonia | 6 (1.54%) | 2 (1.55%) | 2 (1.36%) | 2 (1.77%) | 1.000 |
| Acute respiratory failure | 1 (0.26%) | 0 (0.00%) | 0 (0.00%) | 1 (0.88%) | 0.290 |
| Acute renal failure | 1 (0.26%) | 1 (0.78%) | 0 (0.00%) | 0 (0.00%) | 0.622 |
| Digestive | 3 (0.77%) | 1 (0.78%) | 0 (0.00%) | 2 (1.77%) | 0.197 |
| Cutaneous | 5 (1.29%) | 1 (0.78%) | 4 (2.72%) | 0 (0.00%) | 0.197 |
| <i>IA treatment suspension</i> | | | | | |
| Intolerance | 84 (21.2%) | 19 (14.1%) | 27 (18.5%) | 38 (33.0%) | 0.001 |
| Eradication | 44 (11.1%) | 18 (13.3%) | 15 (10.3%) | 11 (9.57%) | 0.590 |
| Stability | 65 (16.2%) | 24 (17.4%) | 24 (16.2%) | 17 (14.7%) | 0.840 |
| Dose reduction | 16 (4.23%) | 6 (4.62%) | 5 (3.60%) | 5 (4.59%) | 0.901 |

Data are presented *n* (%) or mean [95%CI]. Phenotype 1: normal lung function; Phenotype 2: moderately impaired lung function; Phenotype 3: severely impaired lung function. FEV₁: forced expiratory volume in 1 s; FVC: forced vital capacity; DLCO: diffusing capacity of the lungs for carbon monoxide.

nomogram (Fig. 1 Panel D) were developed to estimate individual risk.

This multicenter, retrospective study identified three clinically relevant phenotypes among patients with chronic bronchial infection (CBI) receiving inhaled antibiotic (IA) therapy, based primarily on lung function status. Phenotype 1 included patients with preserved lung function, mainly women with bronchiectasis; phenotype 2 had moderate obstruction, often with overlapping COPD and bronchiectasis; and phenotype 3 comprised older patients with severe airflow limitation, predominantly older patients with COPD and multiple comorbidities.

While all phenotypes benefited from IA therapy—showing reductions in exacerbations and symptomatic improvement—patients in phenotype 3 experienced poorer outcomes. This group had the highest exacerbation rate after one year, the lowest symptom response, and a greater incidence of adverse effects, leading to treatment discontinuation in one-third of cases. In contrast, phenotypes 1 and 2 showed more favorable responses, with slightly greater reductions in exacerbation frequency compared to prior studies, likely due to the exclusive inclusion of patients with confirmed CBI in this cohort.

The study's key innovation is the development of a multivariable model to predict exacerbation risk one year after IA initiation. The model includes age, comorbidities, prior exacerbations, bronchiectasis type on chest CT (varicose/cystic), and phenotype classification. It showed moderate discriminative capacity (AUC 0.74) and is accessible via a web-based tool and nomogram, facilitating its use in clinical practice. Unlike previous scoring systems (e.g., FACED,¹⁴ E-FACED¹⁵), this model incorporates bronchiecta-

sis type, which has rarely been linked to outcomes but may be a marker of more advanced disease. The inclusion of CT morphology may thus represent a step toward precision medicine in CBI.

The poor response observed in phenotype 3 highlights a clinically relevant subgroup. These patients likely derive limited benefit from IA due to non-modifiable risk factors such as advanced COPD,¹⁶ comorbidities, and age. Importantly, viral infections¹⁷—common drivers of exacerbations in severe COPD—are unaffected by IA, which may explain the limited efficacy in this group. While adverse effects were more frequent in phenotype 3, the rate was comparable to previous studies of IA in COPD, though those cohorts included fewer CBI patients and had higher baseline lung function.^{18,19}

Despite its retrospective design and reliance on electronic medical records, the study includes data from 10 Spanish hospitals, reflecting real-world practice and enhancing generalizability. Limitations include the absence of a control group, incomplete availability of inflammatory markers, and some variability across centers. A prospective multicenter study (REPAIR²⁰) is underway to validate these findings and further assess the utility of the proposed model.

In conclusion, phenotypic analysis in this large cohort of CBI patients treated with IA revealed distinct clinical profiles associated with differential response to therapy. The predictive model developed in this study, incorporating both clinical variables and radiological features, may guide individualized treatment decisions and inform future trial design. While further prospective validation is needed, these findings offer a practical framework for optimizing IA use in patients with CBI.




Fig. 1. Panel A: Distribution of clustered characteristics by phenotype; Panel B: Exacerbation one-year pre-post IA according to phenotypes; Panels C and D: Multivariate model for risk of exacerbation one-year after IA. (A) Radar plot showing the prevalence or standardized mean (0–1) of clustered characteristics for clinical phenotype identification. Clustering was performed using k-prototypes with multiple imputation on sociodemographic information (blue), underlying respiratory diseases (purple), comorbidities (yellow), pulmonary function (green) and pharmacological treatments (red). *Abbreviations:* COPD, chronic obstructive pulmonary disease; FVC, forced vital capacity; FEV₁, forced expiratory volume in 1 second; ICS, inhaled corticosteroids; LABA, long-acting beta2 agonist; LAMA, long-acting muscarinic antagonist. (C) Forest plot with Odds Ratios and confidence interval for predictors of exacerbation model. (D) Nomogram to predict risk of exacerbation after one-year IA.

Authors' Contributions

Conceptualization (BR, IDB, JG, AS), data curation (IDB, AS, AM, IJ-G), formal analysis (IDB, AS, AM), investigation (all), methodology (BR, IDB, JG, AS, DR, GS), project administration (JG, AS, DR, GS), supervision (JG, AS, AM), writing – original draft (BR, JG, IDB, AS), and writing – review & editing (all). All authors provided final approval of the version submitted for publication.

Artificial Intelligence Involvement

The authors declare that no material has been partially or totally produced with the help of artificial intelligence.

Sources of Support

This study has been founded by Instituto de Salud Carlos III (ISCIII) through the project PI23/01381 and co-funded by the European Union. AS was supported by Departament de Salut (Pla Estratègic de Recerca i Innovació en Salut (PERIS): SLT028/23/000191), SS was supported by Departament de Salut (Pla Estratègic de Recerca i Innovació en Salut (PERIS): SLT035/24/000025) and IJ-G was supported by the Instituto de Salud Carlos III (ISCIII) through the predoctoral fellowship FI24/00084 and co-funded by the European Union.

Conflicts of Interest

The authors declare not to have any conflicts of interest that may be considered to influence directly or indirectly the content of the manuscript.

References

- Solarat B, Perea L, Faner R, de La Rosa D, Martínez-García MÁ, Sibila O. Pathophysiology of chronic bronchial infection in bronchiectasis. *Arch Bronconeumol.* 2023;59(2):101–8. <http://dx.doi.org/10.1016/j.arbres.2022.09.004>. English, Spanish.
- Chalmers JD, Aliberti S, Blasi F. Management of bronchiectasis in adults. *Eur Respir J.* 2015;45(5):1446–62. <http://dx.doi.org/10.1183/09031936.00119114>.
- Aliberti S, Lonni S, Dore S, McDonnell MJ, Goeminne PC, Dimakou K, et al. Clinical phenotypes in adult patients with bronchiectasis. *Eur Respir J.* 2016;47(4):1113–22. <http://dx.doi.org/10.1183/13993003.01899-2015>.
- Tay GT, Reid DW, Bell SC. Inhaled antibiotics in cystic fibrosis (CF) and non-CF bronchiectasis. *Semin Respir Crit Care Med.* 2015;36(2):267–86. <http://dx.doi.org/10.1055/s-0035-1547346>.
- Laska IF, Crichton ML, Shoemark A, Chalmers JD. The efficacy and safety of inhaled antibiotics for the treatment of bronchiectasis in adults: a systematic review and meta-analysis. *Lancet Respir Med.* 2019;7(10):855–69. [http://dx.doi.org/10.1016/S2213-2600\(19\)30185-7](http://dx.doi.org/10.1016/S2213-2600(19)30185-7).
- Cordeiro R, Choi H, Haworth CS, Chalmers JD. The efficacy and safety of inhaled antibiotics for the treatment of bronchiectasis in adults: updated systematic review and meta-analysis. *Chest.* 2024;166(1):61–80. <http://dx.doi.org/10.1016/j.chest.2024.01.045>.
- Polverino E, Goeminne PC, McDonnell MJ, Aliberti S, Marshall SE, Loeblinger MR, et al. European Respiratory Society guidelines for the management of adult bronchiectasis. *Eur Respir J.* 2017;50(3):1700629. <http://dx.doi.org/10.1183/13993003.00629-2017>.

8. Martínez-García MÁ, Máiz L, Oliveira C, Girón RM, de la Rosa D, Blanco M, et al. Spanish guidelines on treatment of bronchiectasis in adults. *Arch Bronconeumol*. 2018;54(2):88–98. <http://dx.doi.org/10.1016/j.arbr.2017.07.014>.
9. Hill AT, Sullivan AL, Chalmers JD, De Soya A, Elborn JS, Floto RA, et al. British Thoracic Society guideline for bronchiectasis in adults. *BMJ Open Resp Res*. 2018;5(1):e000348. <http://dx.doi.org/10.1136/bmjresp-2018-000348>.
10. Sibila O, Laserna E, Shoemark A, Perea L, Bilton D, Crichton ML, et al. Heterogeneity of treatment response in bronchiectasis clinical trials. *Eur Respir J*. 2022;59(5):2100777. <http://dx.doi.org/10.1183/13993003.00777-2021>.
11. Chalmers JD, Chotirmall SH. Bronchiectasis: new therapies and new perspectives. *Lancet Respir Med*. 2018;6:715–26. [http://dx.doi.org/10.1016/S2213-2600\(18\)30053-5](http://dx.doi.org/10.1016/S2213-2600(18)30053-5).
12. de la Rosa Carrillo D, López-Campos JL, Alcázar Navarrete B, Calle Rubio M, Cantón Moreno R, García-Rivero JL, et al. Documento de consenso sobre el diagnóstico y tratamiento de la infección bronquial crónica en la enfermedad pulmonar obstructiva crónica. *Arch Bronconeumol*. 2020;56:651–64. <http://dx.doi.org/10.1016/j.arbres.2020.04.023>.
13. Aschenbruck R, Szepannek G, Wilhelm AFX. Imputation strategies for clustering mixed-type data with missing values. *J Classif*. 2023;40:2–24. <http://dx.doi.org/10.1007/s00357-022-09422-y>.
14. Martínez-García MÁ, de Gracia J, Vendrell Relat M, Girón RM, Máiz Carro L, de la Rosa Carrillo D, et al. Multidimensional approach to non-cystic fibrosis bronchiectasis: the FACED score. *Eur Respir J*. 2014;43(5):1357–67. <http://dx.doi.org/10.1183/09031936.00026313>.
15. Martínez-García MA, Athanazio RA, Girón R, Máiz-Carro L, de la Rosa D, Oliveira C, et al. Predicting high risk of exacerbations in bronchiectasis: the E-FACED score. *Int J Chron Obstruct Pulmon Dis*. 2017;12:275–84. <http://dx.doi.org/10.2147/COPD.S121943>.
16. Dharmage SC, Bui DS, Walters EH, Lowe AJ, Thompson B, Bowatte G, et al. Life-time spirometry patterns of obstruction and restriction, and their risk factors and outcomes: a prospective cohort study. *Lancet Respir Med*. 2023;11(3):273–82. [http://dx.doi.org/10.1016/S2213-2600\(22\)00364-2](http://dx.doi.org/10.1016/S2213-2600(22)00364-2).
17. Venkatesan P. GOLD COPD report: 2024 update. *Lancet Respir Med*. 2024;12(1):15–6. [http://dx.doi.org/10.1016/S2213-2600\(23\)00461-7](http://dx.doi.org/10.1016/S2213-2600(23)00461-7).
18. Bruguera-Avila N, Marin A, Garcia-Olive I, Radua J, Prat C, Gil M, et al. Effectiveness of treatment with nebulized colistin in patients with COPD. *Int J Chron Obstruct Pulmon Dis*. 2017;12:2909–15. <http://dx.doi.org/10.2147/COPD.S138428>.
19. De la Rosa Carrillo D, Martínez-García MÁ, Barreiro E, Tabernero Huguet E, Costa Sola R, García-Clemente MM, et al. Effectiveness and safety of inhaled antibiotics in patients with chronic obstructive pulmonary disease. A multicentre observational study. *Arch Bronconeumol*. 2022;58(1):11–21. <http://dx.doi.org/10.1016/j.arbres.2021.03.009>. English, Spanish.
20. de la Rosa-Carrillo D, Miravittles M, Pomares X, Martínez-García MÁ, et al. Unraveling the enigma: Spain takes a pioneering step with REPAIR in the world of inhaled antibiotics. *Arch Bronconeumol*. 2024;60(5):261–2. <http://dx.doi.org/10.1016/j.arbres.2024.01.007>.

Beatriz Raboso Moreno^{a,1}, Iván D. Benítez^{b,c,d,1}, Marta Erro^e, Javier Burgos^f, Alicia Sayés^g, Eduardo J. García-Urrestarazu^g, Cristina Pou^h, Rosa Abril Castañónⁱ, Annie Navarro^j, Ane Martínez-De las Fuentes^f, Joel-Suresh Lakhani^k, Ivan Juez-García^b, Carlos Manzano^b, Sally Santistevé^{b,c}, Arturo Morales^l, Aida Monge^b, Anna Moncusí-Moix^{b,c}, Anna Sánchez-Cucó^b, Ana Belén Alcaide^g, Juan P. de-Torres^{c,h}, Ferran Barbé^{b,c}, Guillermo Suarez-Cuartín^{c,f}, David de la Rosa^{c,k}, Jessica González^{b,c,*}

^a Universidad Europea de Madrid, Faculty of Medicine, Health and Sports, Department of Medicine, Department of Pulmonology, Getafe University Hospital, Madrid, Spain

^b Translational Research in Respiratory Medicine, University Hospital Arnau de Vilanova and Santa Maria, IRBLleida, Lleida, Spain

^c CIBER of Respiratory Diseases (CIBERES), Institute of Health Carlos III, Madrid, Spain

^d Department of Basic Medical Sciences, Lleida Biomedical Research Institute (IRBLleida), University of Lleida, Lleida, Spain

^e Puerta de Hierro University Hospital, Madrid, Spain

^f Bellvitge University Hospital, Bellvitge Biomedical Research Institute (IDIBELL), Universitat de Barcelona, Barcelona, Spain

^g Navarra University Clinic, Pamplona, Spain

^h Research Group NeumoVigo I+i, Instituto de Investigación Sanitaria Galicia Sur (IISGS), Álvaro Cunqueiro Hospital, Vigo, Spain

ⁱ Insular Maternal and Child University Hospital of Gran Canaria, Gran Canaria, Spain

^j Fundació Hospital Sant Joan de Deu de Martorell, Martorell, Spain

^k Hospital de la Santa Creu i Sant Pau, Barcelona, Spain

^l Pharmacy Department, University Hospital Arnau de Vilanova and Santa Maria, Spain

*Corresponding author.

E-mail address: jgonzalezgutierrez88@gmail.com (J. González).

¹ Beatriz Raboso and Iván D. Benítez contributed equally to the manuscript and are co-first authors.