

## Original Article

## Hospital Mortality and Effect of Adjusting PaO<sub>2</sub>/FiO<sub>2</sub> According to Altitude Above the Sea Level in Acclimatized Patients Undergoing Invasive Mechanical Ventilation. A Multicenter Study

Manuel Jibaja<sup>a,b,\*</sup>, Guillermo Ortiz-Ruiz<sup>c,d</sup>, Fernanda García<sup>e</sup>, Manuel Garay-Fernández<sup>f,c</sup>, Felipe de Jesús Montelongo<sup>g</sup>, Jorge Martínez<sup>h</sup>, José Antonio Viruez<sup>i</sup>, Orville Baez-Pravia<sup>j</sup>, Santiago Salazar<sup>k</sup>, Francisco Villacorta-Cordova<sup>l</sup>, Freddy Morales<sup>m,n</sup>, Amilcar Tinoco-Solórzano<sup>o,p</sup>, Carlos Ibañez Guzmán<sup>q</sup>, Bruno Valle Pinheiro<sup>r,s</sup>, Felix Zubia-Olaskoaga<sup>t,u</sup>, Carmelo Dueñas<sup>v,w</sup>, Antonio Lara García<sup>c,f</sup>, Pablo Cardinal-Fernández<sup>x,y</sup>

<sup>a</sup> Intensive Care Unit, Hospital Eugenio Espejo, Quito, Ecuador

<sup>b</sup> Escuela de Medicina, Universidad Internacional del Ecuador, Quito, Ecuador

<sup>c</sup> Universidad El Bosque, Bogotá, Colombia

<sup>d</sup> Universidad de Barcelona, Spain

<sup>e</sup> Intensive Care Unit, Hospital Eugenio Espejo, Quito, Ecuador

<sup>f</sup> Intensive Care Unit, Hospital Santa Clara Bogotá, Colombia

<sup>g</sup> Intensive Care Unit, Hospital General de las Américas, Ecatepec, Mexico

<sup>h</sup> Intensive Care Unit, Hospital San Pedro, Pasto, Colombia

<sup>i</sup> Intensive Care Unit, Hospital Norte, El Alto, Bolivia

<sup>j</sup> Intensive Care Unit, Hospital Universitario HM Sanchinarro, Madrid, Spain

<sup>k</sup> Intensive Care Unit, Hospital Carlos Andrade Marín, Quito, Ecuador

<sup>l</sup> Intensive Care Unit, Hospital General Docente, Riobamba, Ecuador

<sup>m</sup> Intensive Care Unit, Hospital Oncológico Dr. Julio Villacreses Comont, Portoviejo, Ecuador

<sup>n</sup> Intensive Care Unit, Clínica Santa Margarita, Portoviejo, Ecuador

<sup>o</sup> Intensive Care Unit, Hospital Ramiro Prialé Prialé EsSalud, Huancayo, Peru

<sup>p</sup> Centro de Investigación de Medicina de la altura, Facultad de Medicina Humana, Universidad de San Martín de Porres, Lima, Peru

<sup>q</sup> Intensive Care Unit, Hospital Obrero N° 1, La Paz, Bolivia

<sup>r</sup> Pulmonary Research Laboratory, Federal University of Juiz de Fora, Minas Gerais, Brazil

<sup>s</sup> Intensive Care Unit, Hospital Universitário, Universidade Federal de Juiz de Fora, Minas Gerais, Brazil

<sup>t</sup> Intensive Care Department, Donostia University Hospital, San Sebastián, Spain

<sup>u</sup> Departamento de Medicina, Universidad del País Vasco-Euskal Herriko Unibertsitatea, Spain

<sup>v</sup> Intensive Care Department, Gestión Salud, Cartagena, Colombia

<sup>w</sup> Universidad de Cartagena, Cartagena, Colombia

<sup>x</sup> Hospital Universitario HM Sanchinarro, Madrid, Spain

<sup>y</sup> HM Research Foundation, Madrid, Spain

## ARTICLE INFO

## Article history:

Received 26 February 2019

Accepted 28 June 2019

Available online 30 September 2019

## Keywords:

Altitude  
High altitude  
Mortality  
Respiratory failure  
Mechanical ventilation

## ABSTRACT

**Objective:** (i) Analyze the effect of altitude above the sea level on the mortality rate in patients undergoing invasive mechanical ventilation. (ii) Validate the traditional equation for adjusting PaO<sub>2</sub>/FiO<sub>2</sub> according to the altitude.

**Design:** A prospective, observational, multicenter and international study conducted during August 2016.

**Patients:** Inclusion criteria: (i) age between 18 and 90 years old, (ii) admitted to intensive care unit (ICU) situated at the same altitude above the sea level (AASL) in which the patients has stayed, at least, during the previous 40 days and (iii) received invasive MV for at least 12 h.

**Material and methods:** All variables were registered the day of intubation (day 0). Patients were followed until death, ICU discharge or day 28. PaO<sub>2</sub>/FiO<sub>2</sub> ratio was adjusted by the AASL according to: PaO<sub>2</sub>/FiO<sub>2</sub> \* (barometric pressure/760). Categorical variables were compared with  $\chi^2$  and Cochran–Mantel–Haenszel test. Continuous variables with Mann–Whitney. Correlation between continuous variables was analyzed graphically and analytically. Logistic regression model was constructed to identify factors associated to mortality. Kaplan–Meier method was used to estimate the probability of survival according to the altitude. A 2-side *p* value <0.05 was consider significant.

\* Corresponding author.

E-mail address: [mjibaja79@gmail.com](mailto:mjibaja79@gmail.com) (M. Jibaja).

**Results:** 249 patients (<1500 m n = 55; 1500 to <2500 m n = 20; 2500 to <3500 m n = 155 and ≥3500 m n = 19) were included. Adjusted and non-adjusted PaO<sub>2</sub>/FiO<sub>2</sub> were correlated with several respiratory and non respiratory variables. None discordances between non adjusted and adjusted PaO<sub>2</sub>/FiO<sub>2</sub> were identified. However, several correlations were appreciated only in patients situated <1500 m or in >1500 m. Seventy-nine patients died during the ICU stayed (32%). The mortality curve was not affected by the altitude above the sea level. Variables independently associated to mortality are: PEEP, age, systolic arterial blood pressure, and platelet count. AUROC: 0.72.

**Conclusion:** In acclimatized patients undergoing invasive mechanical ventilation, the traditional equation for adjusting PaO<sub>2</sub>/FiO<sub>2</sub> according the elevation above the sea level seems to be inaccurate and the altitude above the sea level does not affect the mortality risk.

© 2019 SEPAR. Published by Elsevier España, S.L.U. All rights reserved.

## Mortalidad hospitalaria y efecto de ajustar el cociente PaO<sub>2</sub>/FiO<sub>2</sub> de acuerdo con la altitud por encima del nivel del mar en pacientes aclimatados que se someten a ventilación mecánica invasiva. Un estudio multicéntrico

### R E S U M E N

#### Palabras clave:

Altitud  
Altitud elevada  
Mortalidad  
Fallo respiratorio  
Ventilación mecánica

**Objetivo:** 1) Analizar el efecto de la altitud por encima del nivel del mar en la tasa de mortalidad de pacientes sometidos a ventilación mecánica invasiva, y 2) Validar la ecuación tradicional de ajuste de PaO<sub>2</sub>/FiO<sub>2</sub>, de acuerdo con la altitud.

**Diseño:** Estudio internacional prospectivo, observacional y multicéntrico realizado durante agosto de 2016.

**Pacientes:** Criterios de inclusión: 1 Edad comprendida entre 18 y 90 años, 2 Haber sido ingresado en una unidad de cuidados intensivos (UCI) situada a la misma altitud por encima del nivel del mar (AASL) en la cual el paciente haya estado durante al menos los 40 días previos al estudio, y 3) Haber recibido ventilación mecánica (VM) durante al menos 12 h.

**Materiales y métodos:** Todas las variables se registraron el día de la intubación (día 0). El seguimiento se realizó hasta la muerte del paciente, el alta de la UCI o el día 28. El cociente PaO<sub>2</sub>/FiO<sub>2</sub> se ajustó según los criterios de la AASL de acuerdo con: PaO<sub>2</sub>/FiO<sub>2</sub> \* (presión barométrica/760). Las variables categóricas se compararon mediante la prueba de  $\chi^2$  y el test Cochran-Mantel-Haenszel, y las variables continuas con el test de Mann-Whitney. La correlación entre las variables continuas se analizó de forma gráfica y analítica. Para identificar los factores asociados a la mortalidad se elaboró un modelo de regresión logística. Se utilizó el método de Kaplan-Meier para estimar la probabilidad de supervivencia de acuerdo con la altitud. Un valor de p < 0,05 en la prueba bilateral se consideró como significativo.

**Resultados:** Se incluyeron 249 pacientes (<1.500 m, n = 55; 1.500 a <2.500 m, n = 20; 2.500 a <3.500 m, n = 155 y ≥3.500 m, n = 19). El cociente PaO<sub>2</sub>/FiO<sub>2</sub> mostró correlación con las variables graves tanto respiratorias como no respiratorias. No se registraron discordancias entre el cociente PaO<sub>2</sub>/FiO<sub>2</sub> ajustado y sin ajustar. Únicamente se observaron diversas correlaciones entre los pacientes situados a <1.500 m o a >1.500 m. Setenta y nueve pacientes (32%) murieron durante la estancia en la UCI. La altitud sobre el nivel del mar no afectó a la curva de mortalidad. Las variables asociadas de forma independiente con la mortalidad fueron la presión positiva al final de la espiración (PEEP), la edad, la presión arterial sistólica y el recuento de plaquetas. El área bajo la curva ROC (AUROC) fue de 0,72.

**Conclusión:** En pacientes aclimatados sometidos a ventilación mecánica invasiva la ecuación tradicional para ajustar el cociente PaO<sub>2</sub>/FiO<sub>2</sub>, de acuerdo con la elevación sobre el nivel del mar parece inexacta. Por otro lado, la altitud por encima del nivel del mar no afecta al riesgo de mortalidad.

© 2019 SEPAR. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

## Introduction

Geographical areas above the sea level are associated to adverse conditions for the Life. For example, although the oxygen concentration remains constant with the trophosphere, the barometric pressure falls, which determines a condition called hypobaric hypoxemia. Indeed, the humidity and temperature, factors that may contribute to airway reactivity, insensible water losses, ventilatory changes and alterations in pulmonary hemodynamics,<sup>1</sup> also decrease in these areas. Fast ascend to elevated areas is associated to well know entities such us mountain sickness, High Altitude Pulmonary Edema (HAPE), High Altitude Cerebral Edema (HACE) and cardiovascular collapse.<sup>2</sup> However, those who stay for a long period of time above the sea level develop mechanisms to compensate this adverse environment; therefore previously mentioned entities are unusual. Acclimatization to altitude is a term use to describe physiological changes that occur over several weeks with the aim

to increase the endurance against the deleterious effect of staying above the sea level.<sup>3–5</sup>

Invasive mechanical ventilation is the cornerstone intervention for supporting severe respiratory failure. Unfortunately, despite 385 million people permanently residing at elevations above 1500 m<sup>6</sup> and adverse effects of chronic exposure to high altitude are well demonstrated<sup>3,5,7</sup>; the knowledge regarding the effect of altitude above the sea level in acclimatized patients under invasive mechanical ventilation is scarce. Furthermore, several worldwide surveys regarding mechanical ventilation have been performed,<sup>8–11</sup> but none of them have considered the effect of altitude above the sea level. It is paramount importance to highlight that mechanical ventilators are non pressurized systems and they function at the same atmospheric pressure as the environment. This mean that, despite the patient is under invasive mechanical ventilation, their alveolar pressure and their arterial pressure is lower than at sea level for the same inspiratory pressure.

On the other hand, with the aim to maintain the biological significance of one variable (e.g. diagnostic criteria for a disease or prognosis factor) sometimes it is necessary to adjust their value according to another variable. For example, it is possible to use the partial oxygen pressure ( $\text{PaO}_2$ ) as a surrogate biomarker of the gas exchange if everybody receives the same oxygen inspiratory fraction ( $\text{FiO}_2$ ). However, as it is possible to prescribe different  $\text{FiO}_2$ , it is common to relativize the  $\text{PaO}_2$  value according to the  $\text{FiO}_2$  ( $\text{PaO}_2/\text{FiO}_2$ ). Partial oxygen pressure is also dramatically influenced by the atmospheric pressure, thus in high altitudes above the sea level the biological significance of a specific  $\text{PaO}_2/\text{FiO}_2$  could differ than at sea level. A mathematical equation has been proposed to adjust the  $\text{PaO}_2/\text{FiO}_2$  according to the altitude ( $\text{PaO}_2/\text{FiO}_2 * [\text{barometric pressure}/760]$ ).<sup>12,13</sup> However, as far as we know, it is an empiric equation and none have validate it.

Studying the effect of the altitude above the sea level is not merely an academic exercise as, if the outcome of lowlanders differs from highlanders, it may imply that one group are exposed to different risks and require special interventions than the other. Indeed, mechanisms for adaptation to hypobaric hypoxia could be useful in other hypoxemic diseases such as chronic obstructive pulmonary diseases (COPD), myocardial ischemia or ARDS.<sup>3,12,14–16</sup> On the other hand, validating the equation for adjusting the  $\text{PaO}_2/\text{FiO}_2$  according to the altitude is also of paramount importance as is the variable used to define, stratify and select the best treatment for patients with ARDS.<sup>12,17–19</sup> Likewise, due to the fact that  $\text{PaO}_2/\text{FiO}_2$  is associated to the outcome in general population under invasive mechanical ventilation, predicting their outcome would be inaccurate in patients living in high altitudes.<sup>10,20</sup>

We hypothesize that altitude above the sea level affects the outcome of acclimatized patients undergoing invasively mechanically ventilated.

The aim of this study is (i) to validate the traditional equation for adjusting  $\text{PaO}_2/\text{FiO}_2$  according to the altitude and (ii) to analyze the effect of altitude above the sea level on the mortality rate in patients undergoing invasive mechanical ventilation.

## Material and method

**Design:** prospective, observational, multicenter and international study conducted during August 2016.

**Inclusion criteria:** age between 18 and 90 years old, admitted to intensive care unit (ICU) situated at the same altitude above the sea level in which the patient has stayed, at least, during the previous 40 days before ICU admission and received invasive MV for at least 12 h. The full list of centers can be appreciated in [supplementary material](#). Two hospitals (“Obrero N° 1” Hospital, Bolivia and Donostia University Hospital, Spain) retrieved the data retrospectively. **Exclusion criteria** include: home oxygen therapy, metastatic cancer, pregnancy, and absence of informed consent.

## Protocol

General and national coordinators recruited local researchers from eligible ICUs. With the aim of decreasing the bias in practice, only research coordinators directly related to the study were aware of the exact purpose, the timing of the study and variables reported.

A website was developed to register demographic characteristics (age, weight and height), comorbidities (COPD, diabetes, chronic heart failure, chronic renal failure, active solid cancer and active hemato-oncologic cancer), clinical data (arterial systolic pressure and APACHE II score), ventilator parameters (tidal volume, peak pressure, plateau pressure, positive end expiratory and respiratory rate) and analytical (arterial blood gases, hemoglobin, leukocytes, platelets, C reactive protein, creatinine

and total bilirubin) of patients on the day of starting invasive mechanical ventilation.

The  $\text{PaO}_2/\text{FiO}_2$  was recorded and then, in patients treated 1000 m above the sea level, adjusted according to following equation:  $[\text{PaO}_2/\text{FiO}_2 * (\text{barometric pressure}/760)]$ .<sup>12,13</sup> Thus, two  $\text{PaO}_2/\text{FiO}_2$  were contemplated: non-adjusted and adjusted.

According to Zubieta-Calleja et al.,<sup>21</sup> acclimation to a specific altitude was defined if the patient permanently stayed during 40 days at the same altitude above sea level. Altitude was expressed in meters above the sea level. Severe Sepsis and septic shock were defined according to the American College of Chest Physicians/Society of Critical Care Medicine<sup>22</sup> as it was the current definition at the time that the study was performed. Active haemato-oncologic or active solid cancer was defined if the patients were diagnosed with cancer (or metastasis), received chemotherapy, radiotherapy or biological therapy during the 60 days before the enrollment.

The study was registered at the Clinical trial website (NCT02871063), was approved by the ethics committee of the Universidad “El Bosque” – Bogota – Colombia and, when required, by the institutional review boards at the individual sites.

## Quality control

With the aim of decreasing errors during data entry on the website, a range of continuous variables and codification for qualitative variables was set up. Likewise, once the recruiting period finished, 10% of the patients were randomly selected for a second review by the general coordinators of the study (MJ, GO, PC). Incomplete records were not considered.

## Statistical analysis

All clinical and analytical variables correspond to the day of oral intubation and initiating mechanical ventilation. Continuous variables were expressed as the median with interquartile range and compared using Mann–Whitney test. Categorical variables were expressed as the absolute frequency with proportions and were compared using the  $\chi^2$  test. The strength of the association was expressed as the odds ratio (OR) and its 95% confidence interval (CI).

With the aim to validate the equation for adjusting the  $\text{PaO}_2/\text{FiO}_2$ , the correlation adjusted and non adjusted  $\text{PaO}_2/\text{FiO}_2$  and other continuous variables were compared in patients situated below and above 1500 m of altitude. Secondly, both  $\text{PaO}_2/\text{FiO}_2$  were stratified according to criteria used by Esteban et al.<sup>10</sup> in: <100; 100–149; 150–199; 200–300 and >300 and then the mortality rate distribution between <1500 m and  $\geq 1500$  m was compared using Cochran–Mantel–Haenszel test.

With the aim to identify the association between altitude and mortality, a logistic regression model was performed. The maximum model included all variables with a  $p$  value <0.1 in the univariate analysis. Then, one by one the variables with the highest  $p$  value were removed until all variables in the model had a  $p$  value <0.05 (backward step procedure). The AUROC, sensitivity, specificity, positive and negative likelihood ratio were calculated. The Kaplan–Meier method was used to estimate the probability of survival according to altitude categorized in <1500; 1500–2500; 2500–3500 and >3500.

A 2-sided  $p$  value less than 0.05 was considered statistically significant. All the analyses were performed using R package.<sup>23</sup>

## Results

Thirty medical–surgical ICUs with 249 patients from 7 countries (Bolivia, Brazil, Colombia, Ecuador, Mexico, Peru and Spain) were

**Table 1**  
Univariate and multivariate analysis for mortality.

	All population (n = 249)	Mortality (n = 79)	Survivors (n = 170)	p value
<i>Altitude above sea level (meters)</i>	2600(2250,2800)	2600(2212,2850)	2600(2250,2800)	0.31
<i>Female</i>	102(0.41)	34(0.43)	68(0.40)	0.75
<i>Body mass index (kg/m<sup>2</sup>)</i>	25(23,28)	25(23,29)	25(23,28)	0.80
<i>Age (years old)</i>	60(43,72)	63(48,76)	57(38,70)	0.04
<i>APACHE II</i>	18(12,26)	23(17,31)	16(12,24)	0.00
<i>Length of mechanical ventilation (days)</i>	4(1,8)	5(2,9)	4(1,7)	0.13
<i>Length of ICU stay (days)</i>	8(4,14)	7(3,13)	8(5,14)	0.06
<i>Length of hospital stay (days)</i>	18(9,30)	10(5,20)	21(13,35)	0.00
<i>Comorbidities</i>				
Chronic obstructive pulmonary disease	52(0.21)	22(0.28)	30(0.18)	0.09
Diabetes	41(0.16)	16(0.20)	25(0.15)	0.36
Chronic heart failure	48(0.19)	10(0.13)	38(0.22)	0.10
Chronic renal failure	40(0.16)	19(0.24)	21(0.12)	0.03
Current solid cancer	21(0.08)	10(0.13)	11(0.06)	0.16
Current hemato-oncologic cancer	10(0.04)	6(0.08)	4(0.02)	0.08
<i>Respiratory variables</i>				
Tidal volume (ml)	480(420,500)	480(407,515)	480(420,500)	0.93
Tidal volume per kilogram (ml/kg)	7 ± 2	7 ± 2	7 ± 2	0.89
Peak pressure (mmHg)	22(19,27)	24(20,29)	22(19,25)	0.03
Plateau pressure (mmHg)	18(14,22)	18(14,22)	18.5(14,22)	0.92
PEEP (mmHg)	6(5,8)	7(5,8)	5(5,8)	0.00
Respiratory rate (rpm)	18(15,20)	18(15,21)	18(15,20)	0.39
PsO <sub>2</sub> /FiO <sub>2</sub> non-adjusted	154(100,216.67)	122(80,194)	158(115,229)	0.00
PsO <sub>2</sub> /FiO <sub>2</sub> adjusted	112(74,175)	91(58,159)	122(84,179)	0.00
PaCO <sub>2</sub> (mmHg)	34(29,42)	34(29,43.5)	34(30,42)	0.99
<i>Systemic variables</i>				
Arterial systolic pressure (mmHg)	115(100,128)	107(95,120)	120(107,130)	0.00
Arterial pH	7.35(7.29,7.42)	7.33(7.27,7.41)	7.36(7.30,7.43)	0.12
Hemoglobin (g/dl)	11.2(9.6,13.0)	11.0(9.5,12.9)	11.3(9.7,13.2)	0.15
Platelets (cells 10 <sup>3</sup> /mml)	204(132,299)	180(118,267)	218(140,303)	0.02
Leukocytes (cells 10 <sup>3</sup> /mml)	12.4(8.9,17.0)	12.5(7.9,16.9)	12.1(9.1,17.1)	0.57
PCR	17(6,79)	18(8,48)	17(6,80)	0.80
Creatinine (mg/dl)	1.1(0.8,2.1)	1.4(0.9,2.5)	1(0.7,1.6)	0.00
Bilirubin (gr/dl)	1.0(0.6,2.0)	1(0.7,2.2)	1.0(0.6,1.7)	0.57

analyzed (Table S1). Patients were assisted at the following altitudes: <1500 m n = 55; 1500–2500 m n = 20; 2500–3500 m n = 155 and >3500 m n = 19. Reason for ICU admission was: pulmonary sepsis n = 111; extrapulmonary sepsis n = 42; trauma n = 16; acute respiratory failure n = 14; stroke n = 8; post-operative n = 6; acute heart failure n = 4; burned n = 3, cardiac arrest n = 3, hemorrhagic shock n = 3, pancreatitis n = 3 and miscellaneous n = 36. Characteristics of all patients can be appreciated in Table 1. Comparison between highlanders and lowlanders can be appreciated in supplementary Table 1.

#### Adjusting PaO<sub>2</sub>/FiO<sub>2</sub> according to the altitude

The correlation between all continuous variables, including PaO<sub>2</sub>/FiO<sub>2</sub> (non adjusted) and PaO<sub>2</sub>/FiO<sub>2</sub> (adjusted), is shown in Fig. 1 and supplementary Fig. 1. Adjusted and non-adjusted PaO<sub>2</sub>/FiO<sub>2</sub> were correlated with PEEP, arterial systolic blood pressure and creatinine concentration below and above 1500 m. However, respiratory rate, peak pressure, plateau pressure, PaCO<sub>2</sub>, arterial pH, tidal volume per kilogram, length of mechanical ventilation, body mass index and bilirubin concentration were correlated with both PaO<sub>2</sub>/FiO<sub>2</sub> only in high altitude patients. PCR concentration and APACHE II score were correlated with both PaO<sub>2</sub>/FiO<sub>2</sub> at sea level but not in altitude patients. None discordances between non adjusted and adjusted PaO<sub>2</sub>/FiO<sub>2</sub> were identified.

#### Mortality

Seventy-nine patients died during the ICU stayed (32%). The mortality curve was not affected by the altitude above the

sea level (Fig. 2, log-rank 0.986). However, when the mortality is stratified according to the PaO<sub>2</sub>/FiO<sub>2</sub> (non adjusted and adjusted) and the altitude dichotomized in the two main groups (<1500 m and ≥1500 m), the observer mortality in each PaO<sub>2</sub>/FiO<sub>2</sub> range was different between both altitudes (Table 2).

The univariate analysis demonstrated that age, haemato-oncology disease, COPD, chronic renal failure, creatinine, platelets count, arterial systolic blood pressure, APACHE II, PaO<sub>2</sub>/FiO<sub>2</sub> (non-adjusted), PaO<sub>2</sub>/FiO<sub>2</sub> (adjusted), peak pressure and PEEP were associated to mortality (Table 1). All these variables except PaO<sub>2</sub>/FiO<sub>2</sub> (non-adjusted) and APACHE II were included in the maximum logistic regression model. Finally, PEEP (OR 1.17; 95% CI 1.06; 1.31; p: <0.01), age (OR 1.02; 95% CI 1.01; 1.04; p: <0.01), systolic arterial blood pressure (OR 0.98; 95% CI 0.96; 0.99; p: <0.01) and platelet count (OR 0.9999974; 95% IC 0.9999949; 0.9999997, p: 0.03) were independently associated to mortality (Table 1). The AUROC of the logistic model was 0.72 (95% CI: 0.65; 0.79), the sensibility 0.27 (0.17; 0.38), specificity 0.91 (0.85; 0.95), positive likelihood ratio 2.82 (1.56; 5.11) and negative likelihood ratio 0.81 (0.70; 0.93) (Fig. 3). The final regression model did no change when PaO<sub>2</sub>/FiO<sub>2</sub> (non-adjusted) was substituted by PaO<sub>2</sub>/FiO<sub>2</sub> (adjusted).

#### Discussion

Main results of this pioneer study are: (i) the recommended formula for adjusting the PaO<sub>2</sub>/FiO<sub>2</sub> ratio according to the altitude above the sea level seems to be inaccurate and (ii) the altitude above the sea level not influence the mortality risk.

According to our results, in term of mortality prediction or correlation with physiological variables, the traditional equation for adjusting the PaO<sub>2</sub>/FiO<sub>2</sub> according to the altitude above the sea level




Fig. 1. Correlation between continuous variables at <1500 m and >=1500 m.




Fig. 2. Survival analysis according to the altitude above the sea level.

Table 2  
Cochran–Mantel–Haenszel test (PaO<sub>2</sub>/FiO<sub>2</sub> and mortality stratified according to altitude above sea level).

PaO <sub>2</sub> /FiO <sub>2</sub>	All patients	≥1500 m		<1500 m	
		Survivors	Non survivors	Survivors	Non survivors
<b>(a) Not adjusted</b>					
<100	59	30	24	2	3
100–149	59	31	23	5	0
150–199	52	36	7	6	3
200–300	53	30	6	8	9
>300	26	6	1	16	3
<b>(b) Adjusted</b>					
<100	100	52	43	2	3
100–149	66	48	12	6	0
150–199	40	22	5	7	6
200–300	27	11	0	10	6
>300	16	0	1	12	3


(a) Not adjusted PaO<sub>2</sub>/FiO<sub>2</sub>  
Cochran–Mantel–Haenszel M<sup>2</sup> = 16.428, df = 4, p-value = 0.002495.

(b) Adjusted PaO<sub>2</sub>/FiO<sub>2</sub>  
Cochran–Mantel–Haenszel M<sup>2</sup> = 19.097, df = 4, p-value = 0.0007523.

does not add any advantage over the raw variable. This fact was expected, as the equation that links both variables is lineal. However, it is important to highlight that some correlations identified in patients at <1500 m were not observed in patients at ≥1500 m and viceversa. Furthermore, the mortality rate, stratified according to PaO<sub>2</sub>/FiO<sub>2</sub>, also differ in both groups of patients. We hypothesize

that these finding evidence that it would be necessary to adjust the PaO<sub>2</sub>/FiO<sub>2</sub> according to the altitude; but the current equation is not accurate. Future studies with more patients and altitude levels should clarify this issue.

On the other hand, one of the hallmark studies performed in non-specific population under invasive mechanical ventilation was



**Fig. 3.** Receiver operating characteristic curve of mortality (logistic regression model).

conducted by Esteban et al.<sup>10</sup> It include 5183 patients from 361 ICU of Europe, North and Latin America and reported that several variable including  $\text{PaO}_2/\text{FiO}_2 < 200$  are independently associated to mortality.<sup>10</sup> Likewise, Sudarsanam et al.<sup>20</sup> analyzed 200 consecutive patients admitted to an medical UCI and reported that type 1 respiratory failure is also independent predictor of mortality. On line with them, we reported that PEEP, age, systolic blood pressure and platelet count were associated to mortality risk. However, in our study neither in the regression model nor in the survival curve the mortality rate was affected by the altitude above the sea level. This result, that should be validated in future studies, rejected the hypothesis of this study and could reflect the efficacy of acclimatization mechanisms to altitude. Based on this result, we speculate that people acclimatized to high altitude have physiologic mechanisms to compensate the deleterious effect that high altitude bear. Considering this, one natural question arises: “At sea level, Is the outcome of patients adapted to high altitude similar to lowlanders?”. Exploring the differences between humans exposed and not exposed to hypobaric hypoxia has the potential to identify mechanisms important in critical illness and perhaps to alter our therapeutic focus toward increasing the efficiency of oxygen utilization rather than improving delivery.<sup>24</sup>

We have to accept that this study has some limitations. Firstly, the number of patients in extreme altitude is very small which may be explained by the low number of ICUs at those geographical areas. This may influence the result because this subgroup of patients and the clinical practice in these ICUs could differ from those at sea level. Likewise, it could be possible that specific subgroups of patients (e.g. ARDS, surgical, etc.) carried specific features that were not identify in this study. Secondly, despite enrolling a large number of ICUs from South America and Spain, our sample does not include all possible altitudes. Thirdly, the study was conducted for one month and may not represent year-round. Fourth, two centers collected the data retrospectively. Although this is an evident protocol deviation, it is very improbable that may influence the results as booth ICUs have an informatics medical records that allow them to include patients and retrieve their data. Fifth, the mechanical ventilation between different hospitals was not standardized. This limitation is common to all observational study. Sixty, recently new variables such as driving pressure and mechanical power have demonstrated a close relation with the outcome. Future studies will have to address the effect altitude above the sea level on these variable. Finally, as this is a pioneer study, the knowledge previously available to interpret

its result could be insufficient or inaccurate. Contrarily, this study has several strengths. Firstly, we used an objectively and previous published definition for considering a patient adapted to a specific altitude.<sup>21</sup> Unfortunately, analytical or genetic biomarkers for defining adaptation to a specific altitude are not available or are inaccurate. Secondly, the outcome we have address (mortality) is objective. Thirdly, we included several procedures (e.g., website, quality-control, etc.) to reduce the risk of bias. Likewise, the proportion of patient excluded from the final analysis, as their data were incomplete reflect the strict of methodology we applied.

## Conclusions

In acclimatized patients undergoing invasive mechanical ventilation, the traditional equation for adjusting  $\text{PaO}_2/\text{FiO}_2$  according the elevation above the sea level seems to be inaccurate and the altitude above the sea level does not affect the mortality risk.

## Funding

This study was partially supported by Escuela de Medicina de la Universidad Internacional del Ecuador (UIDE).

## Authors' contributions

PC, GO, MJ, participated in the design of the study and performed the statistical analysis. PC, GO, MJ, MG conceived of the study, and participated in its design and coordination and helped to draft the manuscript. PC, GO, MJ, MG, FG, FM, JM, JV, OB, SS, FV, FM, JT, CI BV, FZ, AL participate in coordination of their centers and critical care units. All authors read and approved the final manuscript.

## Conflict of interests

The authors declare no conflict of interest.

## Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at [doi:10.1016/j.arbres.2019.06.024](https://doi.org/10.1016/j.arbres.2019.06.024)

## References

- Grissom CK, Jones BE. Respiratory health benefits and risks of living at moderate altitude. *High Alt Med Biol.* 2017.
- Luks AM, Swenson ER, Bartsch P. Acute high-altitude sickness. *Eur Respir Rev.* 2017;26.
- Imray C, Booth A, Wright A, Bradwell A. Acute altitude illnesses. *BMJ.* 2011;343:d4943.
- Apte CR. Determinants of acclimatisation in high altitude. *Med J Armed Forces India.* 2012;68:197.
- Bartsch P, Gibbs JS. Effect of altitude on the heart and the lungs. *Circulation.* 2007;116:2191–202.
- Cohen JE, Small C. Hypsographic demography: the distribution of human population by altitude. *Proc Natl Acad Sci U S A.* 1998;95:14009–14.
- West JB. Physiological effects of chronic hypoxia. *N Engl J Med.* 2017;376:1965–71.
- Neto AS, Barbas CSV, Simonis FD, Artigas-Raventós A, Canet J, Determann RM, et al. Epidemiological characteristics, practice of ventilation, and clinical outcome in patients at risk of acute respiratory distress syndrome in intensive care units from 16 countries (PRoVENT): an international, multicentre, prospective study. *Lancet Respir Med.* 2016;4:882–93.
- Bellani G, Laffey JG, Pham T, et al. Epidemiology patterns of care, and mortality for patients with acute respiratory distress syndrome in intensive care units in 50 countries. *JAMA.* 2016;315:788–800.
- Esteban A, Anzueto A, Frutos F, Alía I, Brochard L, Stewart TE, et al. Characteristics and outcomes in adult patients receiving mechanical ventilation: a 28-day international study. *JAMA.* 2002;287:345–55.
- Esteban A, Frutos-Vivar F, Murriel A, Ferguson ND, Peñuelas O, Abaira V, et al. Evolution of mortality over time in patients receiving mechanical ventilation. *Am J Respir Crit Care Med.* 2013;188:220–30.

12. Ranieri VM, Rubenfeld GD, Thompson B, Ferguson ND, Caldwell E, Fan E, et al. Acute respiratory distress syndrome: the Berlin Definition. *JAMA*. 2012;307:2526–33.
13. West JB. Oxygen enrichment of room air to relieve the hypoxia of high altitude. *Respir Physiol*. 1995;99:225–32.
14. Cardinal-Fernandez P, Lorente JA, Ballen-Barragan A, Matute-Bello G. Acute respiratory distress syndrome and diffuse alveolar damage new insights on a complex relationship. *Ann Am Thorac Soc*. 2017;14:844–50.
15. Rios F, Iscar T, Cardinal-Fernandez P. What every intensivist should know about acute respiratory distress syndrome and diffuse alveolar damage. *Rev Bras Ter Intensiva*. 2017;29:354–63.
16. Avellanas Chavala ML. A journey between high altitude hypoxia and critical patient hypoxia: what can it teach us about compression and the management of critical disease? *Med Intensiva*. 2017.
17. Guerin C, Reigner J, Richard JC, Beuret P, Gacouin A, Boulain T, et al. Prone positioning in severe acute respiratory distress syndrome. *N Engl J Med*. 2013;368:2159–60.
18. Papazian L, Forel JM, Gacouin A, Penot-Ragon C, Perrin G, Loundou A, et al. Neuromuscular blockers in early acute respiratory distress syndrome. *N Engl J Med*. 2010;363:1107–16.
19. Schmidt GA, Girard TD, Kress JP, Morris PE, Ouellette DR, Alhazzani W, et al. Official executive summary of an American Thoracic Society/American College of Chest Physicians Clinical Practice Guideline: liberation from mechanical ventilation in critically ill adults. *Am J Respir Crit Care Med*. 2017;195:115–9.
20. Sudarsanam TD, Jeyaseelan L, Thomas K, John G. Predictors of mortality in mechanically ventilated patients. *Postgrad Med J*. 2005;81:780–3.
21. Zubieta-Calleja GR, Paulev PE, Zubieta-Calleja L, Zubieta-Castillo G. Altitude adaptation through hematocrit changes. *J Physiol Pharmacol*. 2007;58 Suppl. 5 (Pt 2):811–8.
22. Bone RC, Balk RA, Cerra FB, Dellinger RP, Fein AM, Knaus WA, et al. Definitions for sepsis and organ failure and guidelines for the use of innovative therapies in sepsis. The ACCP/SCCM Consensus Conference Committee. American College of Chest Physicians/Society of Critical Care Medicine. *Chest*. 1992;101:1644–55.
23. Zhang Z, Gayle AA, Wang J, Zhang H, Cardinal-Fernandez P. Comparing baseline characteristics between groups: an introduction to the CBCgrps package. *Ann Transl Med*. 2017;5:484.
24. Grocott M, Montgomery H, Vercueil A. High-altitude physiology and pathophysiology: implications and relevance for intensive care medicine. *Crit Care*. 2007;11:203.