



Spirometry Outside the Hospital

La espirometría en el medio extrahospitalario

Dear Editor,

We carefully read the editorial by Burgos Rincón et al., titled Impact of the COVID-19 Pandemic on Lung Function Laboratories: Considerations for “Today” and the “Day After”, recently published in Archivos de Bronconeumología.¹ The study considered important aspects, including the need to reorganize the lung function laboratories to recover the healthcare activity with sufficient security guarantees for both patients and professionals.¹ The study also stated that current COVID-19-related limitations warrant the need for respiratory teams to engage and explore innovative methods for lung function assessment.¹

Home-based spirometry, particularly, those with on-screen encouragement and/or remote guidance by a technician, could be an alternative to conventional spirometry for management and telemonitoring of chronic respiratory diseases. Home-based spirometry transmits the lung function data from patients’ home to a remote monitoring center (generally on a daily basis from previously trained patients) for a cautious evaluation by a respiratory function technician and interpretation by an expert physician, upholding the quality of the lung function assessment.^{2,3} In 1997, Brudeman et al. described a home-based portable spirometry system to detect early signs of asthma deterioration, which improved asthma management, quality of life, and reduced emergency department visits and hospitalizations.² For chronic obstructive pulmonary disease (COPD), Rodriguez-Roisin et al. compared the results of home-based spirometries performed by more than 2000 patients with severe COPD, after face-to-face training, and in clinical settings (using the same spirometry device); they found a good agreement between in-clinic and home-based measurements.³

In contrast, telespirometry conducted by respiratory function technicians in outpatient clinics might be an acceptable alternative after the resolution of regional COVID-19 outbreaks. Telespirometry uses a portable spirometer and transmits the lung function data online from a primary care clinic to a specialized center for interpretation. Consequently, some national networks were implemented, with positive outcomes.^{4–7} Results from the “RespiraNet Program” in Brazil and a Spanish study by Marina et al. showed that telespirometries performed by previously trained technicians, and analyzed and interpreted remotely by expert physicians, is a useful alternative to improve the diagnosis and management of chronic obstructive lung diseases.^{4,5} Furthermore, another Spanish study by Masa et al. concluded that telespirometry performed by a remotely located technician controlling the spirometer computer online enhanced the quality and the reliability of the maneuver.⁶ The “Alliance” study results showed that Italian general practitioners accepted telespirometry as an useful exploratory modality to improve the early diagnosis, monitoring, and management of COPD and asthma.⁷

Thus, home-based spirometry could decrease the burden on hospital services, avoid gathering of patients in high-risk facilities, and obviate the need for transports of patients to central areas during the COVID-19 pandemic. In addition, after the pandemic, telespirometry could be a key add-on to improve the response to future demand related to currently postponed laboratory lung function tests.

Financial Support

No funding was received for the publication of this article.

Conflict of Interest

None declared.

Bibliografía

1. Burgos Rincón F, Martínez Llorens J, Cordovilla Pérez R. Impact of the COVID-19 pandemic on lung function laboratories: considerations for “Today” and the “Day After” [published online ahead of print, 2020 July 7] [Impacto de la pandemia COVID-19 en los laboratorios de función pulmonar: consideraciones sobre el «hoy» y el «día después» [published online ahead of print, 2020 Jul 7]. Arch Bronconeumol. 2020;S0300-2896:30214-23, <http://dx.doi.org/10.1016/j.arbres.2020.07.001>.
2. Brudeman I, Abboud S. Telespirometry: novel system for home monitoring of asthmatic patients. Telemed J. 1997;3:127-33, <http://dx.doi.org/10.1089/tmj.1.1997.3.127>.
3. Rodriguez-Roisin R, Tetzlaff K, Watz H, Wouters E, Disse B, Finnigan H, et al. Daily home-based spirometry during withdrawal of inhaled corticosteroid in severe to very severe chronic obstructive pulmonary disease. Int J Chron Obstruct Pulmon Dis. 2016;11:1973-81, <http://dx.doi.org/10.2147/COPD.S106142> [published 22.08.16].
4. Moreira MA, Benedetto IG, Machado BC, Gonçalves M, Schneider L, Gadenz S, et al. RespiraNet Program: the first tele-spirometry service in South Brazil. Eur Respir J. 2015;46:PA2781, <http://dx.doi.org/10.1183/13993003.congress-2015.PA2781>.
5. Marina MN, López de Santa María E, Gutiérrez A, Bayón JC, García L, Gálvez JB. Telemedicine spirometry training and quality assurance program in primary care centers of a public health system. Telemed J E Health. 2014;20:388-92, <http://dx.doi.org/10.1089/tmj.2013.0111>.
6. Masa JF, González MT, Pereira R, Mota M, Riesco JA, Corral J, et al. Validity of spirometry performed online. Eur Respir J. 2011;37:911-8, <http://dx.doi.org/10.1183/09031936.00011510>.
7. Averame G, Bonavia M, Ferri P, Moretti AM, Fogliani V, Cricelli C, et al. Office spirometry can improve the diagnosis of obstructive airway disease in primary care setting. Respir Med. 2009;103:866-72, <http://dx.doi.org/10.1016/j.rmed.2008.12.017>.

Mário Morais-Almeida ^{a,◊}, Miguel T. Barbosa ^{a,b,*◊}, Cláudia S. Sousa ^{a,c}, Isabel Almeida ^a, Lara Pimenta ^a, Rita Aguiar ^a

^a Allergy Centre, CUF Descobertas Hospital, Lisboa, Portugal

^b Pulmonology Department, Hospital Centre of Barreiro-Montijo, Barreiro, Portugal

^c Pulmonology Department, Central Hospital of Funchal, Portugal

* Corresponding author.

E-mail address: migueltrbarbosa@gmail.com (M.T. Barbosa).

◊ Both authors contributed equally.

<https://doi.org/10.1016/j.arbres.2020.10.011>

0300-2896/ © 2020 SEPAR. Published by Elsevier España, S.L.U. All rights reserved.