




# ARCHIVOS DE Bronconeumología

www.archbronconeumol.org



## Original Article

# Associations of Cardiorespiratory Fitness and Muscular Endurance Fitness With Pulmonary Function in Physically Active Young Adults

Chia-Hsin Liu<sup>a</sup>, Yen-Chen Lin<sup>b</sup>, Wei-Chun Huang<sup>c,d</sup>, Xuemei Sui<sup>e</sup>, Carl J. Lavie<sup>f</sup>, Gen-Min Lin<sup>g,h,\*</sup>

<sup>a</sup> Division of Pulmonary and Critical Care Medicine, Department of Internal Medicine, Tri-Service General Hospital, National Defense Medical Center, Taipei, Taiwan

<sup>b</sup> Department of Internal Medicine, Linkou Chang-Gung Memorial Hospital, Taoyuan, Taiwan

<sup>c</sup> College of Medicine, National Yang Ming Chiao Tung University, Taipei, Taiwan

<sup>d</sup> Department of Critical Care Medicine, Kaohsiung Veterans General Hospital, Kaohsiung, Taiwan

<sup>e</sup> Arnold School of Public Health, University of South Carolina, Columbia, SC 29208, USA

<sup>f</sup> John Ochsner Heart and Vascular Institute, Ochsner Clinical School, The University of Queensland School of Medicine, New Orleans, LA, USA

<sup>g</sup> Department of Medicine, Hualien Armed Forces General Hospital, Hualien City, Taiwan

<sup>h</sup> Department of Medicine, Tri-Service General Hospital and National Defense Medical Center, Taipei, Taiwan

## ARTICLE INFO

### Article history:

Received 19 February 2024

Accepted 12 June 2024

Available online xxx

### Keywords:

Cardiorespiratory fitness

Muscular endurance fitness

Lung function

FEV<sub>1</sub>

FVC

## ABSTRACT

**Rationale:** While the beneficial effects of physical fitness on general health are well-documented, the specific relationship between different types of physical fitness, particularly cardiorespiratory fitness (CRF) and muscular endurance fitness (MEF), and lung function in physically active young adults remains less explored.

**Objective:** This study investigated the relationship between CRF and MEF, and their correlation with lung function in physically active young adults.

**Methods:** This cross-sectional study involved a cohort of 1227 physically active young adults without lung diseases. Lung function was assessed using FEV<sub>1</sub>, FVC, and FEV<sub>1</sub>/FVC measurements. The 3000-m run was used to assess CRF, and the 2-min push-up and sit-up tests were used to assess MEF. Multivariable linear regression analysis was used to evaluate the relationships between these fitness measures and lung function, adjusting for potential covariates.

**Results:** Enhanced CRF was associated with superior FEV<sub>1</sub> and FVC after adjusting for covariates ( $\beta = -.078$ ,  $p = .015$  for FEV<sub>1</sub>;  $\beta = -.086$ ,  $p = .009$  for FVC). Push-ups were positively associated with FEV<sub>1</sub> ( $\beta = .102$ ,  $p = .014$ ), but not with FVC. In contrast, sit-ups showed no significant correlation with lung function in the fully adjusted model.

**Conclusion:** The study demonstrated a clear association between improved physical fitness and better lung function in physically active young adults, with various exercises showing distinct associations with lung metrics. Notably, push-ups were particularly associated with higher FEV<sub>1</sub>. A future prospective study is necessary to determine whether routine exercises, such as push-ups, might lead to greater lung function.

© 2024 SEPAR. Published by Elsevier España, S.L.U. All rights are reserved, including those for text and data mining, AI training, and similar technologies.

## Introduction

Pulmonary function is a fundamental aspect of respiratory health, achieving its highest levels in young adulthood, around the ages of 20–25.<sup>1</sup> Tobacco smoke exposure is a well-documented major risk factor for “impaired respiratory health”—a condition that can progress to chronic lung disease due to diminished lung growth

and function.<sup>2</sup> Furthermore, recent research also suggests associations between impaired respiratory health and factors such as air pollution, obesity, and low physical fitness levels.<sup>2,3</sup> A reduction in lung function not only limits exercise capacity but also increases the risk of respiratory and cardiovascular diseases, lowers health-related quality of life (HRQoL), and results in increased respiratory hospitalizations, mortality, and healthcare demands.<sup>4–6</sup> Therefore, it is crucial to identify and address these modifiable risk factors to preserve and enhance lung function.

Physical fitness serves as a critical indicator of health outcomes and influences long-term health.<sup>7</sup> Studies have evaluated physi-

\* Corresponding author.  
E-mail address: farmer507@yahoo.com.tw (G.-M. Lin).

cal fitness through cardiorespiratory fitness (CRF) and muscular fitness, using various tests to measure them either separately or together.<sup>8</sup> Research suggests that enhanced aerobic fitness in childhood and adolescence plays a significant role in lung development, leading to improvements in FEV<sub>1</sub> and FVC from ages 9 to 26.<sup>9</sup> Additionally, young adults with higher baseline CRF levels, as measured by treadmill tests, showed reduced declines in FEV<sub>1</sub> and FVC over 20 years, according to the Coronary Artery Risk Development in Young Adults (CARDIA) study.<sup>10</sup> While these findings suggest that CRF contributes to lung development and may help protect against future declines in lung function, the relatively small effect sizes reported indicate that the impact of fitness on lung function requires further clarification. Furthermore, the causal relationship between improved lung function and gains in fitness remains lacking. In the cohort, when the data was analyzed with lung function as the independent variable and fitness as the dependent variable, it yielded similar results.<sup>9</sup> These results suggest a bidirectional relationship: individuals with superior lung function might be predisposed to higher physical fitness levels.

Muscular fitness, characterized by both muscular endurance (repeated force exertion) and muscular strength (single maximal effort), is essential for sustained activity and peak performance.<sup>11</sup> In children and adolescents, higher muscular fitness correlates with lower obesity rates, diminished cardiometabolic risks, and improved HRQoL.<sup>12,13</sup> A meta-analysis highlighted that strong muscular fitness during these early years is associated with less body fat and better cardiometabolic health later in life.<sup>14</sup> Additionally, a recent study has found a positive link between muscular strength (measured by handgrip strength and chair stand tests) and lung function in a community general population.<sup>15</sup> However, the exact relationship between CRF, muscular endurance fitness (MEF), and their impact on lung function, especially among physically active individuals, is not fully understood.

The Cardiorespiratory Fitness and Health In Armed Forces (CHIEF) Lung Study aims to evaluate the association between CRF (time for a 3000-m run) and MEF (2-min push-ups and 2-min sit-ups numbers) with pulmonary function, and to clarify their individual impacts on the respiratory health of active young adults. We hypothesize that superior CRF and MEF levels are positively associated with pulmonary function, with each type of fitness making a unique correlation.

## Methods

### Study population

The CHIEF Lung Study meticulously assessed lung function in 1231 military recruits, both male and female, aged 19–38, across Taiwan from 2017 to 2019 (Fig. 1). Participants underwent medical exams for military enlistment, with no history of pulmonary tuberculosis, chronic lung conditions (e.g., lung abscess, emphysema, asthma), diaphragmatic or pleural disorders, chest wall deformities, or surgeries for pneumothorax or segmentectomy. They participated in a rigorous daily exercise regimen at a military base for six months, focusing on muscular endurance training and cardiovascular fitness. The regimen required all military personnel to complete a 3000-m run twice daily at 6:00 AM and 4:00 PM, led by the captain at a pace of 120 m per minute on weekdays. Following each run, all participants were required to perform 20 consecutive push-ups and sit-ups within 30 min. This training course lasted until the end of June. Before the mid-term physical test in July, all participants underwent an annual health checkup in June, which included laboratory tests and physical evaluations. They also filled out a questionnaire about substance use habits, such as cigarette smoking (active and former/never), alcohol consumption, and betel

nut chewing at the Hualien Armed Forces General Hospital. Furthermore, participants underwent pulmonary function testing to assess their FEV<sub>1</sub>, FVC, and the ratio of FEV<sub>1</sub>/FVC before undergoing the annual military exercise tests. The study adhered to stringent ethical standards and was approved by the Institutional Review Board of Mennonite Christian Hospital in Hualien City (No. 16-05-008), ensuring that all participants provided informed consent.

### Blood pressure and anthropometric measurements

Blood pressure (BP) was measured once on each participant's right arm after a 15-min rest, using an automatic device (FT201) via the oscillometric method. Mean Arterial Pressure (MAP) was calculated using the formula: MAP = diastolic BP + 1/3 (systolic BP – diastolic BP). Anthropometric data, including waist circumference (WC), height, and weight, were recorded for standing participants, with body mass index (BMI) calculated as weight (kg) divided by height squared (m<sup>2</sup>).

### MEF and CRF assessments

MEF in participants was assessed using 2-min push-up and sit-up tests, adhering to protocols from previous studies.<sup>16</sup> These evaluations were scheduled between 14:00 and 15:00 at the Military Physical Training and Testing Center, allowing for a flexible sequence. Performance was recorded on sponge pads with electronic or infrared sensors: sit-ups required elbow contact with thigh-mounted sensors, with hands near ears and feet fixed; push-ups demanded correct elbow motion and a straight body alignment, confirmed by sensors, disqualifying any touch outside of hands or toes before time. CRF was gauged through a 3000-m run, performed outdoors on level ground after an hour's rest from MEF tests. This field test, held outdoors on flat terrain at the Military Physical Training and Testing Center, was scheduled for 16:00, following an hour's rest after the muscular endurance tests. Participants wore standard sweat suits and carried no extra items.<sup>17</sup> The entire run for each participant was video-recorded and monitored by eight military sports officers. The run was contingent upon certain environmental factors: the coefficient, determined by the product of the outdoor temperature (in Celsius) and the relative humidity (%) multiplied by 0.1, had to be below 40, and there should not have been any significant rainfall. The completion time of the 3000-m run was used to assess each participant's cardiorespiratory endurance capacity.

### Pulmonary function assessment

Lung function was measured with a Jaeger Vyntus™ Pneumo spirometer as per the American Thoracic Society.<sup>18</sup> Participants, seated with a nose clip, performed the test thrice for accurate FVC, FEV<sub>1</sub>, and FEV<sub>1</sub>/FVC ratios, inhaling fully and exhaling rapidly for 6 s. The best effort was recorded via a flow-volume curve. Lung function is expressed as Z-scores using the Global Lung Initiative reference equations, accounting for height, sex at birth, age, and Asian ethnicity. In addition, the predicted percentage of lung function was also presented.

### Statistical analyses

Continuous variables were presented as mean ± standard deviation (SD) and categorical ones as frequencies and percentages (%). Descriptive statistics described demographics and clinical characteristics. The fitness data are presented as the median (interquartile range). Pearson's correlation and scatter plots with *r* and *p* values analyzed the relationship between pulmonary parameters and the performance of each physical test. Multivariable linear regression analysis was used to examine the associations between physical




Fig. 1. The flow diagram of participant selection and evaluation process for the CHIEF Lung Study.

performance measurements and lung functions. The analysis was stratified into four distinct models, each designed to incrementally build upon the previous by integrating additional variables. Model 1 served as the baseline, adjusting for demographic and physiological factors including age, gender, height, weight, waist circumference, mean arterial pressure, and smoking status. Model 2 expanded on this by incorporating the 3000-m run time and the number of 2-min push-ups, aiming to assess their specific impacts. Model 3 followed a similar format but focused on the 3000-m run time and 2-min sit-ups. Finally, Model 4, the most comprehensive of the series, combined all the metrics from the previous models to provide a holistic view of the effects of combined physical activities on lung function. Results across these models were expressed through standardized beta coefficients, with significance levels noted for key findings ( $*p \leq 0.05$ ,  $**p \leq 0.01$ ). The  $p$ -interaction value was calculated to assess whether there are sex differences in the association between physical fitness and lung function. Analysis was performed using SPSS (version 29).

## Results

Out of the initial cohort, four were excluded due to acute illnesses, leaving 1227 participants in the final analysis. Clinical characteristics of participants are shown in Table 1. The participants, primarily male at 88.2%, had an average age of 27.4 years. They maintained a healthy physique with an average BMI of  $24.6 \text{ kg/m}^2$ , and a waist circumference of 82.3 cm. Smoking status varied, with 38% currently smoking and 54% either former smokers or having never smoked. Regarding physical fitness, the participants completed a 3000-m run in an average time of 15 min and 12 s and demonstrated considerable strength and endurance by performing an average of 44.1 push-ups and 44.6 sit-ups within a two-minute timeframe for each exercise. In the study comparing clinical and physiological characteristics between sexes, notable differences were observed. Males were older, taller, and heavier than females, with correspondingly higher BMI and waist circumferences. Significant differences in smoking habits were evident, with a higher percentage of males being active or former smokers compared to females, where a majority were never smokers. Physical fitness assessments showed that males performed bet-

ter in muscular endurance exercises like push-ups and sit-ups, and demonstrated superior cardiorespiratory fitness, evidenced by shorter times in the 3000-m run compared to females. Lung function measurements revealed higher absolute values of FEV<sub>1</sub> and FVC, but no differences were observed in standardized lung function metrics such as predicted percentages and Z-scores.

Pearson's correlation coefficients were used to assess the relationship between spirometer parameters and physical performance tests, specifically focusing on CRF and MEF. The relationship between pulmonary function and performance in the 3000-m run test, as well as the outcomes of the 2-minute push-up and sit-up tests, is illustrated in Fig. 2. There is a negative correlation between the time for a 3000-m run and pulmonary function, with FEV<sub>1</sub> (Fig. 2A) and FVC (Fig. 2B) both decreasing as the run time increases ( $r = -0.29$ ,  $p < 0.0001$  for FEV<sub>1</sub>;  $r = -0.27$ ,  $p < 0.0001$  for FVC). No significant correlation is observed with the FEV<sub>1</sub>/FVC ratio. Furthermore, there is a positive correlation between the number of 2-min push-ups and pulmonary function, with both FEV<sub>1</sub> (Fig. 2C) and FVC (Fig. 2D) increasing as the number of push-ups increases ( $r = 0.32$ ,  $p < 0.0001$  for FEV<sub>1</sub>;  $r = 0.29$ ,  $p < 0.0001$  for FVC). Again, no significant correlation is observed with the FEV<sub>1</sub>/FVC ratio. Similarly, a positive correlation is shown between the number of 2-minute sit-ups and pulmonary function, with FEV<sub>1</sub> (Fig. 2E) and FVC (Fig. 2F) both increasing with a greater number of sit-ups performed ( $r = 0.25$ ,  $p < 0.0001$  for FEV<sub>1</sub>;  $r = 0.24$ ,  $p < 0.0001$  for FVC). The FEV<sub>1</sub>/FVC ratio does not show a significant correlation. These results suggest that better performance in fitness tests, specifically faster run time and higher numbers of push-ups and sit-ups, is associated with better pulmonary function, as measured by FEV<sub>1</sub> and FVC. However, these fitness performances do not appear to affect the FEV<sub>1</sub>/FVC ratio significantly. We also analyzed the correlation between physical fitness and standardized lung function using Z-scores. The results showed that when lung function was presented as Z-scores, there was no significant correlation between physical fitness and lung function (Fig. E1, Supplementary material). To ensure the results were not influenced by outlier effects, we excluded outliers defined by more than 3 standard deviations from the mean for physical fitness and lung function. Despite this, the correlation between absolute lung function measurements and physical fitness remains significant (Fig. E2, Supplementary material). Given the

**Table 1**  
Clinical characteristics of study population.

Characteristics	Total population (n = 1227)	Male (n = 1082)	Female (n = 145)	p value
<b>Age, years</b>	27.4 ± 6.2	27.6 ± 6.3	26.2 ± 5.2	0.011
<b>Height (cm)</b>	170.7 ± 7.1	172.0 ± 6.1	160.2 ± 4.9	<0.001
<b>Weight (kg)</b>	71.8 ± 13.4	73.5 ± 13.1	59.3 ± 8.2	<0.001
<b>Body mass index (kg/m<sup>2</sup>)</b>	24.6 ± 3.9	24.8 ± 3.9	23.1 ± 3.0	<0.001
<b>Waist (cm)</b>	82.3 ± 10.6	83.3 ± 10.5	74.9 ± 7.7	<0.001
<b>Mean arterial pressure (mmHg)</b>	85.5 ± 10.3	86.1 ± 10.4	81.1 ± 8.7	<0.001
<b>Smoking</b>				
Active smoker	464 (38)	437 (40)	27 (19)	<0.001
Former smoker	61 (5)	57 (5)	4 (3)	0.1192
Never smoker	600 (49)	497 (46)	103 (71)	<0.001
Alcohol	426 (35)	407 (38)	19 (13)	<0.001
Betel nut	145 (12)	144 (13)	1 (1)	<0.001
<b>Physical function measurements</b>				
3000-m run (s)	878 (840–959)	870 (834–930)	1027 (985–1084)	<0.001
2-min push-ups numbers	45 (40–50)	45 (40–50)	25 (21–33)	<0.001
2-min sit-ups numbers	45 (40–50)	45 (40–50)	35 (30–40)	<0.001
<b>Lung function measurements</b>				
FEV <sub>1</sub>				
Actual (L)	3.4 ± 0.6	3.5 ± 0.5	2.6 ± 0.4	<0.001
% predicted	90.8 ± 14.3	90.8 ± 14.7	90.9 ± 10.9	0.926
Z-score	-0.69 ± 1.0	-0.68 ± 1.0	-0.74 ± 0.9	0.494
FVC				
Actual (L)	4.0 ± 0.7	4.2 ± 0.6	3.0 ± 0.5	<0.001
% predicted	93.0 ± 14.3	92.9 ± 13.6	93.5 ± 11.3	0.587
Z-score	-0.55 ± 1.0	-0.55 ± 1.0	-0.54 ± 0.9	0.844
FEV <sub>1</sub> /FVC %				
Actual (%)	84.4 ± 9.3	84.3 ± 9.0	85.3 ± 11.9	0.205
% predicted	86.8 ± 1.9	86.7 ± 2.0	86.9 ± 1.5	0.254
Z-score	-0.25 ± 1.2	-0.23 ± 1.2	-0.38 ± 1.1	0.171

FEV<sub>1</sub>, forced expiratory volume in 1 s; FVC, forced vital capacity. Data are presented as number (percent) for smoking, alcohol, and betel nut. Physical function measurements were presented as median (interquartile range). For all other variables, means ± SD are reported.

established correlation between tobacco consumption and physical capacity, we stratified physical fitness by cigarette smoking status. The results show that active smokers exhibited better physical fitness performance in activities such as the 3000-m run, 2-min push-ups, and 2-min sit-ups (Fig. E3, Supplementary material).

To elucidate the relationship between CRF and MEF, data from scatter plots depicted in Fig. 3A reveal a notable negative correlation ( $r = -0.48, p < 0.0001$ ) between the number of 2-min push-ups and time for a 3000-m run. Higher numbers of push-ups correlate with shorter run times, suggesting enhanced CRF. Similarly, a negative correlation ( $r = -0.40, p < 0.0001$ ) between 2-minute sit-up numbers and run time is shown in Fig. 3B, indicating that greater muscular endurance fitness from sit-ups is associated with improved CRF. The relationship is further underscored in Fig. 3C, which shows a strong positive correlation ( $r = 0.752, p < 0.0001$ ) between performance in push-ups and sit-ups, indicating a consistent link between these two forms of exercise. Collectively, these findings highlight the interrelation of MEF and CRF, as well as the association between endurance in upper and lower limb exercises.


To determine the distinct impacts of various physical fitness modalities on pulmonary function, we utilized Multivariable linear regression analyses, as depicted in Table 2. In Model 1, after adjusting for variables such as age, gender, height, weight, WC, MAP, and smoking status, the results showed that time for a 3000-m run was inversely associated with both FEV<sub>1</sub> (standardized  $\beta$ :  $-0.102, p < .001$ ) and FVC (standardized  $\beta$ :  $-0.117, p < .001$ ). Additionally, 2-min push-ups positively correlated with both FEV<sub>1</sub> (standardized  $\beta$ :  $.093, p = .002$ ) and FVC (standardized  $\beta$ :  $.096, p = .002$ ). In contrast, 2-min sit-ups were only positively associated with FVC (standardized  $\beta$ :  $.063, p = .034$ ) and not with FEV<sub>1</sub>.

To explore the potential associations between these physical activities and lung function, we formulated Model 2–4. These models incorporated the same variables as Model 1, with the addi-


tion of either a 3000-m run, 2-min push-ups, or 2-min sit-ups. In both Model 2 and Model 3, the duration of the 3000-m run consistently demonstrated an inverse correlation with FEV<sub>1</sub> (standardized  $\beta$ :  $-0.071, p = .026$ ; standardized  $\beta$ :  $-0.089, p = .004$ ) and FVC (standardized  $\beta$ :  $-0.087, p = .008$ ; standardized  $\beta$ :  $-0.093, p = .003$ ). Nevertheless, 2-min push-ups only positively correlated with FEV<sub>1</sub> (standardized  $\beta$ :  $.067, p = .043$ ) but not with FVC after adjusting for time for the 3000-m run. After making the same adjustment, the relationship between 2-min sit-ups and lung functions became non-significant.

After adjusting for all covariates and incorporating the three physical activities into the Model 4, time for the 3000-m run continued to show a negative relationship with FEV<sub>1</sub> (standardized  $\beta$ :  $-0.078, p = .015$ ) and FVC (standardized  $\beta$ :  $-0.086, p = .009$ ). Notably, the number of 2-min push-ups maintained its positive relationship with FEV<sub>1</sub> (standardized  $\beta$ :  $.102, p = .014$ ) even after adjusting for time for the 3000-m run and 2-min sit-ups, indicating its unique and independent association. In contrast, when considering the performances of time for the 3000-m run and 2-min push-ups, the number of 2-min sit-ups lacked a significant relationship with both FEV<sub>1</sub> and FVC. In addition, we also calculated a *p*-interaction value in Table E1 (Supplementary material) to determine whether sex moderates the association between physical fitness and lung function. The results showed that sex has no interaction effect on the association between physical fitness and lung function.

To investigate the contribution of lung function to maximum physical fitness, we conducted a multivariate linear regression analysis, adjusting for covariates such as age, gender, height, weight, MAP, and smoking status. The analysis revealed that pulmonary function measures, when expressed as Z-scores, did not show significant correlations with physical fitness tests (Table E2, Supplementary material). In contrast, the absolute values of FEV<sub>1</sub> and FVC were positively correlated with enhanced



**Fig. 2.** Scatter plots representing relationships between lung function metrics and physical tests: (A–C) Correlations of FEV<sub>1</sub>, FVC, and FEV<sub>1</sub>/FVC with 3000-m run time. (D–F) FEV<sub>1</sub>, FVC, and FEV<sub>1</sub>/FVC with 2-min push-up count. (G–I) FEV<sub>1</sub> and FVC with 2-min sit-ups. The correlation coefficient (*r*) and *p*-value for each relationship are indicated above each plot. FEV<sub>1</sub>, forced expiratory volume in 1 s; FVC, forced vital capacity; s, seconds. Time is represented in seconds (s).



**Fig. 3.** Scatter plots illustrating relationships: (A) 3000-m run time vs. 2-min push-ups, (B) 3000-m run time vs. 2-min sit-ups, and (C) 2-min push-ups vs. 2-min sit-ups. The correlation coefficient (*r*) and *p*-value for each relationship are indicated above each plot. Time is represented in seconds (s).



**Table 2**  
Multivariable linear regression analysis for the correlations of various physical fitness modalities with pulmonary functions.

Variables	Model 1				Model 2				Model 3				Model 4			
	FEV <sub>1</sub>		FVC		FEV <sub>1</sub>		FVC		FEV <sub>1</sub>		FVC		FEV <sub>1</sub>		FVC	
	$\beta$	<i>p</i>	$\beta$	<i>p</i>	$\beta$	<i>p</i>	$\beta$	<i>p</i>	$\beta$	<i>p</i>	$\beta$	<i>p</i>	$\beta$	<i>p</i>	$\beta$	<i>p</i>
Time for a 3000-m run (s)	-.102	<.001**	-.117	<.001**	-.071	.026*	-.087	.008**	-.089	.004**	-.093	.003**	-.078	.015*	-.086	.009**
2-min push-ups numbers	.093	.002**	.096	.002**	.067	.043*	.056	.096					.102	.014*	.066	.122
2-min sit-ups numbers	.034	.244	.063	.034*					.009	.775	.033	.289	-.056	.158	-.190	.849

Multivariable linear regression analysis was used to determine the associations between physical performance and lung functions, with the following adjustments, including exposures (different types of physical performance) and confounders. Model 1: age, gender, height, weight, waist, mean blood pressure, smoking; Model 2: age, gender, height, weight, waist, mean blood pressure, smoking, 3000 m-run time, and 2-min push-up numbers; Model 3: age, gender, height, weight, waist, mean blood pressure, smoking, 3000 m-run time, and 2-min sit-up numbers; Model 4: age, gender, height, weight, waist, mean blood pressure, smoking, 3000 m-run time, and 2-min push-up numbers, 2-min sit-up numbers;  $\beta$  denotes standardized  $\beta$  coefficient.

\*  $p \leq 0.05$ .

\*\*  $p \leq 0.01$ .

Abbreviations: FEV<sub>1</sub>, forced expiratory volume in 1 second; FVC, forced vital capacity; L, liter; s, second.

performance in the 3000-m run and push-ups (Table E3, Supplementary material). Additionally, FVC, but not FEV<sub>1</sub>, was associated with improved sit-up performance.

## Discussion

In this cross-sectional study of 1227 active young adults, we explored how physical fitness levels affect lung function. Results showed that better performance in CRF, via 3000-m run times, and MEF, assessed through 2-min push-ups and sit-ups, correlated positively with FEV<sub>1</sub> and FVC, but not with the FEV<sub>1</sub>/FVC ratio. We found a moderate link between 3000-m run times and push-up/sit-up numbers, with a strong association between push-ups and sit-ups. Multivariable linear analysis revealed that shorter run times were correlated with higher FEV<sub>1</sub> and FVC, independent of MEF outcomes. Notably, push-ups positively influenced FEV<sub>1</sub> alone, while sit-ups showed no significant lung function effect in the fully adjusted model.

Our study found that improved CRF positively correlates with FEV<sub>1</sub> and FVC in active young adults, consistent with other research across various demographics.<sup>9,19-22</sup> A longitudinal study tracking children into adulthood affirmed this positive CRF association with FEV<sub>1</sub> and FVC throughout all ages, but not with the FEV<sub>1</sub>/FVC ratio.<sup>9</sup> Enhancements in CRF during youth were linked to improved FEV<sub>1</sub> and FVC in later life. This beneficial association between CRF and ventilatory lung function has been observed in diverse populations, including middle-aged<sup>21</sup> and elderly cohorts,<sup>19</sup> and the general population.<sup>20,22</sup> Importantly, CRF might not just enhance lung volume but could also improve lung diffusing capacity (D<sub>LCO</sub>).<sup>19,21</sup> These consistent findings across studies and among different age groups highlight the substantial and positive impact of CRF on lung function throughout the life span.

The gold standard for assessing CRF is Cardiopulmonary Exercise Testing (CPET), which evaluates V<sub>O2max</sub> or V<sub>O2peak</sub>, and can be estimated from work rates or algorithms.<sup>23</sup> While previous research has linked CRF to lung function via V<sub>O2max</sub><sup>9</sup> or V<sub>O2peak</sub><sup>19-22</sup> measurements through CPET, its requirement for specific facilities and participant mask tolerance during testing limits its practicality, especially for children, the elderly, or large-scale studies.<sup>24</sup> The American Heart Association (AHA) endorses CPET and run field tests for assessing V<sub>O2max</sub> in youths.<sup>25</sup> Our prior research has found a significant correlation ( $r = 0.729, p < 0.001$ ) between 3000-m run times and V<sub>O2max</sub> scaled to body mass in young adults, indicating that the 3000-m run is an effective, accessible method for estimating CRF, suitable for larger populations or settings with limited resources.<sup>17</sup>

The correlation between decreased pulmonary function and impaired peripheral muscle strength and endurance has been established in patients with chronic obstructive pulmonary disease

(COPD).<sup>26-30</sup> A case-control study showed that handgrip muscle strength decreases as the FEV<sub>1</sub> and FVC decrease in patients with COPD compared to the normal subjects.<sup>29</sup> Moreover, it was observed that COPD patients experienced a pronounced decline in muscle endurance compared to muscle strength. Several studies have also confirmed a positive correlation between handgrip muscle strength and lung function across diverse populations, ranging from adolescents to the elderly.<sup>15,24,31-36</sup> These results suggest that an association between muscular strength and lung function was observed not only in COPD patients but also in the general community population. Nevertheless, the impact of various forms of peripheral muscle endurance on lung function has been poorly investigated in previous studies, yielding inconsistent results. One study found a positive association between muscle endurance, assessed by quadriceps, and FEV<sub>1</sub> in COPD patients,<sup>26</sup> but another study failed to identify a similar association when evaluating muscle endurance through handgrip.<sup>29</sup> These discrepancies might arise from the use of various muscle groups and measurement techniques, variability in the study populations, and small sample sizes. Our study examined the relationship between both upper and lower limb muscle endurance, separately measured by the number of 2-min push-ups and sit-ups, and lung function, revealing a positive correlation in a large cohort of physically active young adults without chronic lung diseases. Initially, both 2-min push-ups and sit-ups showed a positive correlation with FEV<sub>1</sub> and FVC in univariate analysis. Nevertheless, the link between 2-minute sit-ups and FEV<sub>1</sub> became insignificant after adjusting for covariates. In contrast, the 2-min push-ups maintained a positive correlation with FEV<sub>1</sub>, even after accounting for additional physical fitness metrics such as 2-min sit-ups and 3000-m run time. These findings indicate that upper limb muscle endurance has a greater impact on lung function, particularly on FEV<sub>1</sub>, compared to lower limb muscle endurance. Consequently, our findings propose that the 2-min push-ups fitness routine, focused on improving upper limb muscle and pectoral muscle endurance, might be beneficial in managing obstructive lung diseases like COPD or asthma with airflow limitation. Particularly, the performance of the 3000-m run remained positively correlated with both FEV<sub>1</sub> and FVC across all models, indicating its robust impact on overall lung function.

The mechanisms connecting improved or enhanced physical fitness with better lung function are not fully understood. It is plausible that regular exercise training, a determinant of CRF, may strengthen the respiratory muscles, enhance pulmonary perfusion as well as surfactant release, and ultimately lead to increased lung volumes.<sup>37-39</sup> Additionally, enhanced fitness may decrease fat distribution, particularly abdominal and thoracic fat mass, which can constrain vital capacity and lead to expiratory flow limitation by restricting lung expansion during inhalation.<sup>40</sup> Moreover, reduced

fat mass may also alleviate lung function decline by reducing inflammatory processes, as adipose tissue is a source of inflammatory mediators that can harm lung tissue and narrow airway diameter.<sup>41</sup> The positive correlation between the enhanced 2-min push-up test and lung function might be attributed to the fact that push-ups activate not only the arms and chest but also the accessory respiratory muscles, including the intercostal muscles, pectoral muscles, and those around the shoulder girdle. Furthermore, the repetitive and sustained strengthening of these muscles could improve their efficiency and capacity, thereby contributing to improved lung function.

In the study, we analyzed the correlation between lung function and physical fitness, presenting lung function as either absolute values or Z-scores, which resulted in inconsistent findings. The inconsistency in the correlation results between absolute lung function values and physical fitness versus standardized lung function (Z-scores) could be partly explained by the attenuation of the association when anthropometric variables were accounted for at an individual level. When lung function is presented with absolute lung volumes, the measurements reflect the remaining ventilatory capacity, which may correlate more directly with physical fitness due to the physical demands on lung capacity during exercise.<sup>42</sup> However, when lung function is standardized into Z-scores, these values are adjusted based on a population mean and standard deviation, accounting for variations due to factors like age, sex, and height.<sup>1</sup> This standardization can obscure the raw differences seen with absolute values. For example, in a cross-sectional study, males and females, even when matched on relative lung volume (like FEV<sub>1</sub>% predicted), had significantly different absolute lung volumes, with males having a higher volume.<sup>43</sup> This discrepancy can alter perceptions of lung capacity and its relation to physical fitness when using standardized scores. Therefore, the inconsistency in the results could arise from these underlying differences in measurement and interpretation, highlighting the need for future studies to determine which lung function measurement is appropriate to correlate with physical fitness.

Several strengths and limitations of our study warrant discussion. This research is significant for being the first to investigate the relationship between different physical fitness aspects, including CRF and MEF, and their specific impacts on lung function in a cohort of physically active young adults. It provides valuable benchmarks for physical performance and lung function metrics within this group, known for their peak physiological capacities, including optimal lung and cardiovascular efficiency. Our research uniquely focuses on active young adults without chronic lung conditions, offering clearer insights into the fitness and lung function link. However, the cross-sectional, observational design limits our ability to assert causality. Although enhanced lung function may be attributed to factors common to both exercise capacity and pulmonary metrics, the FEV<sub>1</sub> measurement remains positively correlated in the fully adjusted model, underscoring a distinct connection between push-ups and FEV<sub>1</sub>. Furthermore, our results indicate that active smokers demonstrated superior physical fitness performance in activities such as the 3000-m run, 2-min push-ups, and 2-min sit-ups. Nevertheless, it is important to note that our participant group is predominantly young, and detailed quantitative data on cigarette consumption and smoking duration are not available. Consequently, we are unable to conclusively determine the effects of smoking on physical performance. This limitation underscores the need for further research with more comprehensive smoking data to robustly assess the impact of tobacco use on physical fitness. Finally, the participant demographic, primarily Asian males, may not reflect the diversity needed for broader applicability. These points highlight the necessity for future studies with a prospective design, encompassing a wide range of ethnic backgrounds and a balanced gender distribution, to comprehensively

understand the causal relationship between MEF and pulmonary function.

In conclusion, this study demonstrated that improved CRF and MEF are positively associated with pulmonary function in active young adults. It notably suggests that push-ups are distinctly associated with increased FEV<sub>1</sub>, highlighting the potential of specific physical activities to influence lung health. Further research is required to determine whether routine exercises like push-ups can consistently enhance lung function among various demographic and health backgrounds.

### Sources of funding

This research was funded by TSGH, through awards TSGH-D-113114, and supported by the Medical Affairs Bureau of the Ministry of National Defense, under awards MND-MAB-D-113200.

### Author's contributions

G.M.L. designed the study. C.H.L. performed the data analysis. C.H.L. and G.M.L. interpreted the results of statistical analysis. C.H.L. wrote the manuscript. G.M.L., Y.C.L., W.C.H., X.S., and C.J.L. revised the manuscript critically for important intellectual contents. All authors contributed to the interpretations of the findings. All authors reviewed the manuscript.

### Role of the sponsors

The sponsors and funders of the study had no involvement or any influence in the study design, in the collection, analysis, and interpretation of data, in the writing of the manuscript, and in the decision to submit the manuscript for publication. The corresponding author confirms that he had full access to all the data and had the final responsibility for the decision to submit for publication.

### Data sharing statement

The datasets supporting the conclusions of this article will be made available by the authors upon reasonable request, subject to necessary approvals and confidentiality agreements. Specific data that are not publicly available due to privacy or ethical restrictions will be accessible subject to conditions outlined by the corresponding author.

### Conflict of interest

No conflicts exist for the authors.

### Acknowledgements

We extend our deepest gratitude to all participants and collaborators who made this study possible. Special thanks to TSGH and Ministry of National Defense R.O.C. for their financial support.

### Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at [doi:10.1016/j.arbres.2024.06.005](https://doi.org/10.1016/j.arbres.2024.06.005).

### References

1. Quanjer PH, Stanojevic S, Cole TJ, Baur X, Hall GL, Culver BH, et al. Multi-ethnic reference values for spirometry for the 3–95-yr age range: the global lung function 2012 equations. *Eur Respir J*. 2012;40(6):1324–43.
2. Reyfman PA, Washko GR, Dransfield MT, Spira A, Han MK, Kalhan R. Defining impaired respiratory health. A paradigm shift for pulmonary medicine. *Am J Respir Crit Care Med*. 2018;198(4):440–6.

3. Liu GY, Kalhan R. Impaired respiratory health and life course transitions from health to chronic lung disease. *Chest*. 2021;160(3):879–89.
4. Roman MA, Rossiter HB, Casaburi R. Exercise, ageing and the lung. *Eur Respir J*. 2016;48(5):1471–86.
5. Duong M, Islam S, Rangarajan S, Leong D, Kurmi O, Teo K, et al. Mortality and cardiovascular and respiratory morbidity in individuals with impaired FEV(1) (PURE): an international, community-based cohort study. *Lancet Glob Health*. 2019;7(5):e613–23.
6. Njoroge MW, Mjojo P, Chirwa C, Rylance S, Nightingale R, Gordon SB, et al. Changing lung function and associated health-related quality-of-life: a five-year cohort study of Malawian adults. *EclinicalMedicine*. 2021;41:101166.
7. Ruiz JR, Castro-Piñero J, Artero EG, Ortega FB, Sjöström M, Suni J, et al. Predictive validity of health-related fitness in youth: a systematic review. *Br J Sports Med*. 2009;43(12):909–23.
8. Appelqvist-Schmidlechner K, Vaara JP, Vasankari T, Hakkinen A, Mantysaari M, Kyrolainen H. Muscular and cardiorespiratory fitness are associated with health-related quality of life among young adult men. *BMC Public Health*. 2020;20(1):842.
9. Hancox RJ, Rasmussen F. Does physical fitness enhance lung function in children and young adults? *Eur Respir J*. 2018;51(2):1701374.
10. Benck LR, Cuttica MJ, Colangelo LA, Sidney S, Dransfield MT, Mannino DM, et al. Association between cardiorespiratory fitness and lung health from young adulthood to middle age. *Am J Respir Crit Care Med*. 2017;195(9):1236–43.
11. Knapik JJ, Rieger W, Palkoska F, Van Camp S, Darakjy S. United States Army physical readiness training: rationale and evaluation of the physical training doctrine. *J Strength Cond Res*. 2009;23(4):1353–62.
12. Garcia-Hermoso A, Ramirez-Campillo R, Izquierdo M. Is muscular fitness associated with future health benefits in children and adolescents? A systematic review and meta-analysis of longitudinal studies. *Sports Med*. 2019;49(7):1079–94.
13. Bermejo-Cantarero A, Alvarez-Bueno C, Martinez-Vizcaino V, Redondo-Tebar A, Pozuelo-Carrascosa DP, Sanchez-Lopez M. Relationship between both cardiorespiratory and muscular fitness and health-related quality of life in children and adolescents: a systematic review and meta-analysis of observational studies. *Health Qual Life Outcomes*. 2021;19(1):127.
14. de Lima TR, Martins PC, Moreno YMF, Chaput JP, Tremblay MS, Sui X, et al. Muscular fitness and cardiometabolic variables in children and adolescents: a systematic review. *Sports Med*. 2022;52(7):1555–75.
15. Landi F, Salini S, Zazzara MB, Martone AM, Fabrizi S, Bianchi M, et al. Relationship between pulmonary function and physical performance among community-living people: results from Look-up 7+ study. *J Cachexia Sarcopenia Muscle*. 2020;11(1):38–45.
16. Lin GM, Tsai KZ, Lee DC, Sui X, Lavie CJ. Mean power output for muscular endurance exercises and maximal oxygen uptake in military young adults. *Medicine (Baltimore)*. 2023;102(42):e35578.
17. Lin GM, Tsai KZ, Sui X, Lavie CJ. Estimated power output for a distance run and maximal oxygen uptake in young adults. *Front Physiol*. 2023;14:1110802.
18. Standardization of spirometry, 1994 update. American Thoracic Society. *Am J Respir Crit Care Med*. 1995;152(3):1107–36.
19. Hassel E, Stensvold D, Halvorsen T, Wisloff U, Langhammer A, Steinshamn S. Association between pulmonary function and peak oxygen uptake in elderly: the Generation 100 study. *Respir Res*. 2015;16:156.
20. Rasch-Halvorsen O, Hassel E, Langhammer A, Brumpton BM, Steinshamn S. The association between dynamic lung volume and peak oxygen uptake in a healthy general population: the HUNT study. *BMC Pulm Med*. 2019;19(1):2.
21. McNeill J, Chernofsky A, Nayor M, Rahaghi FN, San Jose Estepar R, Washko G, et al. The association of lung function and pulmonary vasculature volume with cardiorespiratory fitness in the community. *Eur Respir J*. 2022;60(2):2101821.
22. Ostolin T, Gonze BB, Sperandio EF, Arantes RL, Romiti M, Dourado VZ. Mediator effect of cardiorespiratory fitness on the association between physical activity and lung function in adults: cross-sectional results from the Epimov study. *Int J Environ Res Public Health*. 2022;19(15):9377.
23. Ross R, Blair SN, Arena R, Church TS, Després JP, Franklin BA, et al. Importance of assessing cardiorespiratory fitness in clinical practice: a case for fitness as a clinical vital sign: a scientific statement from the American Heart Association. *Circulation*. 2016;134(24):e653–99.
24. Sillanpää E, Stenroth L, Bijlsma AY, Rantanen T, McPhee JS, Maden-Wilkinson TM, et al. Associations between muscle strength, spirometric pulmonary function and mobility in healthy older adults. *Age (Dordr)*. 2014;36(4):9667.
25. Raghuvver G, Hartz J, Lubans DR, Takken T, Wiltz JL, Mietus-Snyder M, et al. Cardiorespiratory fitness in youth: an important marker of health: a scientific statement from the American Heart Association. *Circulation*. 2020;142(7):e101–18.
26. Serres I, Gautier V, Varray A, Prefaut C. Impaired skeletal muscle endurance related to physical inactivity and altered lung function in COPD patients. *Chest*. 1998;113(4):900–5.
27. Bernard S, LeBlanc P, Whittom F, Carrier G, Jobin J, Belleau R, et al. Peripheral muscle weakness in patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med*. 1998;158(2):629–34.
28. Coronell C, Orozco-Levi M, Mendez R, Ramirez-Sarmiento A, Galdiz JB, Gea J. Relevance of assessing quadriceps endurance in patients with COPD. *Eur Respir J*. 2004;24(1):129–36.
29. Shah S, Nahar P, Vaidya S, Salvi S. Upper limb muscle strength & endurance in chronic obstructive pulmonary disease. *Indian J Med Res*. 2013;138(4):492–6.
30. Jaitovich A, Barreiro E. Skeletal muscle dysfunction in chronic obstructive pulmonary disease. What we know and can do for our patients. *Am J Respir Crit Care Med*. 2018;198(2):175–86.
31. Bahat G, Tufan A, Ozkaya H, Tufan F, Akpınar TS, Akin S, et al. Relation between hand grip strength, respiratory muscle strength and spirometric measures in male nursing home residents. *Aging Male*. 2014;17(3):136–40.
32. Smith MP, Standl M, Berdel D, von Berg A, Bauer CP, Schikowski T, et al. Hand-grip strength is associated with improved spirometry in adolescents. *PLOS ONE*. 2018;13(4):e0194560.
33. Son DH, Yoo JW, Cho MR, Lee YJ. Relationship between handgrip strength and pulmonary function in apparently healthy older women. *J Am Geriatr Soc*. 2018;66(7):1367–71.
34. Mgbemena NC, Aweto HA, Tella BA, Emeto TI, Malau-Aduli BS. Prediction of lung function using handgrip strength in healthy young adults. *Physiol Rep*. 2019;7(1):e13960.
35. Kim S, Yoon HK, Rhee CK, Jung HW, Lee H, Jo YS. Hand grip strength and likelihood of moderate-to-severe airflow limitation in the general population. *Int J Chron Obstruct Pulmon Dis*. 2022;17:1237–45.
36. Hesselberg LM, Kyvsgaard JN, Stokholm J, Bisgaard H, Bonnelykke K, Chaves B. Handgrip strength associates with effort-dependent lung function measures among adolescents with and without asthma. *Sci Rep*. 2023;13(1):13044.
37. Hildebran JN, Goerke J, Clements JA. Surfactant release in excised rat lung is stimulated by air inflation. *J Appl Physiol Respir Environ Exerc Physiol*. 1981;51(4):905–10.
38. Harms CA, Wetter TJ, McClaran SR, Pegelow DF, Nickele GA, Nelson WB, et al. Effects of respiratory muscle work on cardiac output and its distribution during maximal exercise. *J Appl Physiol* (1985). 1998;85(2):609–18.
39. Summerhill EM, Angov N, Garber C, McCool FD. Respiratory muscle strength in the physically active elderly. *Lung*. 2007;185(6):315–20.
40. Salome CM, King GG, Berend N. Physiology of obesity and effects on lung function. *J Appl Physiol* (1985). 2010;108(1):206–11.
41. Hancox RJ, Poulton R, Greene JM, Filsell S, McLachlan CR, Rasmussen F, et al. Systemic inflammation and lung function in young adults. *Thorax*. 2007;62(12):1064–8.
42. Quanjer PH, Tammeling GJ, Cotes JE, Pedersen OF, Peslin R, Yernault JC. Lung volumes and forced ventilatory flows. *Eur Respir J*. 1993;6 Suppl. 16:5–40.
43. Ekström M, Schiöler L, Grønseth R, Johannessen A, Svanes C, Leynaert B, et al. Absolute values of lung function explain the sex difference in breathlessness in the general population. *Eur Respir J*. 2017;49(5):1602047.