

Clinical Image

Pulmonary Hamartoma With Rare Imaging Features

Xu-ping Chen<sup>a</sup>, Xiao-Dong Yang<sup>b</sup>, Shi-xu He<sup>a,\*</sup>

<sup>a</sup> Department of Respiratory and Critical Care Medicine, Guangyuan Central Hospital, Guangyuan, Sichuan Province, China

<sup>b</sup> West China Hospital of Sichuan University, Chengdu, Sichuan, China



**Fig. 1.** CT scan reveals numerous round nodules of varying sizes and densities with “popcorn calcification” shape, diffusely distributed primarily in the upper and middle lobes of the right lung.

A 34-year-old male non-smoker was admitted for incidentally found pulmonary nodules. He had previously been misdiagnosed with “pulmonary tuberculosis”. Computed tomography (Fig. 1) revealed numerous round nodules of varying sizes and densities, some with a characteristic “popcorn” calcification, diffusely distributed primarily in the upper and middle lobes of the right lung. The patient underwent a thoracotomy, during which many hard, grayish-white nodules measuring 0.3–3 cm in diameter were observed. The postoperative pathological results confirmed the diagnosis of pulmonary chondromatous hamartoma (PH) which primarily comprises cartilage. At his 4-year follow-up, the patient remained asymptomatic with no evidence of recurrence. PH is the most common benign tumor of the lung, characterized by solitary nodules on the periphery of the lungs, constituting 5–8% of all solitary pulmonary nodules.<sup>1</sup> However, the PH nodules reported in this case exhibit a popcorn-like diffuse distribution, which is extremely uncommon. Therefore, identification of pulmonary hamartomas is still necessary for cases involving multiple calcified nodules.

**Conflict of Interests**

The authors state that they have no conflict of interests.

**Reference**

1. Sodhi KS, Virmani V, Jindal SK, Khandelwal N. Pulmonary hamartoma. *Ann Acad Med Singap.* 2009;38:1110.

\* Corresponding author.  
E-mail address: [hsx2021@foxmail.com](mailto:hsx2021@foxmail.com) (S.-x. He).