

Journal Pre-proof

A thoracic complication of an abdominal procedure: massive hemothorax following percutaneous transhepatic portal vein angioplasty

Luis Gorospe Rut Romera-Sánchez Gemma María Muñoz-Molina



PII: S0300-2896(24)00419-8

DOI: <https://doi.org/doi:10.1016/j.arbres.2024.10.016>

Reference: ARBRES 3684

To appear in: *Archivos de Bronconeumología*

Received Date: 4 October 2024

Accepted Date: 29 October 2024

Please cite this article as: Gorospe L, Romera-Sánchez R, Muñoz-Molina GM, A thoracic complication of an abdominal procedure: massive hemothorax following percutaneous transhepatic portal vein angioplasty, *Archivos de Bronconeumología* (2024), doi: <https://doi.org/10.1016/j.arbres.2024.10.016>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2024 SEPAR. Published by Elsevier España, S.L.U. All rights are reserved, including those for text and data mining, AI training, and similar technologies.

A thoracic complication of an abdominal procedure: massive hemothorax following percutaneous transhepatic portal vein angioplasty

Luis Gorospe^{*a}, Rut Romera-Sánchez^a, Gemma María Muñoz-Molina^b

^a Department of Radiology, Ramón y Cajal University Hospital, Madrid, Spain

^b Department of Thoracic Surgery, Ramón y Cajal University Hospital, Madrid, Spain

*Corresponding author:

Luis Gorospe

Department of Radiology

Ramón y Cajal University Hospital

Ctra. de Colmenar Viejo Km 9,100

28034 Madrid, Spain

E-mail address: luisgorospe@yahoo.com

A 23-year-old liver transplant patient underwent a percutaneous transhepatic portal vein angioplasty (PTHPVA) as a procedure for correcting a clinically significant portal vein stenosis causing prehepatic portal hypertension (fig. 1A, B). Right after the PTHPVA procedure, the patient developed shortness of breath and hypotension. An emergency CT showed active bleeding originating in the right hemidiaphragm surface (fig 1C-E). Because the patient was hemodynamically unstable, he was immediately transferred to the operating room, where a defect in the right hemidiaphragm was detected and closed. Percutaneous transhepatic portal vein angioplasty (PTHPVA) is a procedure performed in patients with clinically significant portal vein stenosis that involves direct right midaxillary puncture of the liver with the use of fluoroscopic and/or ultrasound guidance.¹ Alternative techniques of portal vein angioplasty include a transsplenic and transjugular approaches, but these are less used.² Hemothorax is an uncommon but well-known complication of PTHPVA; the rather large diameter of the dilators and catheters used in the procedure as well as the short puncture tract (making hemostasis difficult) have been claimed as risk factor for hemothorax. Low insertion of the right diaphragm and a high position of the liver graft are also potential factors that may cause hemothorax.^{1,2} This case reminds us that abdominal percutaneous interventional procedures can cause thoracic complications.

References

1. Naidu SG, Alzubaidi SJ, Patel IJ, Iwuchukwu C, Zurcher KS, Malik DG, et al. Interventional Radiology Management of Adult Liver Transplant Complications. *Radiographics*. 2022;42:1705-23.
2. Haddad MM, Fleming CJ, Thompson SM, Reisenauer CJ, Parvinian A, Frey G, et al. Comparison of Bleeding Complications between Transplenic versus Transhepatic Access of the Portal Venous System. *J Vasc Interv Radiol*. 2018;29:1383-91.

Journal Pre-proof

Figure legend




Figure 1. A) Axial CT image obtained before the PTHPVA shows a portal vein stent (arrows) and a poststent portal vein dilatation (asterisk). B) Digital subtraction angiogram shows the portal vein stent (arrows) and the poststent right portal vein dilatation (asterisk). C) Axial maximum intensity projection (MIP) CT image performed after the PTHPVA procedure shows signs of an active bleeding (arrows) originating in the right hemidiaphragm. Note the presence of a right pleural effusion (asterisk). D) Coronal MIP CT image shows signs of an active bleeding within the pleural space (arrows). Note the presence of a large right pleural effusion (white asterisk) causing a contralateral mediastinal shift (black asterisk).