

Journal Pre-proof

The importance of ZIP code-related average household income on the severity of respiratory syncytial virus infection in infants

Rosa Rodríguez-Fernández MD PhD Jimena Pérez-Moreno MD
PhD Felipe González-Martínez MD PhD Blanca Toledo del Castillo
MD Alicia Fernández González MD María Isabel
González-Sánchez MD



PII: S0300-2896(24)00474-5

DOI: <https://doi.org/doi:10.1016/j.arbres.2024.12.002>

Reference: ARBRES 3705

To appear in: *Archivos de Bronconeumología*

Received Date: 30 August 2024

Accepted Date: 11 December 2024

Please cite this article as: Rodríguez-Fernández R, Pérez-Moreno J, González-Martínez F, Castillo BTd, González AF, González-Sánchez MI, The importance of ZIP code-related average household income on the severity of respiratory syncytial virus infection in infants, *Archivos de Bronconeumología* (2024), doi: <https://doi.org/10.1016/j.arbres.2024.12.002>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2024 SEPAR. Published by Elsevier España, S.L.U. All rights are reserved, including those for text and data mining, AI training, and similar technologies.

TITLE: The importance of ZIP code-related average household income on the severity of respiratory syncytial virus infection in infants.

Scientific letter

Importance of Zip Code in the Severity of Respiratory Syncytial Virus Bronchiolitis in Infants

Authors: Rosa Rodríguez-Fernández, MD, PhD^{1,2}, Jimena Pérez-Moreno MD, PhD^{1,2}, Felipe González-Martínez, MD PhD^{1,2}, Blanca Toledo del Castillo, MD^{1,2} Alicia Fernández González, MD¹, María Isabel González-Sánchez, MD¹.

¹ Servicio de Pediatría, Hospital Infantil Gregorio Marañón; ² Instituto de Investigación Sanitaria Gregorio Marañón (IISGM), Madrid, España;

Corresponding author:

Rosa Rodriguez-Fernández.

Servicio de Pediatría. Hospital Gregorio Marañón. Madrid.

Calle O'Donnell 50, 28007, Madrid, España.

Teléfono: +34619312632

e-mail: rrfernandez@salud.madrid.org

Word count: 1,279 words

To the Editor,

Respiratory syncytial virus (RSV) bronchiolitis is the leading cause of hospitalization in infants, is responsible for more than 3 million of hospital admissions in infants

worldwide and is the second leading cause of death in infants under 12 months of age, especially in countries with low socioeconomic resources^{1, 2}.

Some factors, including age less than three months old, premature birth, chronic illnesses, or malnourishment, have been associated with an increased risk of RSV infection^{2, 3}. It is uncertain, nevertheless, how social determinants of health play a role in the severity of RSV bronchiolitis.

There are some studies suggesting an association between lower socioeconomic status and the incidence of RSV infection, emergency department attendance and hospital admissions^{4, 5}. However, there are very few studies linking socioeconomic status to the severity of RSV bronchiolitis⁶⁻⁸. Addressing health inequity is crucial, especially now, with the availability of RSV prevention strategies with long half-life monoclonal antibodies for all infants or maternal RSV vaccine^{9, 10}.

The main objective of this exploratory study was to determine whether mean household income, related to zip code, could affect the severity of RSV infection in infants admitted to a hospital.

We conducted a retrospective observational study in a single center including all infants under 2 years of age hospitalized with a diagnosis of RSV infection over a period of 7 calendar years from January 1, 2015 to December 31, 2021. Patients were identified using ICD- 10 discharge diagnosis codes, including codes J21.0, acute RSV bronchiolitis; J12.1, RSV pneumonia; and J21.9, unspecified bronchiolitis.

Demographic, epidemiologic, clinical, laboratory, radiologic data, and medical interventions during hospitalization were collected from the electronic health care records (EHR). Disease severity was assessed using length of stay (LOS), need for high flow nasal cannula (HFNC), non mechanical ventilation (NMV), mechanical

ventilation (MV), need for admission to the pediatric intensive care unit (PICU) and the Wood-Downes severity score¹¹. Microbiological diagnosis was done by rapid antigen test (2015-2019) or by RCP (2020-2021).

Socioeconomic data comes from *Geoportal datos del Ayuntamiento de Madrid*, *Portal de datos abiertos del Ayuntamiento de Madrid IGUALA*:

<https://igualamadrid.es/pages/presentacion>, *Ranking de vulnerabilidad según distritos en Madrid en 2020-2021*, Instituto Nacional de Estadística, Ministerio de Hacienda, and Comunidad Autónoma de Madrid.

The average household income in Madrid's districts was 35,510 euros. Each patient was assigned the average household income corresponding to his or her zip code and the database was divided into four groups:

- Group 1: 20,000-29,999 euros
- Group 2: 30,000-39,999 euros
- Group 3: 40,000-49,999 euros
- Group 4: >50,000 euros.

A sensitivity analysis was conducted using the economic vulnerability index, which takes into account three factors: households with incomes below 23,076 Euros, adults working less than 20%, and households unable to afford basic items. We considered a economic vulnerability test high when the value is above the median (>5.32)

Statistical analysis was performed using R statistical software, version 3.5.3.¹² The association between qualitative variables was analyzed using Fisher or Chi-square test and quantitative variables with Kruskal-Wallis or Man-Whitney test and most appropriate post-hoc analyses. To assess the severity of infection, logistic regression

models were built and included biologically important variables or those that were significant in bivariate analyses. Associations of risk factors with qualitative variables (PICU admissions) were displayed as odds ratios (ORs) and 95% CI. A p value <0.05 was considered significant.

From 2015-2021 a total of 1,080 infants under 2 years of age were hospitalized with a laboratory- confirmed diagnosis of RSV bronchiolitis, and 1,058 (98%) were successful geocoded in Comunidad Autonoma de Madrid and therefore included in the analysis. Eight hundred and sixty-fourth children (81,6%) lived in the metropolitan area of Madrid and the rest outside of the city. Median age was 3 (1.5-7) months, 53.4% were male, LOS was 5 (3-7) days and 21.7% required admission to the PICU (Table 1).

Three hundred and two children (28.5%) lived in households with a mean income between 20,000-29,999 euros (Group 1), 37.5% (n=397) had a mean income between 30,000-39,999 euros (Group 2), 11.3% between 40,000-49,999 euros (Group 3) and 22.3% had an income above 50,000 euros (Group 4).

Lower household incomes were associated with lower proportions of breastfeeding higher exposure to tobacco smoke and lower day care attendance ($p<0,01$) (Table1)

Severity of RSV bronchiolitis was analyzed according to average household income and we found that infants with lower family income, had a higher proportion of severe scores [Group1: 129/302(42,7%) vs Group 2: 139/397 (35%) vs Group 3: 35/120 (29,2%) vs Group 4: 68/235 (28,9%) $p<0.01$], need for HFNC [Group 1: 45.7% vs Group 2: 37.8% vs Group 3: 33.3% vs Group 4: 31.1%; $p<0.01$], PICU admission [Group 1: 29.5% vs Group 2: 19.4% vs Group 3: 20% vs Group 4 :16.6%; $p<0.01$] and NMV [Group 1: 27.2% vs Group 2: 18.9% vs Group 3: 15.8% vs Group 4 :15.7%;

$p < 0.01$]. Figure 1. This results were consistent in sensitivity analyses using the economic vulnerability index. (Table 1). There were no deaths during the study period

In logistic regression analysis, infants with lower household income had higher risk of admission to the PICU [OR=1.90 (1.13-3.16), $p=0.01$], HFNC [OR: 1,59 (1,04-2,42); $p=0,02$ and NMV [OR: 1,81 (1,08-3,09); $p=0,02$]

This study provides data associating lower household income with higher severity of RSV bronchiolitis in terms of PICU admission, need of HFNC, NMV and Wood-Downes score severity. More vulnerable infants living in lower income areas were at higher risk of admission to the PICU. This association is probably multifactorial, since infants living in more disadvantaged areas with lower income are associated with higher exposure to tobacco smoke and lower breastfeeding rates, both risk factors for severe RSV infection^{13, 14}.

It is widely reported that poverty can impede access to medical care and hinder the prevention of certain diseases, affecting life expectancy in adulthood¹⁵⁻¹⁷. Recently, it has been published in the United Kingdom that children living in low socioeconomic communities are at greater risk of developing severe persistent asthma, and are more likely to be admitted to hospital¹⁸. In 2020, Esteban y Peña et al, published that some health issues in the City of Madrid are intimately related to social and economic attributes, such as obesity, smoking, or sedentary lifestyles. They suggest implementing targeted policies for prevention and health promotion in vulnerable populations¹⁹.

In contrast to our study, Zheng et al published in 2020 in children admitted for bronchiolitis that a higher mean family income was associated with a higher risk of

admission to the PICU⁷. In this multicenter study, unlike our study, data were collected through a survey that up to 29% of the participants refused to answer. In addition, the socioeconomic context and health care in the United States and Spain are very different, which may at least partly explain these differences. Recently, a study conducted in Tennessee using data collected by the RSV Hospitalization Surveillance Network from 2016 to 2023 demonstrated that RSV hospitalization were linked to socially vulnerable census tracts ⁶.

Therefore, it is essential to have a good understanding of the social characteristics of infants with RSV bronchiolitis in order to ensure that new preventive interventions with long half-life monoclonal antibodies (nirsevimab) or bivalent pre-Fusion maternal RSV vaccine reach the most vulnerable and low-income infants ^{9, 10, 20}.

This study has some limitations; it is a retrospective and single-center study. Family income has been estimated according to the zip code, which can sometimes give an inaccurate income. Our hospital is a referral center and receives patients with severe RSV bronchiolitis from other hospitals, which may bias the results.

In conclusion, the social determinants of health and specifically the mean income associated with the zip code may have an important influence on the clinical course of RSV bronchiolitis.

Key words: RSV severity bronchiolitis, socioeconomic status, inequity, social health determinants, zip code

Financiación

Este estudio ha sido parcialmente financiado por la beca FIS PI21/00840

Conflicto de intereses

RRF declara haber impartido charlas relacionadas con la bronquiolitis por VRS financiadas por Pfizer, Sanofi, Astra Zeneca, y MSD. Ha participado en Advisory boards financiados por Pfizer, Merck, Moderna y Sanofi. Ha recibido becas para asistencia a conferencias y congresos financiado por Pfizer y Sanofi


REFERENCES

1. Lozano R, Naghavi M, Foreman K, et al. Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet (London, England)*. 2012;380:2095-2128.
2. Hall CB, Simões EA, Anderson LJ. Clinical and epidemiologic features of respiratory syncytial virus. *Current topics in microbiology and immunology*. 2013;372:39-57.
3. Rodríguez-Fernández R, González-Sánchez MI, Pérez-Moreno J, et al. Age and respiratory syncytial virus etiology in bronchiolitis clinical outcomes. *The journal of allergy and clinical immunology. Global*. 2022;1:91-98.
4. Kachmar AG, Wypij D, Perry MA, Curley MAQ. Income-driven socioeconomic status and presenting illness severity in children with acute respiratory failure. *Research in nursing & health*. 2021;44:920-930.
5. Foley D, Best E, Reid N, Berry MMJ. Respiratory health inequality starts early: The impact of social determinants on the aetiology and severity of bronchiolitis in infancy. *Journal of paediatrics and child health*. 2019;55:528-532.
6. Thomas CM, Raman R, Schaffner W, et al. Respiratory Syncytial Virus Hospitalizations Associated With Social Vulnerability by Census Tract: An Opportunity for Intervention? *Open forum infectious diseases*. 2024;11:ofae184.
7. Zheng DX, Mitri EJ, Garg V, et al. Socioeconomic Status and Bronchiolitis Severity Among Hospitalized Infants. *Academic pediatrics*. 2020;20:348-355.
8. Jansson L, Nilsson P, Olsson M. Socioeconomic environmental factors and hospitalization for acute bronchiolitis during infancy. *Acta paediatrica (Oslo, Norway : 1992)*. 2002;91:335-338.
9. Drysdale SB, Cathie K, Flamein F, et al. Nirsevimab for Prevention of Hospitalizations Due to RSV in Infants. *The New England journal of medicine*. 2023;389:2425-2435.
10. Kampmann B, Madhi SA, Munjal I, et al. Bivalent Prefusion F Vaccine in Pregnancy to Prevent RSV Illness in Infants. *The New England journal of medicine*. 2023;388:1451-1464.

11. Ferrés Mataró J, Mangues Bafalluy MA, Farré Riba R, Juliá Brugues A, Bonal de Falgas J. [Subcutaneous adrenaline versus inhaled salbutamol in the treatment of childhood asthmatic crisis]. *An Esp Pediatr.* 1987;27:37-40.
12. R Core Team (2018). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>; 2018.
13. Maedel C, Kainz K, Frischer T, Reinweber M, Zacharasiewicz A. Increased severity of respiratory syncytial virus airway infection due to passive smoke exposure. *Pediatric pulmonology.* 2018;53:1299-1306.
14. Mineva GM, Purtill H, Dunne CP, Philip RK. Impact of breastfeeding on the incidence and severity of respiratory syncytial virus (RSV)-associated acute lower respiratory infections in infants: a systematic review highlighting the global relevance of primary prevention. *BMJ global health.* 2023;8.
15. Chandrasekhar R, Sloan C, Mitchel E, et al. Social determinants of influenza hospitalization in the United States. *Influenza and other respiratory viruses.* 2017;11:479-488.
16. Hussain T, van den Berg S, Ziesemer KA, Markhorst DG, Vijverberg SJH, Kapitein B. The influence of disparities on intensive care outcomes in children with respiratory diseases: A systematic review. *Pediatric pulmonology.* 2023; 1-9. doi:10.1002/ppul.26629
17. Beamer PI, Lothrop N, Lu Z, et al. Spatial clusters of child lower respiratory illnesses associated with community-level risk factors. *Pediatric pulmonology.* 2016;51:633-642.
18. Pinot de Moira A, Taylor-Robinson D. Social Inequalities in Asthma: The Cold Facts. *Archivos de bronconeumologia.* 2023;59:791-792.
19. Esteban YPMM, Fernández Velasco E, Jiménez García R, Hernández Barrera V, Fernandez Del Pozo I. [Health, incidence and differences in the territorial vulnerability city of Madrid]. *Revista española de salud pública.* 2020;94.
20. Ares-Gómez S, Mallah N, Santiago-Pérez MI, et al. Effectiveness and impact of universal prophylaxis with nirsevimab in infants against hospitalisation for respiratory syncytial virus in Galicia, Spain: initial results of a population-based longitudinal study. *The Lancet. Infectious diseases.* 2024 Aug; 24(8):817-828. doi: 10.1016/S1473-3099(24)00215-9.

Figure 1 caption. The figure shows four groups of infants admitted for RSV bronchiolitis according to their family income by zip code and their differences in PICU hospitalizations. Asterisks indicate statistical significance

FIGURE 1



Journal Pre

Table 1.- Differences according socioeconomic status and economic vulnerability index in hospitalized RSV infections in infants

Variable	Average household income					Economic vulnerability index		
	20-29.999 euros (Group 1)	30-39.999 euros (Group 2)	40-49.999 euros (Group 3)	>50.000 euros (Group 5)	p ¹	High Index (>5,32)	Low index (≤ 5,32)	p ²
Age (months)	3 (1,7-7)	2,8 (1,3-7)	3 (1,5-7)	3,5 (1,5-8)	0,40	3,5 (1,5-6,5)	3 (1,5-7)	0,47
Sex (male)	176/302 (58,3%)	206/397 (51,9%)	70/120 (58,3%)	112/235 (47,7%)	0,06	231/391 (59,1%)	226/465 (48,6%)	0,04
Breastfeeding	168/264 (63,6%) [#]	260/361 (72%)	89/114 (78,1%) [#]	167/216 (77,3%) [#]	<0,01	244/355 (68,7%)	322/431 (74,7%)	0,06
Second hand smoke	90/248 (36,3%) [#]	120/324 (37%) [#]	27/97 (27,8%)	37/197 (18,8%) [#]	<0,01	113/319 (35,4%)	107/393 (27,2%)	0,02
Day care attendance	17/254 (6,7%) [#]	31/344 (9%)	14/107 (13,1%)	30/212 (14,2%) [#]	0,03	32/342 (9,4%)	46/412 (11,2%)	0,47
Preterm birth	59/302 (19,5%)	55/397 (13,9%)	18/120 (15%)	43/235 (18,3%)	0,18	59/391 (15,1%)	68/465 (14,6%)	0,84
Underlying condition	64/302 (21,2%)	61/397 (15,4%)	28/120 (23,3%)	46/235 (19,6%)	0,11	73/391 (18,7%)	74/465 (19,1%)	0,31
Severity score								
-Mild	20/302 (6,6%)	35/397 (8,8%)	7/120 (5,8%)	13/235 (5,5%)		27/391 (6,9%)	38/465 (8,2%)	
-Moderate	153/302 (50,7%)	223/397 (56,2%)	78/120 (65%)	154/235 (65,5%)	0,01	221/391 (56,5%)	299/465 (64,3%)	0,01
-Severe	129/302(42,7%) [#]	139/397 (35%)	35/120 (29,2%)	68/235 (28,9%) [#]		143/391 (36,6%)	128/465 (27,5%)	
Respiratory rate	50 (41-60)	50 (44-60)	50 (42-60)	50 (42-60)	0,20	50 (42-60)	50 (42-60)	0,15
Heart rate	156 (140-170)	156 (140-170)	154,5 (140-170)	154 (138-170)	0,82	156 (140-170)	155 (139-170)	0,34

LOS (days)	5 (3-7)	5 (3-8)	5 (3-7)	5 (3-7)	0,29	5 (3-8)	5 (3-7)	0,16
Enteral feeding	99/302 (32,8%) [#]	82/397 (20,7%) [#]	23/120 (19,2%) [#]	45/235 (19,1%) [#]	<0,01	105/391 (26,9%)	71/465 (15,3%)	<0,01
Apnea	32/302 (10,6%)	55/397 (13,9%)	18/120 (15%)	24/235 (10,2%)	0,32	45/391 (11,5%)	49/465 (10,5%)	0,66
Pneumonia in X-Ray	57/302 (18,9%)	70/397 (17,6%)	16/120 (13,3%)	39/235 (21,4%)	0,45	59/457 (15,1%)	78/391 (16,8%)	0,10
HFNC	138/302 (45,7%) [#]	150/397 (37,8%)	40/120 (33,3%)	73/235 (31,1%) [#]	<0,01	153/391 (39,1%)	143/465 (30,8%)	0,01
PICU	89/302 (29,5%) [#]	77/397 (19,4%) [#]	24/120 (20%)	39/235 (16,6%) [#]	<0,01	96/391 (24,6%)	58/465 (12,5%)	<0,01
NMV	82/302 (27,2%) [#]	75/397 (18,9%)	19/120 (15,8%)	37/235 (15,7%) [#]	<0,01	87/391 (22,3%)	56/465 (12%)	<0,01
MV	10/302 (3,3%)	10/397 (2,5%)	2/120 (1,7%)	4/235 (1,7%)	0,67	11/391 (2,8%)	5/465 (1,2%)	0,07

Abbreviations: LOS: length of stay, HFNC: high flow nasal cannula, PICU: Pediatric intensive care unit, NMV: Non-mechanical ventilation, MV: Mechanical ventilation.

Continuous variables were analyzed using the U-Mann-Whitney or Kruskal-Wallis tests and categorical variables using Fisher's exact or Chi-square test. Data are reported as medians, 25%-75% interquartile range.

Bold highlights the significant.

[#] Indicates significance in analysis post hoc after Chi-square test with Bonferroni correction (Specifically, there are differences between groups 1-3 and 1-4 in terms of breastfeeding, there are differences between groups 1-4 and 2-4 in terms of passive exposure to tobacco smoke, and there are differences between groups 1-4 in daycare attendance. Regarding the severity score, there are differences between groups 1-4 in the % of severe bronchiolitis. Regarding PICU there are differences between groups 1-2 and, 1-4, in HFNC between groups 1-4, in NMV between groups 1-4 and in enteral feeding between groups 1-2, 1-3 and 1-4)

p¹: Comparison according average household income

p²: Comparison according economic vulnerability index

Journal Pre-proof