

## Original Article

## Montelukast, Leukotriene Inhibitor, Reduces LPS-Induced Acute Lung Inflammation and Human Neutrophil Activation



Jaime Eduardo Davino-Chiovatto<sup>a,◇</sup>, Manoel Carneiro Oliveira-Junior<sup>b,◇</sup>, BreAnne MacKenzie<sup>b,◇</sup>, Alana Santos-Dias<sup>b</sup>, Ana Roberta Almeida-Oliveira<sup>b</sup>, Jefferson Comin Jonco Aquino-Junior<sup>b</sup>, Auriléia Aparecida Brito<sup>a</sup>, Nicole Cristine Rigonato-Oliveira<sup>a</sup>, Nilsa Regina Damaceno-Rodrigues<sup>c</sup>, Ana Paula Ligeiro Oliveira<sup>a</sup>, Alessandro Pereira Silva<sup>d</sup>, Fernanda Marciano Consolim-Colombo<sup>a</sup>, Flavio Aimbire<sup>e</sup>, Hugo Caire Castro-Faria-Neto<sup>f</sup>, Rodolfo Paula Vieira<sup>b,g,h,i,\*</sup>

<sup>a</sup> Nove de Julho University (UNINOVE), São Paulo, SP, Brazil

<sup>b</sup> Brazilian Institute of Teaching and Research in Pulmonary and Exercise Immunology (IBEPIPE), São José dos Campos, SP, Brazil

<sup>c</sup> University of São Paulo, School of Medicine, Department of Pathology (LIM 59), São Paulo, SP, Brazil

<sup>d</sup> Post-graduation Program in Biomedical Engineering, University of Mogi das Cruzes, Mogi das Cruzes, SP, Brazil

<sup>e</sup> Federal University of São Paulo (UNIFESP), São José dos Campos, SP, Brazil

<sup>f</sup> Laboratory of Immunopharmacology, Osvaldo Cruz Institute (IOC), Osvaldo Cruz Foundation, Rio de Janeiro, RJ, Brazil

<sup>g</sup> Universidade Brasil, Post-graduation Program in Bioengineering and in Biomedical Engineering, São Paulo, SP, Brazil

<sup>h</sup> Federal University of São Paulo (UNIFESP), Post-graduation Program in Sciences of Human Movement and Rehabilitation, Santos, SP, Brazil

<sup>i</sup> Anhembi Morumbi University, School of Medicine, Avenida Deputado Benedito Matarazzo 4050, São José dos Campos, SP, Brazil

## ARTICLE INFO

## Article history:

Received 3 July 2018

Accepted 1 May 2019

## Keywords:

Inflammation  
Neutrophils  
Acute lung injury  
Leukotriene  
Cytokines  
Growth factors

## ABSTRACT

**Objectives:** Some pro-inflammatory lipids derived from 1 lipooxygenase enzyme are potent neutrophil chemoattractant, a cell centrally involved in acute respiratory distress syndrome (ARDS); a syndrome lacking effective treatment. Considering the beneficial effects of the leukotriene receptor inhibitor, montelukast, on other lung diseases, whether montelukast attenuates inflammation in a mouse model of ARDS, and whether it reduces LPS stimulated activation of human neutrophils was investigated.

**Methods:** Thirty-five C57Bl/6 mice were distributed into control (PBS) + 24 h, LPS + 24 h (10 µg/mouse), control + 48 h, LPS + 48 h, and LPS 48 h + Montelukast (10 mg/kg). In addition, human neutrophils were incubated with LPS (1 µg/mL) and treated with montelukast (10 µM).

**Results:** Oral-tracheal administration of montelukast significantly attenuated total cells ( $P < .05$ ), macrophages ( $P < .05$ ), neutrophils ( $P < .01$ ), lymphocytes ( $P < .001$ ) and total protein levels in BAL ( $P < .05$ ), as well as IL-6 ( $P < .05$ ), CXCL1/KC ( $P < .05$ ), IL-17 ( $P < .05$ ) and TNF- $\alpha$  ( $P < .05$ ). Furthermore, montelukast reduced neutrophils ( $P < .001$ ), lymphocytes ( $P < .01$ ) and macrophages ( $P < .01$ ) in the lung parenchyma. In addition, montelukast restored BAL VEGF levels ( $P < .05$ ). LTB4 receptor expression ( $P < .001$ ) as well as NF- $\kappa$ B ( $P < .001$ ), a downstream target of LPS, were also reduced in lung parenchymal leukocytes. Furthermore, montelukast reduced IL-8 ( $P < .001$ ) production by LPS-treated human neutrophils.

**Conclusion:** In conclusion, montelukast efficiently attenuated both LPS-induced lung inflammation in a mouse model of ARDS and in LPS challenged human neutrophils.

© 2019 SEPAR. Published by Elsevier España, S.L.U. All rights reserved.

### El montelukast, un inhibidor de leucotrienos, reduce la inflamación pulmonar aguda inducida por LPS y la activación de neutrófilos humanos

## RESUMEN

**Objetivos:** Algunos lípidos proinflamatorios derivados de la enzima lipooxigenasa 1 son potentes quimioatrayentes de neutrófilos, un tipo celular con una implicación principal en el síndrome de distrés

## Palabras clave:

Inflamación

\* Corresponding author.

E-mail address: [rodrelena@yahoo.com.br](mailto:rodrelena@yahoo.com.br) (R.P. Vieira).

◇ These authors are equally contributed to this study.

Neutrófilos  
Daño pulmonar agudo  
Leucotrieno  
Citoquinas  
Factores de crecimiento

respiratorio agudo (SDRA), para el que no hay tratamiento efectivo. Considerando los efectos beneficiosos del inhibidor de los receptores de leucotrienos montelukast en otras enfermedades pulmonares, se investigó si este fármaco era capaz de atenuar la inflamación en un modelo de ratón de SDRA y de reducir la activación de los neutrófilos humanos inducida por LPS.

**Métodos:** Se utilizaron 35 ratones C57BL/6 distribuidos en los siguientes grupos: control (PBS) + 24 h, LPS + (24 h [10 µg/ratón]), control + 48 h y LPS 48 h + montelukast (10 mg/kg). Por otro lado, se incubaron neutrófilos humanos con LPS (1 µg/ml) y se trataron con montelukast (10 µM).

**Resultados:** La administración orotraqueal de montelukast redujo el número total de células ( $p < 0,05$ ), de macrófagos ( $p < 0,05$ ), de neutrófilos ( $p < 0,01$ ), de linfocitos ( $p < 0,001$ ) y los niveles totales de proteína en el lavado broncoalveolar ( $p < 0,05$ ), así como de IL-6 ( $p < 0,05$ ), CXCL1/KC ( $p < 0,05$ ), IL-17 ( $p < 0,05$ ) y TNF- $\alpha$  ( $p < 0,05$ ). Además, el montelukast redujo los neutrófilos ( $p < 0,001$ ), los linfocitos ( $p < 0,01$ ) y los macrófagos ( $p < 0,01$ ) en el parénquima pulmonar. Asimismo, restauró los niveles de VEGF en el lavado broncoalveolar ( $p < 0,05$ ) y disminuyó la expresión del receptor LTB4 ( $p < 0,001$ ) y de NF- $\kappa$ B ( $p < 0,001$ ), una diana *downstream* del LPS, en los leucocitos del parénquima pulmonar. Por último, redujo la producción de IL-8 por parte de los neutrófilos humanos tratados con LPS.

**Conclusión:** En conclusión, el montelukast atenuó de manera eficaz tanto la inflamación pulmonar inducida por LPS en un modelo de ratón de SDRA como en neutrófilos humanos estimulados con LPS.

© 2019 SEPAR. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

## Introduction

Acute respiratory distress syndrome (ARDS) is characterized by hypoxemic respiratory failure associated with acute pulmonary inflammation and edema presenting high mortality rates.<sup>1</sup> Different etiologies, such as head, chest and other major injuries, as well as sepsis, inhalation of harmful substances and severe pneumonia may result in the development of ARDS.<sup>1–3</sup> Although the mechanisms underlying the pathophysiology of ARDS are not completely understood, in all cases; especially in bacterial infections, an exacerbated inflammatory response plays a central role.<sup>1–3</sup>

While several animal models have been established to study ARDS, lipopolysaccharide (LPS) is the most widely used as it reproduces several important ARDS features, such as the accrual of neutrophils in alveolar and in interstitial space and in bronchoalveolar lavage (BAL), the accumulation of proteinaceous debris in alveolar spaces, thickening of the alveolar wall, and increased concentration of total proteins and pro-inflammatory cytokines in BAL.<sup>2–5</sup>

LPS signals via toll like receptor 4 (TLR4) and activates nuclear factor kappa B (NF- $\kappa$ B), a key regulator of the inflammatory process.<sup>3–6</sup> NF- $\kappa$ B is a transcription factor regulating several aspects of ARDS pathophysiology, such as production of pro-inflammatory cytokines (i.e. IL-1 $\beta$ , IL-6, IL-8/CXCL-1 and TNF- $\alpha$ ), and also the lung fibroproliferative response in murine models of ARDS.<sup>7–10</sup> NF- $\kappa$ B also regulates human and mouse fibroblast differentiation,<sup>11</sup> a key event that occurs during fibroproliferation, an important process following lung injury. Therefore, pharmacological approaches that inhibit NF- $\kappa$ B expression and activation may attenuate lung inflammatory and fibrotic responses following injury.<sup>3–5,10–12</sup>

Cys leukotrienes receptor-1 (cysLTR1) antagonists montelukast, pranlukast and zafirlukast are small molecules that have demonstrated secondary, off-target, anti-inflammatory effects including the inhibition of cyclic nucleotides phosphodiesterases and 5'-lypoxygenase as well as NF- $\kappa$ B downregulation.<sup>13</sup>

Taken together, this study tested the hypothesis that montelukast inhibits both acute lung injury induced by LPS in mice and LPS-induced human neutrophil activation.

## Results

### Montelukast Reduces Leukocyte Number in Bronchoalveolar Lavage (BAL)

Similarly to the leukocyte response during ARDS, oral-tracheal administration of LPS results in a significant increase in the

number of lung leukocytes. At 24 h, BAL leukocytes were significantly increased in mice injured with LPS (Fig. 1A). To mimic an ARDS therapeutic scenario, mice received oral-tracheal administration of montelukast 24 h post-LPS injury and were sacrificed 24 h later. At 48 h, total number of BAL cells remained significantly increased, however, montelukast significantly reduced the number of cells in the BAL (Fig. 1B). At 24 h post-LPS treatment, differential cell counts of BAL fluid revealed a significant increase in the number of macrophages, neutrophils and lymphocytes compared to PBS treated controls (Fig. 1C, E, G). Montelukast treatment at 24 h, significantly attenuated the accumulation of macrophages (Fig. 1D), neutrophils (Fig. 1F), and lymphocytes at 48 h (Fig. 1H).

### Montelukast Reduces Vascular Permeability and Cytokines in Bronchoalveolar Lavage (BAL)


Compared to PBS treated mice, increased vascular permeability was evident by increased protein concentration in BAL fluid at 24 h post-LPS treatment (Fig. 2A) and remained at 48 h (Fig. 2B), while Montelukast treatment at 24 h post-LPS reduced the total amount of BAL proteins (Fig. 2B). At 24 h post-LPS treatment, ELISA detected increased levels of proinflammatory cytokines IL-6 (Fig. 2C), CXCL1/KC (Fig. 2E), TNF- $\alpha$  (Fig. 2G) and IL-17 (Fig. 2I). At 48 h post-LPS treatment, cytokines remained high: IL-6 (Fig. 2D), CXCL1/KC (Fig. 2F), TNF- $\alpha$  (Fig. 2H) and IL-17 (Fig. 2J), but were effectively reduced by therapeutic administration of Montelukast.

### Montelukast Reduces Neutrophils, Lymphocytes and Macrophages Accumulation in the Lung Parenchyma

Indeed, at 24 and 48 h post-LPS treatment, morphometric analysis revealed increased numbers of parenchymal neutrophils (Fig. 3A, B), lymphocytes (Fig. 3C, D), and macrophages (Fig. 3E, F). Treatment with Montelukast significantly reduced the number of immune cells in the lung parenchyma (Fig. 3B, D, F). Representative images of PBS (Fig. 3G), LPS + 24 h (Fig. 3I), LPS + 48 h (Fig. 3H), and 96 LPS + Montelukast (Fig. 3J).

### Montelukast Inhibits NF- $\kappa$ B and Leukotriene B4 Receptor (LTB4R) Expression in the Lung Parenchyma

In the present study, increased expression of NF- $\kappa$ B by parenchymal leukocytes for both periods studied (24 h and 48 h) post-LPS stimulation compared with PBS groups (Fig. 4A and B)



**Fig. 1.** Therapeutic administration of montelukast reduced leukocyte number in bronchoalveolar lavage (BAL). Results are expressed as mean  $\pm$  SEM. For (A and B) \*  $P < .05$  compared with PBS48 h and PBS48h + MK group. For (C) \*\* $P < .01$  compared with PBS48 h and LPS48h + MK group. For (D) \*\*\* $P < .001$  compared with PBS48 h group and LPS48h + MK group.

was observed. The results also revealed that Montelukast efficiently reduced the NF- $\kappa$ B expression (Fig. 4B). LTB<sub>4</sub> signals mainly via LTB<sub>4</sub>R, which results in the recruitment of neutrophils. The present study demonstrated that LPS stimulation at 24 h and 48 h resulted in increased expression of LTB<sub>4</sub>R by parenchymal leukocytes (Fig. 4C and D) while Montelukast significantly reduced the LTB<sub>4</sub>R expression (Fig. 4D). Fig. 4E–H shows representative photomicrographs of NF- $\kappa$ B for PBS48 h, LPS24 h, LPS48 h and LPS48 h + montelukast groups, respectively.

#### Montelukast Restores VEGF Expression in LPS-injured Lungs

Treatment with LPS decreased VEGF expression at 24 h (Fig. 5A) and furthermore at 48 h while Montelukast administration at 24 h resulted in a recovery of VEGF expression in BAL compared to control (PBS) treated animals (Fig. 5B).

#### Montelukast Reduces IL-8 Release by Human Neutrophils

Human neutrophils were stimulated with LPS (1.5  $\mu$ g/mL) for 1 h and then incubated with Montelukast (10  $\mu$ M in PBS) for an additional 5 h. ELISA for IL-8 was performed on the supernatant. Montelukast treatment significantly inhibited LPS-stimulated production of IL-8 (Fig. 6A).


#### Discussion

The present study demonstrates for the first time that leukotriene inhibitor Montelukast reduces both acute LPS-induced lung inflammation in mice as well as LPS-induced human neutrophils activation. Exacerbated inflammation plays a central role in the pathogenesis of ARDS; therefore in the quest to develop effective ARDS therapies, reducing inflammation has been a main goal. The off-target effects of Montelukast on LTB<sub>4</sub> receptors present on neutrophils could have beneficial implications in

ARDS therapy. Leukotriene B<sub>4</sub> (LTB<sub>4</sub>) is synthesized primarily by activated basophils, eosinophils, monocytes and macrophages and acts in both an autocrine and paracrine manner by signaling to structural cells, neutrophils and TH<sub>2</sub> lymphocytes.<sup>14</sup> Though neutrophils express a low level of LTB<sub>4</sub> receptor, LTB<sub>4</sub> acts as a strong chemoattractant of neutrophils, a cell centrally involved in ARDS.

While the recognition of the involvement of leukotriene pathways in asthma prompted the development of leukotriene receptor inhibitors, drugs such as Montelukast, a leukotriene inhibitor, have not yet been tested in the context of ARDS.<sup>15</sup> Twenty-five years ago, a swine LPS model of acute lung injury (ALI) demonstrated that the LTBR1 competitive receptor antagonist LY255283 reduced ALI.<sup>16</sup> However, unlike Montelukast, this agent did not exhibit potent off target anti-inflammatory effects such as the inhibition of cyclooxygenase or 5-lipoxygenase enzymes.<sup>17</sup> Therefore, in the case of ARDS, Montelukast may be an effective inhibitor of inflammation not only because of its ability to inhibit leukotriene signaling, but also via off-target effects which result in a net increase in cyclic adenosine monophosphate (cAMP),<sup>18</sup> and the suppression of NF- $\kappa$ B<sup>19</sup> which leads to the attenuation of cytokine production and a dispersal of lung immune cells.

This study used the LPS-induced acute lung injury model in mice to test the hypothesis that the Montelukast would attenuate lung inflammation. Our results indicated that Montelukast reduced the LPS-induced increase in total immune cells both in the BAL, suggesting decreased vascular permeability, and in the lung parenchyma. In addition, Montelukast treated mice displayed significantly reduced levels of cytokines IL-6, CXCL1/KC, IL-17 and TNF- $\alpha$  suggesting attenuation of inflammatory processes due to LPS. LPS-induced pulmonary dysfunction was associated with increased neutrophil count, leukotriene (LT) B<sub>4</sub>, and tumor necrosis factor (TNF)- $\alpha$  in BALF. These results suggest that treatment with Montelukast can be useful in chronic airway inflammatory diseases including COPD poorly responsive to glucocorticoids.<sup>20</sup> In addition, Montelukast treated mice also displayed reduced LTB<sub>4</sub>R




**Fig. 2.** Therapeutic administration of montelukast reduced vascular permeability and cytokine production in BAL. The results are expressed as mean  $\pm$  SEM. For (A–E) \*  $P < .05$  compared with PBS48 h and LPS48 h + MK group.

and NF- $\kappa$ B expression in the lung parenchyma, which correlates to decreased leukotriene signaling and decreased inflammation and Montelukast was able to block LTD(4)-induced stimulation. Allergen challenge leads to a significant increase in sCD14 concentrations in BAL and might modulate the allergen-induced inflammation. In addition, LTD(4) might play a role in the release of sCD14, and it could be speculated that sCD14 reduction by LTRA might contribute to the mechanisms of LTRA in the treatment of allergic asthma.<sup>21</sup> Moreover, LTB(4)- and LTD(4)-induced eosinophil activation was attenuated by CP-105,696 and the Cys-LT(1) receptor antagonist montelukast, respectively, highlighting specific receptor dependency.<sup>6</sup> Thus, mediator-triggered granulocyte activation and antiapoptotic pathways are distinct events that can be differentially regulated.<sup>22</sup> Lastly, our results presented herein indicates that Montelukast treatment resulted in a recovery of VEGF expression in the BAL, a measurement which is associated with recovery in ARDS patients.<sup>23–25</sup>

While neutrophils themselves do not produce LTB<sub>4</sub>, and LTB<sub>4</sub> receptors are expressed only at low levels, exposure to LTB<sub>4</sub> primes neutrophils to produce copious amounts of reactive oxygen species (ROS), matrix metalloproteinases (MMPs) and other cytokines upon stimulation by additional cytokines.

Human neutrophils activated with the chemoattractant N-formyl-L-methionyl-L-leucyl-L-phenylalanine (fMLP) in combination with cytochalasin B resulted in abrupt and sustained increases in cytosolic Ca<sup>2+</sup>(+), as well as release of elastase and production of superoxide and LTB<sub>4</sub>, and expression of macrophage complement receptor 3 (CR3). These effects were attenuated with Montelukast treatment and the literature point out its association with significant increases in cyclic AMP.<sup>26</sup> In this study, human neutrophils were stimulated with LPS which signals via TLR4 and activates NF- $\kappa$ B-signaling resulting in the production of cytokines, particularly IL-8.<sup>27</sup> Stimulation with LPS for 1 h, followed by one treatment of Montelukast resulted in a significant reduction of IL-8 in the cell supernatant. Given the low level of LTBR4 receptor expression by neutrophils, this study also suggests that an important off-target effect of Montelukast is the ability to attenuate NF- $\kappa$ B signaling, likely through upstream mechanisms that increase intracellular cAMP.

An interesting point that should be highlighted is the fact that our results showed herein a reduction of circulating levels of IL-17 in Montelukast-treated mice with acute lung inflammation. Therefore, from our results, is possible suggest that the Th17 cells could be a bystander mediator of Montelukast effect in vivo, since



**Fig. 3.** Therapeutic administration of montelukast reduced neutrophils, macrophages and lymphocytes in the lung parenchyma. Results are expressed as mean  $\pm$  SEM. For (A and B) \*\*\*  $P < .001$  compared with PBS48 h and LPS48 h + MK group. For (C) \*\*\*  $P < .001$  compared with PBS48 h and \*\*  $P < .01$  compared with LPS48 h + MK group. (D, E and F) are representative photomicrographs of HE lung stained slides in PBS48 h, LPS48 h and LPS48 h + MK groups, respectively.

that IL-17 is produced by Th17 cells, and this subset expresses high levels of LTB<sub>4</sub>R1 and CysTLR1, being attracted by leukotrienes (especially LTD<sub>4</sub>).<sup>28</sup>

In conclusion, this is the first study to report the effects of Montelukast in an LPS mouse model of ARDS and in LPS-stimulated human neutrophils activation. These results concur not only the potent anti-inflammatory nature of Montelukast but also its ability to signal via LTB<sub>4</sub>R independent pathways. Future studies should further explore the nature of these mechanisms especially in the context of human neutrophils, a central inflammation mediator of ARDS.

## Material and methods


### Experimental Design of Animal Experiments

This study was approved by the ethical committee of Nove de Julho University (AN0002/2015) and were carried out in accordance to Guide for the Care and Use of Laboratory Animals, published by the U.S. National Institutes of Health (NIH


publication no. 85-23, revised 1996). Thirty-eight C57Bl/6 male mice weighing between 20 and 25 g were distributed in Control 24 h (PBS24 h;  $n = 7$ ), LPS 24 h (LPS24 h;  $n = 7$ ), Control 48 h (PBS48 h;  $n = 8$ ), LPS 48 h (LPS48 h;  $n = 8$ ) and LPS 48 h + montelukast (LPS 48 h + MK;  $n = 8$ ). For Montelukast administration, animals were anesthetized by intra-peritoneal injection of ketamine (100 mg/kg) and xylazine (10 mg/kg). The PBS24 h and LPS24 h groups were euthanized 24 h after vehicle (PBS 50  $\mu$ l) or LPS (10  $\mu$ g/mouse/50  $\mu$ l PBS) administration. The PBS48 h, LPS48 h and LPS 48 h + MK groups were euthanized 48 h after vehicle (PBS; 50  $\mu$ l) or LPS (10  $\mu$ g/mouse/50  $\mu$ l PBS) administration. Montelukast (10  $\mu$ M/mouse/50  $\mu$ l PBS) was orotracheally administered 24 h after LPS administration and animals were euthanized 24 h later.

### Assessment of Lung Inflammation


Lung inflammation was assessed through the collection and analysis of bronchoalveolar lavage (BAL). Quantitative analysis of parenchymal inflammation was performed using histomorphometrical technique.<sup>29–31</sup> The numbers of total and differential cells



**Fig. 4.** NF-κB and LTB4R expression by leukocytes in lung parenchyma were reduced by therapeutic montelukast administration. Results are expressed as mean ± SEM. For (A) \*\*\*  $P < .001$  compared with PBS48h group and LPS48h + MK group. For (B) \*\*\*  $P < .001$  compared with PBS48 h and \*  $P < .05$  LPS48h + MK group. (C, D and E) are representative photomicrographs of immunohistochemistry for NF-κB in PBS48h, LPS48h, LPS48h and LPS48h + montelukast groups, respectively.



**Fig. 5.** Therapeutic montelukast administration resulted in a recovery of VEGF levels in lung tissue homogenate. Results are expressed as mean ± SEM. For this figure \*\*  $P < .01$  compared with LPS48 h group and \*  $P < .05$  compared with PBS48h + MK group.



**Fig. 6.** Montelukast suppressed IL-8 levels production by human neutrophils stimulated with LPS. IL-8 levels in neutrophil supernatant were measured using ELISA. The results are expressed as mean ± SEM. For this figure \*\*\*  $P < .001$  compared with non-stimulated (Medium) and LPS-stimulated and treated with montelukast (LPS + MK group).

in BAL were evaluated in the material recovered from 3 gentle washes of 0.5 mL of sterile PBS, by using a Neubauer chamber (total cells) and cytopsin preparations (differential cell count).<sup>29–31</sup> The cytopsin were stained with Diff Quick (Medion Diagnostics, Düringen, Switzerland), as previously described.<sup>27–32</sup> In summary,

300 cells per slide per mouse were analyzed using an optical microscope and were counted according to the classical hematological criteria.<sup>29–34</sup>

The density of neutrophils, lymphocytes and macrophages in the lung parenchyma was evaluated as previously described.<sup>29–31</sup> Briefly, the lungs were excised in block, fixed in 10% formalin solution, at a constant pressure (20 cmH<sub>2</sub>O) for 24 h, and submitted to histological routine.<sup>33</sup>

Five micrometer lung slices were stained with hematoxylin and eosin and 20 photomicrographs at 400× magnification of each animal of all groups were obtained using an Olympus BX43-L-FL microscope with a camera XM-10. The area of lung tissue in the lung parenchyma was obtained by subtracting the airspace area from the total photo area. Then, the number of neutrophils, lymphocytes and macrophages were counted in each photo according to the morphological criteria. The results were expressed in number of cells/mm<sup>2</sup> of lung tissue.<sup>30–33</sup>

#### Total Proteins in BAL

The levels of total proteins in BAL was measured using the BCA Protein Assay Kit (Thermo Scientific, USA) and was used as an index of vascular permeability.<sup>31</sup>

#### Cytokines in BAL, Serum and in Cell Culture Supernatants

The levels of IL-6 (Biolegend, Code 431306; Detection limit 7.8 pg/mL), CXCL1/KC (R&D Systems, Code DY453; Detection limit 15.6 pg/mL), IL-8 (Biolegend, Code 431506; Detection limit 15.6 pg/mL), IL-17 (R&D Systems, Code DY421; Detection limit 15.6 pg/mL) and TNF- $\alpha$  (Biolegend, Code 430906; Detection limit 7.8 pg/mL) in BAL and in cell culture supernatants were measured using ELISA kits from Biolegends (USA) and R&D Systems (USA) as indicated, according to the manufacturers' recommendations.<sup>33</sup> The measurements were done in triplicates in all samples.

#### Immunohistochemical Study

The quantitative analysis of the expression of leukotriene B<sub>4</sub> receptor (LTB<sub>4</sub>R) (sc-98841; Santa Cruz, CA, USA) and of NF- $\kappa$ B p65 (sc-109; Santa Cruz, CA, USA) by parenchymal leukocytes was performed using classical immunohistochemistry protocol, as previously described.<sup>30,33,34</sup> Briefly, the area of lung tissue in the lung parenchyma was obtained by subtracting the airspace area from the total photo area. Then, the number of leukocytes positive to LTB<sub>4</sub>R and NF- $\kappa$ B p65 were counted in each photo. The results were expressed in number of positive cells/mm<sup>2</sup> of lung tissue.<sup>29–32</sup>

#### Isolation and Culture of Human Neutrophils

Eight milliliter of peripheral blood was diluted 1:1 in sterile PBS and added to a tube containing Ficoll Paque gradient and submitted to 20 min centrifugation (1200 × g). Red cells were lysed, polymorphonuclear cells collected and neutrophils were separated from eosinophils using EasySep™ Human Neutrophil Enrichment Kit (#19257; StemCell Technologies, USA). The purity was assessed through cyto-spin analysis and was higher than 98%. Neutrophils (1 × 10<sup>6</sup>/2 mL medium/well) were incubated in 48 well plates in RPMI 1640 medium and were stimulated with LPS (1.5  $\mu$ g/mL medium) for 1 h and then incubated with montelukast (10  $\mu$ M in PBS) for an additional 5 h. Supernatant was recovered for measurements of IL-8 by ELISA.

#### Statistical Analysis

If not stated otherwise, two-way analysis of variance (TWO-WAY ANOVA) followed by Bonferroni post hoc test was used.

Significance levels were considered for  $P < .05$ . Values were expressed as mean  $\pm$  SEM.

#### Authorship

JEDC, MCOJ, BM, AD, ARAO, JCJAJ, AAB, NCRO, NRDR, contributed performing the experiments and analysis. APLO, APS, FMCC, FA, HCCFN, and RPV have written the manuscript and critically revised the manuscript and performed the statistical analysis. RPV have designed the study. All authors have reviewed and approved the final version of the manuscript prior to submission.

#### Acknowledgements

This study was supported by Sao Paulo Research Foundation (FAPESP), grant 2012/15165-2. MCOJ holds a PhD fellowship from FAPESP (2014/14604-8). BM holds a post-doctoral fellowship from FAPESP (2014/23196-0). ARAO holds a MSc fellowship from FAPESP (2014/07500-1). JCJAJ holds a MSc fellowship from FAPESP (2014/12755-9). APB holds a MSc fellowship from CAPES. NCRO holds a PhD fellowship from CAPES.

#### References

- Ranieri VM, Rubenfeld GD, Thompson BT, Ferguson ND, Caldwell E, Fan E, et al. Acute respiratory distress syndrome: the Berlin Definition. *JAMA*. 2012;307:2526–33.
- Matute-Bello G, Downey G, Moore BB, Groshong SD, Matthay MA, Slutsky AS, et al. An official American Thoracic Society workshop report: features and measurements of experimental acute lung injury in animals. *Am J Respir Cell Mol Biol*. 2011;44:725–38.
- Guo L, Li S, Zhao Y, Qian P, Ji F, Qian L, et al. Silencing Angiopoietin-Like Protein 4 (ANGPTL4) protects against lipopolysaccharide-induced acute lung injury via regulating SIRT1/NF- $\kappa$ B pathway. *J Cell Physiol*. 2015;230:2390–402.
- Lin MH, Chen MC, Chen TH, Chang HY, Chou TC. Magnolol ameliorates lipopolysaccharide-induced acute lung injury in rats through PPAR $\gamma$ -dependent inhibition of NF- $\kappa$ B activation. *Int Immunopharmacol*. 2015;28:270–8.
- Meng F, Meliton A, Moldobaeva N, Mutlu G, Kawasaki Y, Akiyama T, et al. Asef mediates HGF protective effects against LPS induced lung injury and endothelial barrier dysfunction. *Am J Physiol Lung Cell Mol Physiol*. 2015;308:L452–63.
- Lu YC, Yeh WC, Ohashi PS. LPS/TLR4 signal transduction pathway. *Cytokine*. 2008;42:145–51.
- Christman JW, Venkatakrisnan A, Wheeler AP, Bernard GR, Blackwell TS. NF- $\kappa$ B binding activity in mixed Bal cells from patients with SIRS and ARDS. *Sepsis*. 2008;4:35–41.
- Caamano J, Hunter CA. NF-B family 317 of transcription factors: central regulators of innate and adaptive immune functions. *Clin Microbiol Rev*. 2002;15:414–29.
- Hoesel B, Schmid JA. The complexity of NF- $\kappa$ B signaling in inflammation and cancer. *Mol Cancer*. 2013;2:12–86.
- Lawrence T. The nuclear factor NF- $\kappa$ B pathway in inflammation. *Cold Spring Harb Perspect Biol*. 2009;1:a001651.
- Sun X, Chen E, Dong R, Chen W, Hu Y. Nuclear factor (NF)- $\kappa$ B p65 regulates differentiation of human and mouse lung fibroblasts mediated by TGF $\beta$ . *Life Sci*. 2015;122:8–14.
- Avasarala S, Zhang F, Liu G, Wang R, London SD, London L. Curcumin modulates the inflammatory response and inhibits subsequent fibrosis in a mouse model of viral-induced acute respiratory distress syndrome. *PLoS ONE*. 2013;8:e57285.
- Theron AJ, Steel HC, Tintinger GR, Gravett CM, Anderson R, Feldman C. Cysteinyl leukotriene receptor-1 antagonists as modulators of innate immune cell function. *J Immunol Res*. 2014;2014:608930.
- Latz E, Visintin A, Lien E, Fitzgerald KA, Monks BG, Kurt-Jones EA, et al. Lipopolysaccharide rapidly traffics to and from the Golgi apparatus with the toll-like receptor 4-MD-2-CD14 complex in a process that is distinct from the initiation of signal transduction. *J Biol Chem*. 2002;277:47834–43.
- Cheng H, Leff JA, Amin R, Gertz BJ, De Smet M, Noonan N, et al. Pharmacokinetics, bioavailability, 358 and safety of montelukast sodium (MK-0476) in healthy males and females. *Pharm Res*. 1996;13:445–8.
- Wollert PS, Menconi MJ, O'Sullivan BP, Wang H, Larkin V, Fink MP, et al. LY255283, a novel leukotriene B<sub>4</sub> receptor antagonist, limits activation of neutrophils and prevents acute lung injury induced by endotoxin in pigs. *Surgery*. 1993;114:191–8.
- Silbaugh SA, Stengel PW, Cockerham SL, Roman CR, Saussy DL Jr, Spaethe SM, et al. Pulmonary actions of LY255283, a leukotriene B<sub>4</sub> receptor antagonist. *Eur J Pharmacol*. 1992;13:57–64.
- Flamand N, Surette ME, Picard S, Bourgoin S, Borgeat P. Cyclic AMP mediated inhibition of 5-lipoxygenase translocation and leukotriene biosynthesis in human neutrophils. *Mol Pharmacol*. 2002;62:250–6.

19. Takahashi N, Tetsuka T, Uranishi H, Okamoto T. Inhibition of the NFκB transcriptional activity by protein kinase A. *Eur J Biochem.* 2002;269:4559–65.
20. Abdel Kawy HS. Montelukast versus dexamethasone treatment in a guinea pig model of chronic pulmonary neutrophilic inflammation. *COPD.* 2016;13:455–63.
21. Julius P, Grosse-Thie C, Kuepper M, Bratke K, Virchow JC. sCD14 in bronchoalveolar lavage 18, 42 and 162 hours after segmental allergen provocation. *Scand J Immunol.* 2010;71:304–11.
22. Murray J, Ward C, O'Flaherty JT, Dransfield I, Haslett C, Chilvers ER, et al. Role of leukotrienes in the regulation of human granulocyte behaviour: dissociation between agonist-induced activation and retardation of apoptosis. *Br J Pharmacol.* 2003;139:388–98.
23. Medford ARL, Millar AB. Vascular endothelial 337 growth factor (VEGF) in acute lung injury (ALI) and acute respiratory distress syndrome (ARDS): paradox or paradigm? *Thorax.* 2006;61:621–6.
24. Maitre B, Boussat S, Jean D, Gouge M, Brochard L, Housset B, et al. Vascular endothelial growth factor synthesis in the acute phase of experimental and clinical lung injury. *Eur Respir J.* 2001;18:100–6.
25. Thickett DR, Armstrong L, Millar AB. A role for vascular endothelial growth factor in acute and resolving lung injury. *Am J Respir Crit Care Med.* 2002;166:1332–7.
26. Gravett CM, Theron AJ, Steel HC, Tintinger GR, Cockeran R, Feldman C, et al. Interactive inhibitory effects of formoterol and montelukast on activated human neutrophils. *Eur Respir J.* 2010;36:1417–24.
27. Harada A, Sekido N, Akahoshi T, Wada T, Mukaida N, Matsushima K, et al. Essential involvement of interleukin-8 (IL-8) in acute inflammation. *J Leukoc Biol.* 1994;56:559–64.
28. Lee W, Su Kim H, Lee GR. Leukotrienes induce the migration of Th17 cells. *Immunol Cell Biol.* 2015;93:472–9.
29. Ramos DS, Olivo CR, Quirino Santos Lopes FD, Toledo AC, Martins MA, Lazo Osório RA, et al. Low-intensity swimming training partially inhibits lipopolysaccharide-induced acute lung injury. *Med Sci Sports Exerc.* 2010;42:113–9.
30. Reis GCT, Reis GCG, de Almeida FM, Lopes FD, dos Santos DAC, dos Santos FA, et al. Protective effects of aerobic exercise on acute lung injury induced by LPS in mice. *Crit Care.* 2012;18:R199.
31. Oliveira MC Jr, Greiffo FR, Rigonato-Oliveira NC, Custódio RW, Silva VR, Damaceno-Rodrigues NR, et al. Low level laser therapy reduces acute lung inflammation in a model of pulmonary and extrapulmonary LPS-induced ARDS. *J Photochem Photobiol B.* 2014;5:57–63.
32. Baudiß K, Ayata CK, Lazar Z, Cicko S, Beckert J, Meyer A, et al. Ceramide-1-phosphate inhibits cigarette smoke-induced airway inflammation. *Eur Respir J.* 2015;45:1669–80.
33. Vieira RP, de Andrade VF, Duarte AC, Dos Santos AB, Mauad T, Martins MA, et al. Aerobic conditioning and allergic pulmonary inflammation in mice II. Effects on lung vascular and parenchymal inflammation and remodeling. *Am J Physiol Lung Cell Mol Physiol.* 2008;295:L670–9.
34. Vieira RP, Toledo AC, Silva LB, Almeida FM, Damaceno-Rodrigues NR, Caldini EG, et al. Anti-inflammatory effects of aerobic exercise in mice exposed to air pollution. *Med Sci Sports Exerc.* 2012;44:1227–34.