

## Original Article

## Aerobic Capacity of Preterm Children with Very Low Birth Weight at School Age and its Associated Factors

 Sabrina Pinheiro Tsopanoglou<sup>a</sup>, Josy Davidson<sup>a,b,\*</sup>, Victor Zuniga Dourado<sup>a</sup>, Ana Lucia Goulart<sup>a</sup>, Marina Carvalho de Moraes Barros<sup>a</sup>, Amélia Miyashiro Nunes dos Santos<sup>a</sup>
<sup>a</sup> Universidade Federal de São Paulo – UNIFESP, São Paulo, Brazil

<sup>b</sup> Centro Universitário São Camilo, São Paulo, Brazil

## ARTICLE INFO

## Article history:

Received 13 March 2019

Accepted 26 May 2019

## Keywords:

 Preterm children  
 Very low birth weight  
 Aerobic capacity

## ABSTRACT

**Introduction:** Respiratory morbidities of preterm infants can cause significant ventilatory impairment thus compromising the aerobic capacity in childhood and adolescence. Therefore, the present study was conducted to evaluate the aerobic capacity in school age preterm children with VLBW and its associated factors.

**Methods:** A cross-sectional study was conducted among preterm born with VLBW and term children, both aged 6–9 years. An individualized symptom-limited treadmill testing protocol performed aerobic capacity. Measured variables: oxygen pulse (PuO<sub>2</sub>), percentage of maximum heart rate for age (%HR max), tidal volume/inspiratory capacity ratio (TV/IC), oxygen consumption (VO<sub>2</sub>) peak, and the ratio of the anaerobic threshold of gas exchange to the predicted percentage of maximum VO<sub>2</sub> (VO<sub>2</sub>@LA/%VO<sub>2</sub> max.pred.) were compared between groups. Univariate and multiple linear regression analyses were used to determine the factors associated with aerobic capacity.

**Results:** Thirty-four preterm and 32 term children were included. Similar VO<sub>2</sub> peak and the other variables were observed. The development of bronchopulmonary dysplasia (BPD) and being obese/overweight was positively associated with %HR max. The Z-score for height/age and birth weight <1000 g was positively associated with PuO<sub>2</sub> and peak VO<sub>2</sub>, and negatively associated with overweight/obesity and female sex. **Conclusions:** Aerobic capacity was similar in both groups. Sex, development of BPD, birth weight <1000 g and factors related to body growth, such as Z-score for height/age and overweight/obesity, were associated with aerobic capacity in preterm children with VLBW.

© 2019 SEPAR. Published by Elsevier España, S.L.U. All rights reserved.

### Capacidad aeróbica de niños prematuros de muy bajo peso al nacer y edad escolar y sus factores asociados

## RESUMEN

**Introducción:** Las enfermedades respiratorias de los niños prematuros pueden causar importantes impedimentos ventilatorios que comprometen la capacidad aeróbica en la infancia y en la adolescencia. El presente estudio se llevó a cabo para evaluar la capacidad aeróbica de niños prematuros en edad escolar de muy bajo peso al nacer (BPN) y los factores asociados.

**Métodos:** Se llevó a cabo un estudio transversal con niños prematuros de muy BPN y con niños a término, ambos grupos con edades comprendidas entre los 6 y 9 años. Las siguientes variables se compararon entre los 2 grupos: el pulso de oxígeno (PuO<sub>2</sub>), el porcentaje de frecuencia cardíaca máxima (%FC máx.), la relación entre el volumen corriente y la capacidad inspiratoria (TV/IC), el consumo pico de oxígeno (VO<sub>2</sub>) y la relación entre el umbral anaeróbico de intercambio de gas y el porcentaje estimado de VO<sub>2</sub> máximo (VO<sub>2</sub>@LA/%VO<sub>2</sub> máx. pred.). Se llevaron a cabo análisis de regresión lineal univariante y multivariante para determinar los factores asociados con la capacidad aeróbica.

## Palabras clave:

 Niños prematuros  
 Muy bajo peso al nacer  
 Capacidad aeróbica

\* Corresponding author.

E-mail address: [josy davidson@yahoo.com.br](mailto:josy davidson@yahoo.com.br) (J. Davidson).

**Resultados:** Se incluyeron 34 niños prematuros y 32 niños a término. Se registraron valores similares de  $VO_2$  pico y de otras variables. El desarrollo de displasia broncopulmonar (BPD) y de obesidad/sobrepeso mostró una asociación positiva con el %FC máx. El Z-score para la altura/edad y el peso al nacer < 1.000 g se asoció positivamente con la  $SAO_2$  y el  $VO_2$  y negativamente con el sobrepeso/obesidad y el sexo femenino. **Conclusiones:** La capacidad aeróbica fue similar entre los 2 grupos. El sexo, el desarrollo de BPD, peso al nacer < 1.000 g y factores relacionados con el crecimiento corporal, tales como el Z-score para la altura/edad y para el sobrepeso/obesidad se asociaron con la capacidad aeróbica en niños prematuros de muy BPN.

© 2019 SEPAR. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

## Introduction

Respiratory morbidities remain the most common problem among preterm infants<sup>1</sup> despite the recent improvements in post-natal care.<sup>2</sup> Among respiratory morbidities, bronchopulmonary dysplasia (BPD) is the most important impairment and its pulmonary structural alterations and ventilatory and gas exchange limitations could decrease muscle strength<sup>3–5</sup> and then, functional exercise capacity in early childhood and adolescence.<sup>6,7</sup>

During exercise, one of the main factors that determine exercise capacity is the maximum oxygen uptake ( $VO_2$  peak). Studies on adults and adolescents born extremely preterm have shown that peak  $VO_2$  values may be within the normal range.<sup>4,8</sup> However, some authors have reported that decreased pulmonary diffusion capacity or altered pulmonary function may compromise the aerobic capacity in preterm children and adolescents, with a decreased peak  $VO_2$  or distance covered being observed on the cardiopulmonary exercise test (CPET) that is performed on a treadmill or cycle ergometer.<sup>8–10</sup> Hence, the actual aerobic capacity and the associated factors with it in school age preterm children with very low birth weight (VLBW) remain controversial.

Therefore, the present study was conducted to evaluate the aerobic capacity in school age preterm children with VLBW and its associated factors.

## Methods

A cross-sectional study was conducted on two groups of 6–9-year-old children of both sex, recruited between February 2013 and April 2015. The preterm group consisted of children born with gestational age of <37 weeks and birth weight of <1500 g, accompanied at the multidisciplinary outpatient follow-up clinic of the institution. Term group consisted of children born with gestational age of  $\geq 37$  weeks, both sexes, without respiratory impairment or outpatient follow-up, and being relatives or friends of premature children attended in our outpatient clinic. Children with congenital malformations; neuromuscular diseases; neurological, visual, and/or hearing disorders; and/or other problems that may lead to difficulties on the execution of the test were excluded. Children with signs or symptoms of acute or chronic respiratory disease in the last 2 weeks or those with abnormal hemodynamic condition at rest were also excluded.<sup>11</sup>

This study followed the declaration of Helsinki. The Ethics and Research Committee of the Institution approved this study (Protocol: 173.275/12). The consent form was signed by the parents or guardians of the children, and the assent form was signed by the children.

Included children were evaluated according to a clinical questionnaire answered by their parents, and anthropometric variables were evaluated before cardiopulmonary exercise test. Additionally, the clinical records of preterm group during the neonatal period and their clinical status after discharge from the neonatal unit were also collected. In the term group, neonatal data were obtained from the birth card complemented by an interview with parents and/or legal guardians conducted by the researcher. The

socioeconomic status and frequency of weekly physical activity of the child were evaluated through a questionnaire answered by the parents or guardians on the scheduled day of the test.<sup>12</sup> The child's Body Mass Index (BMI) during the recruitment was determined according to the recommendations of the World Health Organization.<sup>13</sup>

A cardiopulmonary exercise test (CPET) and a symptom-limited treadmill test with a rapid incremental protocol and individualized increase in load was used to evaluate the aerobic capacity.<sup>14</sup> The slope was calculated individually<sup>15</sup> based on the fixed velocity of 3.5 mph (children aged between 6 and 7 years) and 4.0 mph (those aged between 8 and 9 years) and predicted maximum  $VO_2$  ( $VO_2$  max). For each age group,  $VO_2$  max was calculated according to child's sex and weight.<sup>16</sup>

Prior to the test, the child remained seated for 5 min, and then vital signs (blood pressure, heart rate, and peripheral oxygen saturation) and degree of dyspnea were evaluated. The non-rebreathing valve face mask (Hans Rudolph, Kansa) of the gas analyzer (K4b2, Cosmed, Pavnadi Albano, Itália) was placed on the face of the child, and the device analyzer was placed in front of the treadmill in order not to impose a workload during the CPET. The gas analyzer was previously calibrated at the beginning of the tests according to the manufacturer's recommendations.

The CPET was started with 3-min warm-up phase<sup>17</sup> maintaining the speed of 1.0 mph without grade and during the last minute, the speed was raised according to the child's age. During the exercise phase, the grade was increased by 2% every minute until a maximum grade of 10%. If at this grade the child did not meet the maximum effort criteria,<sup>14,18</sup> the speed was increased by 0.5 mph every minute until meeting the interruption criteria. In the recovery phase, the child remained walking for 5 min in the warm-up speed and grade.

During the test, expired gases were collected breath-by-breath, and data recording was filtered every 15 s.<sup>19</sup> Variables were analyzed for two periods: at rest and at the peak of the CPET, considering the values during the last 15–30 s at the end of the test, based on arithmetic means.

The anaerobic threshold (AT) was obtained using the V-slope method<sup>16</sup> after the concordance of two previously trained observers. The interpretation of anaerobic threshold of gas exchange ( $VO_2@LA$ ) was obtained based on the relationship between anaerobic threshold of gas exchange and percentage of maximum oxygen consumption predicted according to age ( $VO_2@LA/\% VO_2$  max.pred.).

To assess the aerobic capacity, the following cardiovascular, respiratory, and metabolic variables were analyzed: oxygen pulse ( $PuO_2$ ) obtained by the ratio  $\Delta FC/\Delta VO_2$ ; %HR max; tidal volume and inspiratory capacity ratio (TV/IC peak);  $VO_2$  peak, and the relationship between anaerobic threshold of gas exchange and percentage of maximum oxygen consumption predicted according to age ( $VO_2@LA/\% VO_2$  max.pred.).<sup>16,19,20</sup>

In order to consider the maximum CPET, the child was required to meet three of the following criteria: maximum HR  $\geq 90\%$  of the predicted HR according to age; signs of exhaustion, gaseous exchange ratio (R) of >1.0, and  $VO_2$  maximum of  $\geq 85\%$ .<sup>14,18</sup> In




Fig. 1. Descriptive flowchart of the total sample.

addition, CPET was discontinued if some of the following interurrences were identified: request for discontinuation by the child, failure of the monitoring device, pulse oxygenation decrease of  $\geq 4\%$  compared to the resting phase.<sup>16,21</sup>

The same protocol was used in all children, and a pulmonologist and/or pediatrician was present in the laboratory during the tests, together with the researchers of the study.

#### Statistical analysis

Sample size (G-Power 3.0.10) was calculated according to  $VO_2$  difference of 7.3 mL/kg/min<sup>14</sup> between groups, showed the need of 22 participants in each group, assuming an alpha error of 0.05 and beta of 0.80.

Numerical variables were compared using Student's *t* test (normal distribution) or Mann–Whitney test (non-normal distribution). Categorical variables were compared using the chi-square test or Fisher's exact test. The normality of variables was evaluated using the Kolmogorov–Smirnov test.

Regarding the analysis of factors associated with children's aerobic capacity, univariate and multivariate linear regressions were used considering the cardiovascular, respiratory, and metabolic variables related to an individual's aerobic capacity as dependent variables. Variables with statistical ( $p < 0.2$ ) and clinical significance in the univariate regression were included in the multiple linear regression models, excluding collinear variables in the same model. Statistical analyses were performed using the SPSS for Win/v.17.0 program (IBM SPSS Statistics, Somers, NY), and  $p < 0.05$  was considered statistically significant.

#### Results

During the study period, 95 preterm children were attended in our outpatient clinic, however 33 (34.7%) of them did not meet inclusion criteria. Then, 62 preterm children were eligible and 34 term children (relatives or preterm children's friends) were invited to the study but at the end of the study 34 preterm and 32 term children finished the exercise test (Fig. 1).

The 22 (35.5%) eligible preterm children (but not included) were compared to 34 included preterm children, and no difference in the demographic and clinical characteristics was observed.

The mean birth weight and gestational age were  $3072 \pm 609$  g and  $38.5 \pm 1.4$  weeks in the term group, and  $1131 \pm 228$  g and  $29.8 \pm 2.5$  weeks in the preterm group. Clinical and demographic characteristics of both groups were similar, except for the lower Z score height for age and an increased need for hospitalization in the preterm group (Table 1).

During the CPET, all children reached the maximum speed and grade recommended in the protocol, and in some cases, the speed

Table 1

Clinical and demographic characteristics of the sample.

	Preterm group (n = 34)	Term group (n = 32)	p value
Age (years) <sup>a</sup>	8.0 ± 1.0	7.9 ± 1.1	0.773
Female (%) <sup>c</sup>	17 (50.0%)	14 (43.8%)	0.631
Baseline weight (kg) <sup>a</sup>	27.2 ± 6.5	28.4 ± 7.6	0.499
Z score weight for age <sup>a</sup>	0.14 ± 1.44	0.43 ± 1.26	0.389
Height (cm) <sup>a</sup>	125.7 ± 7.7	128.3 ± 9.2	0.211
Z score height for age <sup>a</sup>	−0.35 ± 1.17	0.24 ± 0.96	0.032
BMI (kg/cm <sup>2</sup> ) <sup>a</sup>	17.0 ± 2.6	17.0 ± 2.8	0.944
BMI (Z-score) <sup>b</sup>	0.44 ± 1.3	0.40 ± 1.2	0.710
Overweight, n (%) <sup>d</sup>	8 (23.5%)	4 (12.5%)	0.342
Obesity, n (%) <sup>d</sup>	5 (14.7%)	4 (12.5%)	1.000
Hospitalization after NICU discharge, n (%) <sup>d</sup>	14 (45.2%)	5 (15.6%)	0.014
Physical activity (h/week) <sup>b</sup>	3.3 ± 1.4	3.8 ± 1.5	0.190

BMI: body mass index.

<sup>a</sup> *t* student.

<sup>b</sup> Mann–Whitney.

<sup>c</sup> Chi-square.

<sup>d</sup> Exact Fisher test.

Table 2

Cardiovascular, respiratory, and metabolic variables.

	Preterm group (n = 34)	Term group (n = 32)	p value
PuO <sub>2</sub> (mL/bpm) <sup>a</sup>	6.0 ± 1.4	6.2 ± 1.4	0.535
%HR max <sup>b</sup>	87.9 ± 4.1	86.1 ± 5.9	0.244
TV/IC peak (L) <sup>b</sup>	0.615 ± 0.146	0.592 ± 0.131	0.735
VO <sub>2</sub> peak (mL/min) <sup>a</sup>	1085 ± 273	1102 ± 277	0.798
VO <sub>2</sub> peak (mL/kg/min) <sup>a</sup>	40.4 ± 6.5	39.5 ± 6.4	0.541
VO <sub>2</sub> @LA/%VO <sub>2</sub> max.pred. (%) <sup>a</sup>	67.8 ± 18.1	69.0 ± 10.7	0.738

PuO<sub>2</sub>: oxygen pulse, expressed by the VO<sub>2</sub>/HR ratio; HR: heart rate; %HR max: percentage of maximum heart rate of age; TV: tidal volume; IC: inspiratory capacity; VO<sub>2</sub> peak: maximal oxygen uptake; VO<sub>2</sub>@LA: VO<sub>2</sub> at anaerobic threshold; %VO<sub>2</sub> max.pred.: percentage of predicted VO<sub>2</sub> max. p value:

<sup>a</sup> *t* student.

<sup>b</sup> Mann–Whitney.

was also increased meeting all the criteria for the maximum CPET. Despite this, the speed variation of the treadmill was similar between groups (preterm group:  $23.5 \pm 13.5\%$  vs. term:  $26.0 \pm 9.6\%$ ,  $p = 0.312$ ). All children presented sinus rhythm on the electrocardiogram from the beginning until the recovery phase.

Aerobic pattern was similar between groups according to cardiovascular, respiratory, and metabolic variables, as described in Table 2.

Multiple linear regression analyses for possible factors associated with aerobic capacity are shown in Tables 3–5. The %HR max was positively associated to the dependence of O<sub>2</sub> for more than 28 days and overweight/obesity. Oxygen pulse was positively associated to Z score height/age and negatively to female sex. In the same

**Table 3**

Multiple regression analysis of factors associated with the percentage of maximum heart rate for age (%HR max).

	Beta	95% CI	p
Oxygen dependence at 28 days of life	3.153	0.244 to 6.063	0.034
Overweight/obese	2.628	-0.181 to 5.438	0.066

Controlled model for birth weight of <1000 g, respiratory distress syndrome, oxygen dependence at 28 days of life, duration of mechanical ventilation, days of hospitalization, overweight/obese. Model significance,  $p = 0.006$ ,  $r^2 = 0.165$ .

**Table 4**

Multiple regression analysis of factors associated with oxygen pulse (PuO<sub>2</sub>).

	Beta	95% CI	p value
Female	-0.683	-1.266 to -0.100	0.022
Z score height for age	0.712	0.447 to 0.977	<0.001

Controlled model for sex, Z score height for age, and overweight/obese. Model significance,  $p < 0.001$ ,  $r^2 = 0.338$ .

**Table 5**

Multiple regression analysis of factors associated with VO<sub>2</sub> peak (mL/kg/min).

	Beta	95% CI	p
Female	-4.326	-0.337 to -3.111	0.003
Birth weight of <1000 g	4.690	0.257 to 2.370	0.021
Overweight/obese	-5.624	-0.409 to -3.837	<0.001

Controlled model for sex, Z score height for age, overweight/obesity, birth weight of <1000 g. Model significance,  $p < 0.001$ ,  $r^2 = 0.300$ .

way, VO<sub>2</sub> peak (mL/kg/min) was negatively associated to female and overweight/obese but positively to birth weight < 1000 g.

The following respiratory and metabolic parameters, the relationship between VC/IC peak, VO<sub>2</sub> in absolute values, and VO<sub>2</sub> LA/%VO<sub>2</sub> max.pred., did not show significant association in the univariate linear regression; thus, they were not included in the multiple regression analysis.

## Discussion

The present study showed that children aged between 6 and 9 years with VLBW had similar aerobic capacity to children born at term. Nevertheless, after adjusting the covariates, these parameters were associated with the history of bronchopulmonary dysplasia, birth weight of <1000 g, sex and factors related to body growth, such as Z score height for age and overweight/obesity.

The results in the univariate analysis demonstrate that at school age, preterm children had an aerobic capacity similar to that of full-term children. Some hypotheses may explain this similarity as evidenced by the fact that the included preterm children were born after undergoing measures to reduce mortality and neonatal morbidity, such as the use of exogenous surfactant, antenatal corticosteroids, and gentle ventilation modes.<sup>22</sup> These measures have certainly improved the development and lung growth of preterm infants, reducing the severity of pulmonary complications such as bronchopulmonary dysplasia.<sup>23</sup> Another hypothesis would be the low frequency of physical activity of the children in both groups, which may have influenced their aerobic responses.<sup>24</sup> These children had a very low physical activity (approximately 3.5 h/week in both groups) when compared to the recommended physical activity for this group age (60 min daily) to prevent sedentary lifestyle.<sup>24</sup> Additionally, the final hypothesis is the possibility of these preterm children to be like adults with chronic obstructive pulmonary disease, which increases the work of respiratory muscles during exercise to maintain or reach their aerobic capacity, i.e., within the normal range.<sup>25</sup>

Although the results of physical capacity seemed similar between groups, other factors were important according to the

multivariate regression analysis. The presence of bronchopulmonary dysplasia increased by 3% maximum heart rate for age. This association could be explained by reduced ventilatory reserve and decreased lung diffusion capacity in preterm children, which can be supplied by increased cardiovascular system demand during exercise, represented by the increase in the %HR max.<sup>2,10</sup> Moreover, children with history of pulmonary disease during the neonatal period have greater degree of respiratory impairment. Maybe, if we have had a greater sample of preterm children with bronchopulmonary dysplasia, the difference in cardiopulmonary conditions could possibly be showed.

Other important neonatal variable was associated with physical capacity as lower weight of birth. Birth weight of <1000 g increased the VO<sub>2</sub> peak in approximately 4 mL/kg/min. It is known that preterm infants, especially the extremely ones may present air trapping during exercise, which involves pulmonary mechanics in an indirect way.<sup>5</sup> In the present study, premature birth or BPD were not associated with any ventilatory variable, such as VC/IC peak, however, we cannot affirm that preterm infants do not present ventilatory or pulmonary mechanics alterations during exercise, since these variables were not evaluated by spirometry, being a limitation of the present study.

In addition to factors inherent to prematurity, sex and factors related to body growth, such as Z-score for height/age and overweight/obesity seemed to be associated with cardiovascular and metabolic conditions during exercise. The nutritional status, more specifically, being obese or overweight, reduced the VO<sub>2</sub> peak by approximately 6 mL/kg/min and increased by about 3% %HR for age. Zwirner<sup>26</sup> describes that the peak VO<sub>2</sub> expectedly increases based on an individual's growth; however, when excessive body weight gain occurs, as in the case of overweight or obesity, the association may be negative because the percentage of body fat is higher than body growth.<sup>27</sup> Our results are consistent with the study of Welsh et al.<sup>8</sup> that studied preterm children aged between 10 and 11 years by cycloergometer test. They also showed that impaired body growth was associated with decreased aerobic capacity. Our results show the importance of a good monitoring of nutritional status, especially in relation to obesity and overweight during follow-up of these premature infants during childhood.

Another important factor that presented a positive association was oxygen pulse. Despite this association, it did not clinically alter the PuO<sub>2</sub> behavior in these children. This association was expected as studies had demonstrated the association between increased VO<sub>2</sub> peak and body growth. As already known, PuO<sub>2</sub> is dependent on VO<sub>2</sub>, which were expected to increase together.<sup>26,28</sup>

Finally, the last factor associated with cardiovascular and metabolic parameters was sex, which demonstrated that being female reduced the PuO<sub>2</sub> and VO<sub>2</sub> peak (mL/kg/min). As previously demonstrated, PuO<sub>2</sub> is dependent on the VO<sub>2</sub> peak, as girls had lesser PuO<sub>2</sub>, may be associated with the lower VO<sub>2</sub> peak presented.<sup>26,27</sup>

Ventilatory reserve expressed as TV/IC peak ratio is expected to be associated with some independent variables because of the presence of BPD or prematurity itself, as previously reported in the literature.<sup>2,4,8</sup> This association was expected because preterm infants may present pulmonary hyperinflation, which would reduce the tidal volume and inspiratory capacity to exercise. Then, preterm infants in their study sample possibly did not present significant pulmonary sequelae due to BPD or prematurity per se.

Regarding metabolic parameters, absolute values of VO<sub>2</sub> and VO<sub>2</sub>@LA/%VO<sub>2</sub> max.pred. ratio<sup>4</sup> studies had shown controversial results, which were expected to be not associated with preterm parameters.<sup>2,28,29</sup>

The varied results presented in our study and the literature can be explained by the inclusion criteria. Most studies included heterogeneous population<sup>4</sup> and different types of protocols and

ergometers that could also influence the results.<sup>4,9</sup> Thus, including children with homogeneous age group and similar growth and developmental characteristics as in our study could increase the internal validity and reliability of the presented data, because studies have shown that sexual maturation may play an important role in the performance of exercise capacity in preterm children.<sup>30</sup> Although the groups presenting statistical difference in the Z score height/age ratio, their mean values are within normal limits, showing the similarity of growth and development between groups. The application of an individualized CPET loading protocol in the treadmill also favored the validity of results, because this type of ergometer protocol can better identify the peak VO<sub>2</sub> in children.<sup>17</sup> Aside from this factor, controlling the weekly physical activity of children, which would be a possible confounding factor, increased the quality of results in this study.

The percentage of loss and the number of children that had not met criteria to participate to the study could be a limitation in this study and it is possible that if we had a larger sample we could reach more precise conclusions. However, the analysis of the main demographic and clinical characteristics comparing the included and not included preterm children showed no significant difference, especially in relation to variables that could be biased, such as BPD and VLBW. Another limitation was the inclusion of a convenience sample, excluding preterm children with severe motor and cognitive sequelae. Those with severe neurodevelopmental impairments are also possibly the extremely preterm ones and with higher occurrence of bronchopulmonary dysplasia, and thus they could show a decreased aerobic capacity. However, these patients cannot be included due to the characteristics required in the protocol.

## Conclusions

Preterm children with VLBW at school age showed aerobic capacity similar to that of full-term children. However, the intensity of responses during the cardiopulmonary exercise test was associated with sex, BPD, lower birth weight and factors related to body growth, demonstrating the influence of neonatal factors and body growth on the aerobic capacity in these children.

## Funding

This work was supported by the Brazilian Government Scholarship—CAPES—Coordenação de Aperfeiçoamento de Pessoal de Nível Superior, in which Sabrina Pinheiro Tsopanoglou was supported by as a postgraduate student.

## Authorship

Tsopanoglou S.P., Davidson J., Dourado V.Z., Goulart A.L., Barros M.C.M., Santos A.M.N.: the conception and design of the study; acquisition of data; analysis and interpretation of data; drafting the article; revising it critically for important intellectual content and final approval of the version to be submitted.

## Conflict of interests

The authors declare no conflict of interests.

## Acknowledgments

Sabrina Pinheiro Tsopanoglou was supported by the Brazilian Government Scholarship—CAPES—Coordenação de

Aperfeiçoamento de Pessoal de Nível Superior, as a postgraduate student.

## References

- Schmalisch G, Wilitzki S, Roehr CC, Proquitté H, Bühner C. Differential effects of immaturity and neonatal lung disease on the lung function of very low birth weight infants at 48–52 postconceptional weeks. *Pediatr Pulmonol.* 2013;48:1214–23.
- Clemm HH, Vollsæter M, Røksund OD, Eide GE, Markestad T, Halvorsen T. Exercise capacity after extremely preterm birth. Development from adolescence to adulthood. *Ann Am Thorac Soc.* 2014;11:537–45.
- Er I, Gunlemez A, Uyan ZS, Aydogan M, Oruc M, Isik O, et al. Evaluation of lung function on impulse oscillometry in preschool children born late preterm. *Pediatr Int.* 2016;58:274–8.
- Kaplan E, Bar-Yishay E, Prais D, Klinger G, Mei-Zahav M, Mussaffi H, et al. Encouraging pulmonary outcome for surviving, neurologically intact, extremely premature infants in the postsurfactant era. *Chest.* 2012;142:725–33.
- Maiorana A, O’Dea CA, Banton GL, Wilson AC, Logie K, Pillow JJ, et al. Increased prevalence of expiratory flow limitation during exercise in children with bronchopulmonary dysplasia. *ERJ Open Res.* 2018. <http://dx.doi.org/10.1183/23120541.00048-2018>.
- Smith LJ, van Asperen PP, McKay KO, Selvadurai H, Fitzgerald DA. Reduced exercise capacity in children born very preterm. *Pediatrics.* 2008;122:e287–93.
- Tsopanoglou SP, Davidson J, Goulart AL, Barros MC de M, dos Santos AMN. Functional capacity during exercise in very-low-birth-weight premature children. *Pediatr Pulmonol.* 2014;49:91–8.
- Welsh L, Kirkby J, Lum S, Odendaal D, Marlow N, Derrick G, et al. The EPICure study: maximal exercise and physical activity in school children born extremely preterm. *Thorax.* 2010;65:165–72.
- Vollsæter M, Clemm HH, Satrell E, Eide GE, Røksund OD, Markestad T, et al. Adult respiratory outcomes of extreme preterm birth. A regional cohort study. *Ann Am Thorac Soc.* 2015;12:313–22.
- Kilbride HWMD, Gelatt MCM, Sabath RJE. Pulmonary function and exercise capacity for ELBW survivors in preadolescence: effect of neonatal chronic lung disease. *J Pediatr.* 2003;143:488–93.
- National High Blood Pressure Education Program Working Group on High Blood Pressure in Children and Adolescents. The fourth report on the diagnosis, evaluation, and treatment of high blood pressure in children and adolescents. *Pediatrics.* 2004;114 Suppl., 4th report:555–76.
- Pardini R, Matsudo S, Matsudo V, Andrade E, Braggion G, Andrade D. Validação do questionário internacional de nível de atividade física (IPAQ – versão 6): estudo piloto em adultos jovens brasileiros. *Rev Bras Ciên Mov Bras.* 2001;9:45–51.
- World Health Organization. The Child Growth Standards; 2007. <http://www.who.int/childgrowth/standards/en/>
- Karila C, de Blic J, Waernessyckle S, Benoist MR, Scheinmann P. Cardiopulmonary exercise testing in children: an individualized protocol for workload increase. *Chest.* 2001;120:81–7.
- Graves LEF, Batterham AM, Foweather L, McWhannell N, Hopkins ND, Boddy LM, et al. Scaling of peak oxygen uptake in children: a comparison of three body size index models. *Med Sci Sports Exerc.* 2013;45:2341–5.
- Wasserman K, Hansen JE, Sue DY, Casaburi RWB. Prova de esforço: princípios e interpretação. In: Revinter, editor. Rio de Janeiro: Revisão técnica: Pedro Loureiro. 3<sup>ª</sup> ed. Tradução: Maurício de Arruda Campos; 2005.
- Armstrong N, Kirby BJ, McManus AM, Welsman JR. Prepubescents’ ventilatory responses to exercise with reference to sex and body size. *Chest.* 1997;112:1554–60.
- Welsman JR, Armstrong N. The measurement and interpretation of aerobic fitness in children: current issues. *J R Soc Med.* 1996;89, 281P–5P.
- Cooper CBST. Teste ergométrico: aplicações práticas e interpretação. In: Revinter, editor. Rio de Janeiro: Revisão técnica: Maurício Nunes Rodrigues. 1<sup>ª</sup> ed. Tradução: Daniela Reis Aubaut; 2005. p. 1–114.
- Pereira. Espirometria. *J Pneumol.* 2002;28:S1–82.
- Paridon SM, Alpert BS, Boas SR, Cabrera ME, Calderera LL, Daniels SR, et al. Clinical stress testing in the pediatric age group: a statement from the American Heart Association Council on Cardiovascular Disease in the Young, Committee on Atherosclerosis Hypertension, and Obesity in Youth. *Circulation.* 2006;113:1905–20.
- Glass HC, Costarino AT, Stayer SA, Brett CM, Cladis F, Davis PJ. Outcomes for extremely premature infants. *Anesth Analg.* 2015;120:1337–51.
- Northway WH, Rosan RC, Porter DY. Pulmonary disease following respirator therapy of hyaline-membrane disease. Bronchopulmonary dysplasia. *N Engl J Med.* 1967;276:357–68.
- Riddoch CJ, Mattocks C, Deere K, Saunders J, Kirkby J, Tilling K, et al. Objective measurement of levels and patterns of physical activity. *Arch Dis Child.* 2007;92:963–9.
- Lovering AT, Laurie SS, Elliott JE, Beasley KM, Yang X, Gust CE, et al. Normal pulmonary gas exchange efficiency and absence of exercise-induced arterial hypoxemia in adults with bronchopulmonary dysplasia. *J Appl Physiol.* 2013;115:1050–6.
- Zwiren L. Anaerobic and aerobic capacities of children. *Pediatr Exerc Sci.* 1989;31–44.

27. Rowland TW. *Fisiologia do exercício na criança*. In: Manole, editor. *Fisiologia do exercício na criança*. 2<sup>a</sup> ed. São Paulo: Tradução Maria Salette Tilelli; 2008. p. 68–87.
28. Kathegesu E, Beucher J, Daniel V, Guillot S, Lefeuvre S, Deneuille E, et al. Devenir respiratoire à l'âge scolaire de la dysplasie bronchopulmonaire. *Arch Pédiatr*. 2016;23:325–32.
29. Vollsæter M, Røksund OD, Eide GE, Markestad T, Halvorsen T. Lung function after preterm birth: development from mid-childhood to adulthood. *Thorax*. 2013;68.
30. Robič Pikel T, Starc G, Strel J, Kovač M, Babnik J, Golja P. Impact of prematurity on exercise capacity and agility of children and youth aged 8 to 18. *Early Hum Dev*. 2017;110:39–45.