

## Original Article

## Influence of the *CYP2J2* Gene Polymorphisms on Chronic Obstructive Pulmonary Disease Risk in the Chinese Han Population



Hui Lu<sup>a,1</sup>, Yixiu Yang<sup>b,1</sup>, Xianghong Chen<sup>c</sup>, Cibing Wu<sup>d</sup>, Jie Zhao<sup>d</sup>, Qiong Feng<sup>d</sup>, Xiaoli Zhou<sup>b</sup>, Dongchuan Xu<sup>e</sup>, Quanni Li<sup>b</sup>, Huan Niu<sup>e</sup>, Ping He<sup>e</sup>, Jianfang Liu<sup>d</sup>, Hongxia Yao<sup>b</sup>, Yipeng Ding<sup>b,\*</sup>

<sup>a</sup> Hainan Provincial Key Laboratory for Human Reproductive Medicine and Genetic Research, The First Affiliated Hospital of Hainan Medical University, Hainan Medical University, Haikou 570102, Hainan, China

<sup>b</sup> Department of General Practice, Hainan General Hospital, Haikou 570311, Hainan, China

<sup>c</sup> Department of General Practice, The Second Affiliated Hospital of Hainan Medical University, Haikou 570311, Hainan, China

<sup>d</sup> Hainan General Hospital, University of South China, Haikou 570311, Hainan, China

<sup>e</sup> Department of Emergency, Hainan General Hospital, Haikou 570311, Hainan, China

## ARTICLE INFO

## Article history:

Received 14 October 2019

Accepted 30 November 2019

## Keywords:

*CYP2J2*

Chronic obstructive pulmonary disease

Genetic polymorphism

## ABSTRACT

**Introduction:** Cytochrome P450 (CYP) 2J2 is a major enzyme that controls epoxyeicosatrienoic acids biosynthesis, which may play a role in chronic obstructive pulmonary disease (COPD) development. In this study, we aimed to assess the influence of *CYP2J2* polymorphisms with COPD susceptibility.

**Material and methods:** A case-control study enrolled 313 COPD cases and 508 controls was to investigate the association between *CYP2J2* polymorphisms and COPD risk. Agena MassARRAY platform was used to genotype *CYP2J2* polymorphisms. Odds ratios (OR) and 95% confidence intervals (CI) were calculated to evaluate the association between *CYP2J2* polymorphisms and COPD risk.

**Results:** We observed rs11207535 (homozygote: OR=0.08, 95%CI=0.01–0.96,  $p=0.047$ ; recessive: OR=0.08, 95%CI=0.01–0.94,  $p=0.044$ ), rs10889159 (homozygote: OR=0.08, 95%CI=0.01–0.92,  $p=0.043$ ; recessive: OR=0.08, 95%CI=0.01–0.90,  $p=0.040$ ) and rs1155002 (heterozygote: OR=1.63, 95%CI=1.13–2.36,  $p=0.009$ ; dominant: OR=1.64, 95%CI=1.15–2.35,  $p=0.006$ ; additive: OR=1.45, 95%CI=1.09–1.92,  $p=0.011$ ) were significantly associated with COPD risk. Allelic tests showed T allele of rs2280274 was related to a decreased risk of COPD and T allele of rs1155002 was associated with an increased COPD risk. Stratified analyses indicated the effects of *CYP2J2* polymorphisms and COPD risk were dependent on gender and smoking status ( $p<0.05$ ). Additionally, two haplotypes ( $A_{rs11207535}C_{rs10889159}T_{rs1155002}$  and  $A_{rs11207535}C_{rs10889159}C_{rs1155002}$ ) significantly decreased COPD risk.

**Conclusion:** It suggested *CYP2J2* polymorphisms were associated with COPD susceptibility in the Chinese Han population.

© 2020 Published by Elsevier España, S.L.U. on behalf of SEPAR.

## Influencia de los polimorfismos del gen *CYP2J2* en el riesgo de desarrollo de enfermedad pulmonar obstructiva crónica en la población Han China

## RESUMEN

**Introducción:** El citocromo P450 (CYP) 2J2 es una enzima importante que controla la biosíntesis de los ácidos epoxyeicosatrienoicos, y que podría desempeñar un papel en el desarrollo de la enfermedad pulmonar obstructiva crónica (EPOC). En este estudio, nuestro objetivo fue evaluar la influencia de los polimorfismos de *CYP2J2* en la susceptibilidad a la EPOC.

**Materiales y métodos:** Se realizó un estudio de casos y controles que incluyó 313 casos de EPOC y 508 controles para investigar la asociación entre los polimorfismos de *CYP2J2* y el riesgo de desarrollar EPOC. Se utilizó la plataforma Agena MassARRAY para genotipar los polimorfismos de *CYP2J2*. Se calcularon los *odds ratio* (OR) con unos intervalos de confianza (IC) del 95% para valorar la asociación entre los polimorfismos de *CYP2J2* y el riesgo de la EPOC.

## Palabras clave:

*CYP2J2*

Enfermedad pulmonar obstructiva crónica

Polimorfismo genético

\* Corresponding author.

E-mail address: [DingYP66@163.com](mailto:DingYP66@163.com) (Y. Ding).

<sup>1</sup> These authors contributed equally to this work.

**Resultados:** Observamos que rs11207535 (homocigoto: OR: 0,08, IC 95%: 0,01–0,96; p=0,047; recesivo: OR: 0,08; IC 95%: 0,01–0,94; p=0,044), rs10889159 (homocigoto: OR: 0,08, IC 95%: 0,01–0,92; p=0,043; recesivo: OR: 0,08, IC 95%: 0,01–0,90; p=0,040) y rs1155002 (heterocigoto: OR: 1,63, IC 95%: 1,13–2,36; p=0,009; dominante: OR: 1,64, IC 95%: 1,15–2,35; p=0,006; aditivo: OR: 1,45, IC 95%: 1,09–1,92; p=0,011) se asociaron significativamente con el riesgo de EPOC. Las pruebas alélicas mostraron que el alelo T de rs2280274 estaba relacionado con una disminución del riesgo de EPOC y el alelo T de rs1155002 se asoció con un mayor riesgo de EPOC. Los análisis estratificados indicaron que los efectos de los polimorfismos de *CYP2J2* y el riesgo de EPOC dependían del sexo y del tabaquismo ( $p < 0,05$ ). Además, 2 haplotipos ( $A_{rs11207535}C_{rs10889159}T_{rs1155002}$  y  $A_{rs11207535}C_{rs10889159}C_{rs1155002}$ ) reducían significativamente el riesgo de la EPOC.

**Conclusión:** El estudio sugirió que los polimorfismos de *CYP2J2* se asociaban con la susceptibilidad a la EPOC en la población Han China.

© 2020 Publicado por Elsevier España, S.L.U. en nombre de SEPAR.

## Introduction

Chronic obstructive pulmonary disease (COPD) is the leading cause of morbidity and mortality worldwide, and the prevalence of COPD is rising in China.<sup>1,2</sup> COPD is characterized by persistent respiratory symptoms and reversible airflow limitation.<sup>3</sup> The development of COPD is caused by the interactions of environmental and genetic factors.<sup>4</sup> Cigarette smoking has long been considered a significant risk factor of COPD. Recently, an increasing number of studies proved that genetic characteristics play a vital role in the susceptibility to COPD.<sup>5</sup>

*Cytochrome P450 (CYP) 2J2* gene belongs to the CYP superfamily, which metabolizes arachidonic acid to four regioisomeric epoxyeicosatrienoic acids (5,6-, 8,9-, 11,12-, and 14,15-EET).<sup>6</sup> *CYP2J2* enzyme has influence on metabolism of endogenous and exogenous compounds. *CYP2J2* is the major CYP expressed in heart, placenta, lung, kidney and gastrointestinal tissues.<sup>7</sup> Moreover, *CYP2J2* is responsible for the drug metabolizing.<sup>8</sup> In previous studies, *CYP2J2* had been reported to exert biological effects in the cardiovascular system due to its role in endobiotic metabolism.<sup>9</sup> Notably, *CYP2J2* is found to be overexpressed in various cancers.<sup>10</sup> Furthermore, several *CYP2J2* polymorphisms have been reported to be associated with susceptibility to multiple diseases.<sup>11,12</sup> G-50T is a common single nucleotide polymorphisms (SNP) of *CYP2J2*, which leads to a decrease in the gene expression and results in an altered epoxygenase-dependent arachidonic acid metabolism of eicosanoids that possess important biological functions in the lung and airways.<sup>13</sup> Among the candidate SNPs in our study, rs1155002 was the most popular polymorphism. Studies revealed that rs1155002 was associated with increased risk of late-onset Alzheimer's disease and hypertension.<sup>14,15</sup> However, a limited number of studies have been conducted to investigate the contributions of *CYP2J2* polymorphisms to COPD risk.<sup>16–18</sup>

Thus, it is necessary to estimate the relationships of COPD risk and *CYP2J2* polymorphisms. The present study was designed to explore whether *CYP2J2* polymorphisms (rs2280274, rs4388726, rs11207535, rs10889159, and rs1155002) had relationship with the susceptibility of COPD based on a case-control study.

## Materials and methods

### Study participants

A total of 313 COPD patients (238 men and 75 women) and 508 healthy controls (337 men and 171 women) were consecutively recruited from Hainan General Hospital. All COPD patients had airway obstruction on the spirometry and clinical features of COPD (cough, sputum production, dyspnea), and they were confirmed as post-bronchodilator FEV<sub>1</sub>/FVC < 70% according to the criteria

established by the National Heart, Lung and Blood Institute/World Health Organization Global Initiative for Chronic Obstructive Lung Disease.<sup>19</sup> Patients with other respiratory diseases, autoimmune diseases or cancers were excluded from the study. Healthy controls with FEV<sub>1</sub>/FVC > 70% were randomly recruited from the same hospital with patients during the same period, which included the subjects without diseases or family history of any diseases. Informed consents were obtained from all participants. And, our study protocol was strictly conformed to the Declaration of Helsinki and was carried out with the approval from the ethics committee of Hainan General Hospital.

### SNP selection and genotyping

Combined previous studies, five SNPs of *CYP2J2* had minor allele frequencies (MAFs) > 5% in the HapMap Chinese Han Beijing population were selected in this study. Genomic DNA was isolated from whole-blood samples using the blood DNA kit (GoldMag Co. Ltd., Xi'an, China), and DNA concentrations were determined by Nanodrop 2000 (Thermo Scientific, Waltham, MA, USA).<sup>20</sup> Agena MassARRAY Assay Design 3.0 software was performed to design primers used in the study (Table 1). Genotyping was conducted by Agena MassARRAY system (Agena, San Diego, CA, USA), according to the manufacturer's protocol. In addition, we performed the data management and analysis using the Agena Typer 4.0 Software (San Diego, CA, USA).<sup>21</sup>

### Statistical analysis

We used SPSS 22.0 statistical package (SPSS, Chicago, IL, USA) to perform statistical analyses. All continuous data were presented as mean ± standard deviation. The chi square test and Student's *t*-test were used to compare the differences of categorical and continuous variables. Hardy-Weinberg equilibrium (HWE) for each SNP was assessed using Fisher's exact test in control group. For evaluating the impact of candidate SNPs on COPD risk, we used logistic regression analyses with adjustment for age and gender to estimate the odds ratio (OR) and 95% confidence interval (CI). Then, genetic models (co-dominant, dominant, recessive and additive) were used to evaluate the associations between *CYP2J2* polymorphisms and COPD risk by PLINK software (version 1.07). Furthermore, we used PLINK software (version 1.07) and Haploview software (version 4.2) for haplotype analysis and linkage disequilibrium (LD).<sup>22</sup> All *p* values shown in our study were two sided, and  $p < 0,05$  was regarded as statistically significant.

**Table 1**  
Primers used for this study.

SNP	First PCR primer	Second PCR primer	Uep.Dir	Use Seq
rs2280274	ACGTTGGATGCAAACCACTTAAAGCTCAC	ACGTTGGATGATCTAGGAGATGAAAAGAGG	R	GGAGATGAAAAGAGGATAATG
rs4388726	GTTGGATGCTCTTTTCATCTCTAGA	ACGTTGGATGAACCTAGATCCAAATCCCAGC	R	AAATCCCAGCTCTACTGTCT
rs11207535	ACGTTGGATGGCTGAATTGAATCTCTGCC	ACGTTGGATGACAGCTGACACTAGCATTG	F	cgGTTATGTGGGTGAACCTG
rs10889159	ACGTTGGATGCTGTGATCTGGTTAATCTC	ACGTTGGATGTAATGCTACACCCAAAAG	R	gatCCACCCAAAAGATTTTCTTTA
rs1155002	ACGTTGGATGAGGCAGACTCTATTACAGC	ACGTTGGATGTCATCCATCCAAAGGTGTC	R	attttCAGCACTTGGGGCAGGACA

Abbreviations: SNP, single nucleotide polymorphism; USE SEQ, unextended mini-sequencing primer.

**Table 2**  
The characteristic of case and control.

Variable	Case (N = 313)	Control (N = 508)	p
Age (years, SD)	71.80 ± 10.09	60.05 ± 6.48	<b>&lt;0.001</b>
Gender			<b>0.004</b>
Men	238 (76.04%)	337 (66.34%)	
Women	75 (23.96%)	171 (33.66%)	
FEV <sub>1</sub> (L)	1.37 ± 0.32	2.12 ± 0.89	<b>&lt;0.001</b>
FEV <sub>1</sub> (%)	62.0 ± 15.3	84.6 ± 12.6	<b>&lt;0.001</b>
FVC (L)	2.45 ± 0.64	2.68 ± 0.73	<b>&lt;0.001</b>
FVC (%)	53.0 ± 16.7	83.0 ± 10.2	<b>&lt;0.001</b>
FEV <sub>1</sub> /FVC	0.56 ± 0.05	0.79 ± 0.04	<b>&lt;0.001</b>
BMI, kg/m <sup>2</sup>	24.67 ± 4.62	24.35 ± 4.58	0.587
Smoking status			0.082
Yes	149 (47.60%)	216 (42.5%)	
No	164 (52.40%)	292 (42.5%)	
Pack-years smoked	32.96 ± 20.53	35.12 ± 22.08	
Comorbidity			
Yes	92 (29.39%)		
No	221 (70.61%)		

Abbreviations: SD, standard deviation; FEV<sub>1</sub>, forced expiratory volume in 1 second; FVC, forced volume capacity; BMI, body mass index.

No smoking means never smoked.

p < 0.05 indicates statistical significance.

Bold data mean significant difference.

## Results

### Characteristics of the participants

The basic characteristics of 821 participants in the study are provided in Table 2. The mean ages of cases and controls were 71.80 ± 10.09 and 60.05 ± 6.48 years old. The majority of patients and controls were men (76.04% and 66.34%, respectively). The p values for age and gender were less than 0.05. Compared with the control group, the cases had significantly lower FEV<sub>1</sub> (L), FEV<sub>1</sub> (%), FVC (L), FVC (%) and FEV<sub>1</sub>/FVC (p < 0.001). There was no significant distribution of BMI in two groups (p = 0.587). There are 149 (47.60%) and 216 (42.5%) smokers in cases and controls, individually. The pack-years smoked were 32.96 ± 20.53 and 35.12 ± 22.08. Among the patients with COPD, 92 (29.39%) had comorbidity.

### Association between CYP2J2 SNPs and COPD susceptibility

The basic information of the selected SNPs is shown in Table 3. All SNPs were in accord with HWE in the controls (p > 0.05). The minor allele frequencies (MAFs) of the analyzed SNPs in the cases and controls were also presented. Then, we compared the differences in frequency distributions of alleles between two groups, and we found rs2280274 and rs1155002 were significantly associated with COPD risk. Among them, the minor allele “T” of rs2280274 decreased the risk of COPD (OR = 0.69, 95%CI = 0.48–0.99, p = 0.041), whereas the individuals carried “T” of rs1155002 were related to higher risk of COPD (OR = 1.50, 95%CI = 1.21–1.87, p < 0.001). After adjustment for age and gender, the genotype frequencies of CYP2J2 polymorphisms and their association with COPD risk in the

genetic models are shown in Table 4. We found that GG genotype of rs11207535 was associated with a decreased risk of COPD in the homozygote model (OR = 0.08, 95%CI = 0.01–0.96, p = 0.047) and recessive model (OR = 0.08, 95%CI = 0.01–0.94, p = 0.044). Meanwhile, rs10889159 had a strong relationship with lower COPD risk in homozygote model (OR = 0.08, 95%CI = 0.01–0.92, p = 0.043) and recessive model (OR = 0.08, 95%CI = 0.01–0.90, p = 0.040). In addition, multiple model analyses revealed that rs1155002 was positively associated with higher susceptibility of COPD (heterozygote: OR = 1.63, 95%CI = 1.13–2.36, p = 0.009; dominant: OR = 1.64, 95%CI = 1.15–2.35, p = 0.006; additive: OR = 1.45, 95%CI = 1.09–1.92, p = 0.011).

### Stratification analysis

The association between CYP2J2 polymorphisms and COPD risk was further assessed in subgroup of gender (Table 5). For men, rs11207535 and rs10889159 of CYP2J2 were associated with a decreased COPD risk in homozygote (rs11207535: OR = 0.07, 95%CI = 0.01–0.99, p = 0.049; rs10889159: OR = 0.07, 95%CI = 0.01–0.96, p = 0.046) and recessive (rs11207535: OR = 0.07, 95%CI = 0.01–0.92, p = 0.043) models, when compared to the respective reference groups. And, rs1155002 was associated with higher COPD risk in dominant (OR = 1.57, 95%CI = 1.01–2.46, p = 0.046) and allele (OR = 1.45, 95%CI = 1.13–1.86, p = 0.004) models. The other SNPs did not have effects on COPD susceptibility for all participants (p > 0.05).

Then, we did the association of CYP2J2 SNPs and COPD risk stratified by smoking status (Table 6). For smoker, rs11207535 and rs1155002 had strong relationship with increased risk of COPD (rs11207535, heterozygote: OR = 2.81, 95%CI = 1.17–6.75, p = 0.021, dominant: OR = 2.56, 95%CI = 1.09–6.02, p = 0.031; rs1155002: OR = 1.40, 95%CI = 1.01–1.93, p = 0.043). Rs2280274 and rs1155002 were associated with COPD risk among non-smoker, rs2280274 significantly decreased COPD risk in dominant (OR = 0.48, 95%CI = 0.25–0.89, p = 0.020), additive (OR = 0.46, 95%CI = 0.25–0.83, p = 0.010) and allele (OR = 0.45, 95%CI = 0.27–0.76, p = 0.002) models, rs1155002 was significantly associated with higher risk of COPD (heterozygote: OR = 2.12, 95%CI = 1.31–3.43, p = 0.002; dominant: OR = 2.11, 95%CI = 1.32–3.37, p = 0.002; additive: OR = 1.71, 95%CI = 1.18–2.49, p = 0.005; allele: OR = 1.63, 95%CI = 1.22–2.18, p = 0.001).

### Haplotype analysis

Finally, we did the LD block and found two strong blocks (Fig. 1). The association between haplotypes of CYP2J2 SNPs and COPD risk is listed in Table 7. The “A-C-T” and “A-C-C” haplotypes of rs11207535, rs10889159 and rs1155002 were significantly decreased the COPD risk (OR = 0.69, 95%CI = 0.52–0.92, p = 0.011; OR = 0.76, 95%CI = 0.58–0.99, p = 0.044; respectively).

**Table 3**  
Basic information of candidate SNPs.

SNP	Chromosome	Position	Alleles	MAF		HWE- <i>p</i>	OR (95%CI)	<i>p</i>
				Case	Control			
rs2280274	1	59893405	A/T	0.077	0.107	0.820	0.69 (0.48–0.99)	<b>0.041</b>
rs4388726	1	59893484	A/G	0.054	0.065	0.713	0.83 (0.54–1.27)	0.381
rs11207535	1	59894523	C/T	0.101	0.108	1.000	0.92 (0.66–1.28)	0.625
rs10889159	1	59894850	C/T	0.113	0.134	0.848	0.82 (0.60–1.12)	0.211
rs1155002	1	59908103	C/T	0.348	0.262	0.422	1.50 (1.21–1.87)	<b>&lt;0.001</b>

Abbreviations: SNP, single nucleotide polymorphism; MAF, minor allele frequency; HWE, Hardy–Weinberg equilibrium; OR, odds ratio; CI, confidence interval.  
*p* < 0.05 indicates statistical significance.  
 Bold data mean significant difference.

**Table 4**  
Genotype frequencies of the SNPs and their associations with risk of COPD.

SNP	Model	Genotype	Genotype frequencies		OR (95%CI)	<i>p</i>	
			Case	Control			
rs2280274	Homozygote	TT	1 (0.3%)	6 (1.2%)	0.08 (0.01–1.01)	0.051	
		Heterozygote	TA	46 (14.7%)	97 (19.1%)	0.89 (0.55–1.44)	0.625
		AA	266 (85.0%)	405 (79.7%)	1.00		
	Dominant	TT-TA	47 (15.0%)	103 (20.3%)	0.81 (0.50–1.30)	0.380	
		AA	266 (85.0%)	405 (79.7%)	1.00		
	Recessive	TT	1 (0.3%)	6 (1.2%)	0.09 (0.01–1.03)	0.052	
TA-AA		312 (99.7%)	502 (98.8%)	1.00			
rs4388726	Additive	AA	0 (-)	1 (0.2%)	–	–	
		Homozygote	AG	34 (10.9%)	64 (12.6%)	0.89 (0.50–1.58)	0.691
		Heterozygote	GG	279 (89.1%)	443 (87.2%)	1.00	
	Dominant	AA-AG	34 (10.9%)	65 (12.8%)	0.87 (0.49–1.53)	0.623	
		GG	279 (89.1%)	443 (87.2%)	1.00		
	Recessive	AA	0 (-)	1 (0.2%)	–	–	
AG-GG		313 (100%)	507 (99.8%)	1.00			
rs11207535	Additive	GG	1 (0.3%)	6 (1.2%)	0.85 (0.49–1.48)	0.559	
		Homozygote	GA	61 (19.5%)	98 (19.3%)	0.08 (0.01–0.96)	<b>0.047</b>
		Heterozygote	AA	251 (80.2%)	404 (79.5%)	1.15 (0.73–1.81)	0.542
	Dominant	GG-GA	62 (19.8%)	104 (20.5%)	1.04 (0.67–1.63)	0.853	
		AA	251 (80.2%)	404 (79.5%)	1.00		
	Recessive	GG	1 (0.3%)	6 (1.2%)	0.08 (0.01–0.94)	<b>0.044</b>	
GA-AA		312 (99.7%)	502 (98.8%)	1.00			
rs10889159	Additive	TT	1 (0.3%)	8 (1.6%)	0.94 (0.62–1.43)	0.766	
		Homozygote	TC	68 (21.9%)	119 (23.6%)	0.08 (0.01–0.92)	<b>0.043</b>
		Heterozygote	CC	242 (77.8%)	378 (74.8%)	1.13 (0.74–1.73)	0.570
	Dominant	TT-TC	69 (22.2%)	127 (25.2%)	1.03 (0.68–1.57)	0.881	
		CC	242 (77.8%)	378 (74.8%)	1.00		
	Recessive	TT	1 (0.3%)	8 (1.6%)	0.08 (0.01–0.90)	<b>0.040</b>	
TC-CC		310 (99.7%)	497 (98.4%)	1.00			
rs1155002	Additive	TT	28 (8.9%)	31 (6.1%)	0.93 (0.63–1.38)	0.732	
		Homozygote	TC	162 (51.8%)	204 (40.2%)	1.71 (0.86–3.42)	0.128
		Heterozygote	CC	123 (39.3%)	272 (53.7%)	1.63 (1.13–2.36)	<b>0.009</b>
	Dominant	TT-TC	190 (60.7%)	235 (46.3%)	1.00		
		CC	123 (39.3%)	272 (53.7%)	1.64 (1.15–2.35)	<b>0.006</b>	
	Recessive	TT	28 (8.9%)	31 (6.1%)	1.34 (0.69–2.61)	0.389	
TC-CC		285 (91.1%)	476 (93.9%)	1.00			
Additive	TT	28 (8.9%)	31 (6.1%)	1.45 (1.09–1.92)	<b>0.011</b>		
	TC-CC	285 (91.1%)	476 (93.9%)	1.00			

Abbreviations: SNP, single nucleotide polymorphism; COPD, chronic obstructive pulmonary disease; OR, odds ratio; CI, confidence interval.  
*p* < 0.05 indicates statistical significance.  
 Bold data mean significant difference.

## Discussion

In this study, we investigated the associations between five SNPs in *CYP2J2* and COPD risk among the Chinese Han population. Our results indicated that rs2280274, rs11207535, rs10889159 and rs1155002 were closely associated with the risk of COPD. The subgroup analysis revealed that men with GG genotype of rs11207535 and TT genotype of rs10889159 had a lower COPD risk, whereas individuals with T allele of rs1155002 significantly increased the risk of COPD. In addition, haplotype analysis showed that rs11207535, rs10889159 and rs1155002 worked together, the

haplotype ACT and ACC could be the protective factors for COPD. It gives us a clue that *CYP2J2* polymorphisms could be useful targets for personalized therapy on COPD.

*CYP2J2* is one of *CYP* enzyme, which is expressed in human pulmonary endothelium.<sup>23</sup> *CYP2J2* plays a vital physiological part in the pulmonary regulation of both vascular and bronchial tone.<sup>18</sup> Recent studies indicated that *CYP2J2* mediated EETs could reduce excessive inflammation in the liver and lung by using mouse models.<sup>24,25</sup> Satoshi et al. observed that 50% reduction in *CYP2J2* mRNA expression in alveolar epithelial type II cells isolated from COPD patients compared to smokers without COPD.<sup>26</sup>

**Table 5**  
The association between *CYP2J2* SNPs and COPD risk stratified by sex.

SNP	Model	Men		Women	
		OR (95%CI)	p	OR (95%CI)	p
rs2280274	Homozygote	0.07 (0.01–1.03)	0.052	–	–
	Heterozygote	1.04 (0.56–1.94)	0.893	0.65 (0.29–1.44)	0.287
	Dominant	0.92 (0.50–1.70)	0.798	0.61 (0.28–1.34)	0.219
	Recessive	0.07 (0.01–1.02)	0.052	–	–
	Additive	0.82 (0.46–1.46)	0.506	0.59 (0.28–1.27)	0.177
	Allele	0.79 (0.52–1.20)	0.272	0.54 (0.27–1.07)	0.073
rs4388726	Homozygote	–	–	–	–
	Heterozygote	–	–	0.67 (0.25–1.78)	0.421
	Dominant	1.02 (0.50–2.11)	0.950	0.63 (0.24–1.66)	0.349
	Recessive	–	–	–	–
	Additive	1.02 (0.50–2.11)	0.950	0.61 (0.24–1.56)	0.303
	Allele	0.93 (0.56–1.53)	0.771	0.62 (0.26–1.47)	0.274
rs11207535	Homozygote	0.07 (0.01–0.99)	<b>0.049</b>	–	–
	Heterozygote	1.24 (0.69–2.24)	0.468	1.03 (0.50–2.09)	0.945
	Dominant	1.10 (0.62–1.96)	0.752	0.95 (0.47–1.92)	0.893
	Recessive	0.07 (0.01–0.96)	<b>0.046</b>	–	–
	Additive	0.96 (0.56–1.65)	0.882	0.88 (0.46–1.71)	0.715
	Allele	0.99 (0.66–1.47)	0.942	0.87 (0.48–1.57)	0.635
rs10889159	Homozygote	0.07 (0.01–0.96)	<b>0.046</b>	–	–
	Heterozygote	1.24 (0.72–2.14)	0.438	0.97 (0.49–1.93)	0.935
	Dominant	1.11 (0.65–1.89)	0.704	0.90 (0.46–1.79)	0.772
	Recessive	0.07 (0.01–0.92)	<b>0.043</b>	–	–
	Additive	0.98 (0.59–1.62)	0.929	0.84 (0.45–1.59)	0.598
	Allele	0.87 (0.60–1.26)	0.464	0.77 (0.43–1.36)	0.364
rs1155002	Homozygote	1.72 (0.76–3.94)	0.196	1.80 (0.47–6.90)	0.393
	Heterozygote	1.55 (0.97–2.46)	0.065	1.76 (0.95–3.26)	0.073
	Dominant	1.57 (1.01–2.46)	<b>0.046</b>	1.76 (0.97–3.21)	0.064
	Recessive	1.38 (0.63–3.05)	0.421	1.39 (0.37–5.17)	0.622
	Additive	1.41 (0.99–1.99)	0.055	1.56 (0.95–2.56)	0.082
	Allele	1.45 (1.13–1.86)	<b>0.004</b>	1.50 (0.97–2.31)	0.065

Abbreviations: SNP, single nucleotide polymorphism; COPD, chronic obstructive pulmonary disease; OR, odds ratio; CI, confidence interval.

$p < 0.05$  indicates statistical significance.

Bold data mean significant difference.


**Table 6**  
The association between *CYP2J2* SNPs and COPD risk stratified by smoking status.

SNP	Model	Smoker		Non-smoker	
		OR (95%CI)	p	OR (95%CI)	p
rs2280274	Homozygote	–	–	–	–
	Heterozygote	2.38 (0.95–5.96)	0.063	0.54 (0.28–1.01)	0.054
	Dominant	2.38 (0.95–5.93)	0.063	0.48 (0.25–0.89)	<b>0.020</b>
	Recessive	–	–	–	–
	Additive	2.33 (0.95–5.72)	0.065	0.46 (0.25–0.83)	<b>0.010</b>
	Allele	1.11 (0.66–1.88)	0.690	0.45 (0.27–0.76)	<b>0.002</b>
rs4388726	Homozygote	–	–	–	–
	Heterozygote	–	–	0.63 (0.29–1.34)	0.229
	Dominant	1.88 (0.66–5.33)	0.238	0.59 (0.28–1.27)	0.180
	Recessive	–	–	–	–
	Additive	1.88 (0.66–5.33)	0.238	0.58 (0.28–1.21)	0.148
	Allele	1.24 (0.67–2.29)	0.489	0.58 (0.31–1.07)	0.078
rs11207535	Homozygote	0.32 (0.01–19.63)	0.589	–	–
	Heterozygote	2.81 (1.17–6.75)	<b>0.021</b>	0.84 (0.48–1.48)	0.548
	Dominant	2.56 (1.09–6.02)	<b>0.031</b>	0.76 (0.43–1.33)	0.338
	Recessive	0.28 (0.01–15.95)	0.540	–	–
	Additive	2.14 (0.97–4.73)	0.061	0.70 (0.41–1.19)	0.184
	Allele	1.08 (0.66–1.78)	0.751	0.81 (0.52–1.26)	0.342
rs10889159	Homozygote	0.33 (0.01–19.74)	0.592	–	–
	Heterozygote	2.16 (0.99–4.71)	0.053	0.86 (0.50–1.49)	0.597
	Dominant	2.03 (0.94–4.38)	0.071	0.78 (0.46–1.34)	0.369
	Recessive	0.29 (0.01–16.12)	0.543	–	–
	Additive	1.8 (0.87–3.70)	0.113	0.72 (0.43–1.19)	0.199
	Allele	0.94 (0.60–1.49)	0.800	0.73 (0.48–1.11)	0.138
rs1155002	Homozygote	2.21 (0.68–7.18)	0.188	2.06 (0.84–5.08)	0.116
	Heterozygote	1.36 (0.71–2.59)	0.350	2.12 (1.31–3.43)	<b>0.002</b>
	Dominant	1.47 (0.79–2.72)	0.224	2.11 (1.32–3.37)	<b>0.002</b>
	Recessive	1.92 (0.61–5.99)	0.263	1.37 (0.58–3.24)	0.472
	Additive	1.43 (0.88–2.33)	0.152	1.71 (1.18–2.49)	<b>0.005</b>
	Allele	1.40 (1.01–1.93)	<b>0.043</b>	1.63 (1.22–2.18)	<b>0.001</b>

Abbreviations: SNP, single nucleotide polymorphism; COPD, chronic obstructive pulmonary disease; OR, odds ratio; CI, confidence interval.

$p < 0.05$  indicates statistical significance.

Bold data mean significant difference.



**Fig. 1.** Haplotype block map for the *CYP2J2* polymorphisms.

Block 1 includes rs2280274 and rs4388726. Block 2 includes rs11207535, rs10889159 and rs1155002. The LD between two SNPs is standardized  $D'$ .

**Table 7**

The haplotype of *CYP2J2* SNPs and COPD risk.

Block	SNP	Haplotype	Haplotype frequency		OR (95%CI)	<i>p</i>
			Case	Control		
1	rs2280274 rs4388726	TA	0.946	0.935	1.18 (0.68–2.06)	0.559
1	rs2280274 rs4388726	TG	0.978	0.958	1.64 (0.77–3.51)	0.201
1	rs2280274 rs4388726	AG	0.923	0.893	1.33 (0.85–2.09)	0.208
2	rs11207535 rs10889159 rs1155002	ACT	0.652	0.738	0.69 (0.52–0.92)	<b>0.011</b>
2	rs11207535 rs10889159 rs1155002	GTC	0.899	0.893	1.06 (0.69–1.60)	0.801
2	rs11207535 rs10889159 rs1155002	ATC	0.987	0.971	1.31 (0.53–3.27)	0.561
2	rs11207535 rs10889159 rs1155002	ACC	0.538	0.601	0.76 (0.58–0.99)	<b>0.044</b>

**Abbreviations:** SNP, single nucleotide polymorphism; COPD, chronic obstructive pulmonary disease; OR, odds ratio; CI, confidence interval.

$p < 0.05$  indicates statistical significance.

Bold data mean significant difference.

Nevertheless, polymorphisms in *CYP2J2* gene could affect the levels of transcription, translation, enzyme activity and result in various substances, which may affect the development of COPD. For instance, *CYP2J2* rs890293 had been analyzed on the contribution to respiratory diseases among Russian and they did not find the association between *CYP2J2* rs890293 and respiratory diseases.<sup>27</sup> In this study, we found rs2280274, rs11207535 and rs10889159 decreased COPD risk, rs1155002 reversely increased the risk of COPD among the Chinese Han population, the differences with previous findings may be attributed to study population, sample size and research method. We also found that rs11207535, rs10889159 and rs1155002 were associated with COPD susceptibility in the subgroup of men. It demonstrated that *CYP2J2* polymorphisms had a relationship with COPD risk and the effects of polymorphisms on COPD is gender dependent, suggesting the occurrence of COPD was related to hormone levels.

In addition, smoking has a significant association with risk of lung diseases. Therefore, we estimated the relationship of *CYP2J2* polymorphisms with COPD risk. We observed a protective effect of rs2280274 on COPD risk among non-smokers. It may attribute to the lower levels inflammation from the lack of smoking. Rs11207535 increased the COPD risk of smokers, and rs1155002 had a higher risk of COPD in two groups. Our results suggest that smoking status has significant influence on COPD risk. However, our results are required to be verified by further work.

Some limitations of this study should not be ignored. First, the limitation of our sample size may influence our results, particularly

in the stratified analysis. Second, several potential environmental factors were not included in this study, which may have impact on COPD risk. Third, we did not study the effects of some other diseases on COPD due to the limited information. Finally, the polymorphisms of *CYP2J2* studied in this study were limited, more information on the associations between *CYP2J2* gene and COPD risk needed to be studied in the future.

## Conclusion

Our findings suggest that polymorphisms of *CYP2J2* gene are associated with COPD susceptibility and their effects on COPD risk are sex- and smoking status- specific. Further studies are required to provide accurate evidences about the influence of *CYP2J2* polymorphisms on COPD risk.

## Clinical implications

The prevalence of COPD is increasing in China. This case-control study focused on the associations of *CYP2J2* polymorphisms (rs2280274, rs4388726, rs11207535, rs10889159 and rs1155002) and COPD risk. We found that rs2280274, rs11207535 and rs10889159 significantly decreased COPD risk, indicating they could protect individuals from COPD. *CYP2J2* rs1155002 was associated with higher risk of COPD, suggesting it may serve as a biomarker for detecting or treating COPD. In addition, stratified

analysis showed that the effects of *CYP2J2* polymorphisms on COPD risk were related to gender and smoking status, it is helpful for prevention, diagnosis and individual treatment of COPD. In conclusion, our results provided information on exploring the mechanism and targeted therapy of COPD, it also promotes the development of precision medicine on COPD.

### Funding

This article was financially supported by the National Natural Science Foundation of China (Nos. 81660013 and 81860015), Key Research and Development Plan of Hainan Province (No. ZDYF2018116) and Hainan Provincial Natural Science Foundation of China (No. 818QN315).

### Conflict of interests

The authors declare no conflict of interest.

### Acknowledgement

We sincerely thank all participants in this study.

### References

- Ding Y, Xu H, Yao J, Xu D, He P, Yi S, et al. Association between *RTEL1* gene polymorphisms and COPD susceptibility in a Chinese Han population. *Int J Chronic Obstr Pulm Dis*. 2017;12:931–6.
- Fincham JE. An unfortunate and avoidable component of American pharmacy: tobacco. *Am J Pharm Educ*. 2008;72:57.
- Vogelmeier CF, Criner GJ, Martinez FJ, Anzueto A, Barnes PJ, Bourbeau J, et al. Global strategy for the diagnosis, management, and prevention of chronic obstructive lung disease 2017 report. GOLD executive summary. *Am J Respir Crit Care Med*. 2017;195:557.
- Silverman EK, Vestbo J, Agustí A, Anderson W, Bakke PS, Barnes KC, et al. Opportunities and challenges in the genetics of COPD 2010: an international COPD genetics conference report. *COPD*. 2011;8:121–35.
- Zhang Y, Qiu J, Zhang P, Zhang J, Jiang M, Ma Z. Genetic variants in *FAM13A* and *IREB2* are associated with the susceptibility to COPD in a Chinese rural population: a case–control study. *Int J Chronic Obstr Pulm Dis*. 2018;13:1735–45.
- Yang Y, Dong R, Chen Z, Hu D, Fu M, Tang Y, et al. Endothelium-specific *CYP2J2* overexpression attenuates age-related insulin resistance. *Aging Cell*. 2018;17:e12718.
- Bièche I, Narjoz C, Asselah T, Vacher S, Marcellin P, Lidereau R, et al. Reverse transcriptase-PCR quantification of mRNA levels from cytochrome (*CYP*)1, *CYP2* and *CYP3* families in 22 different human tissues. *Pharmacogenet Genomics*. 2007;17:731.
- Lee E, Kim JH, Shon JC, Wu Z, Kim HJ, Gim M, et al. Terfenadone is a strong inhibitor of *CYP2J2* present in the human liver and intestinal microsomes. *Drug Metab Pharmacokinet*. 2018;33:159–63.
- Aliwarga T, Evangelista EA, Sotoodehnia N, Lemaitre RN, Totah RA. Regulation of *CYP2J2* and *EET* levels in cardiac disease and diabetes. *Int J Mol Sci*. 2018;19:1916.
- Karkhanis A, Hong Y, Chan ECY. Inhibition and inactivation of human *CYP2J2*: implications in cardiac pathophysiology and opportunities in cancer therapy. *Biochem Pharmacol*. 2017;135:12–21.
- Geng S, Wang Y, Sun Y, Li J, Yin H, Zeng Z, et al. Gene–gene interaction between *CYP2J2* and *PPAR-γ* gene on late-onset Alzheimer's disease in the eastern Chinese Han population. *Behav Brain Res*. 2017;322:362–7.
- Zhu Q, Fu Z, Ma Y, Yang H, Huang D, Xie X, et al. A novel polymorphism of the *CYP2J2* gene is associated with coronary artery disease in Uyghur population in China. *Clin Biochem*. 2013;46:1047–54.
- Polonikov AV, Ivanov VP, Solodilova MA, Khoroshaya IV, Kozhuhov MA, Panfilov VI. Promoter polymorphism G-50T of a human *CYP2J2* epoxygenase gene is associated with common susceptibility to asthma. *Chest*. 2007;132:120–6.
- Geng S, Wang Y, Sun Y, Li J, Yin H, Zeng Z, et al. Gene–gene interaction between *CYP2J2* and *PPAR-γ* gene on late-onset Alzheimer's disease in the eastern Chinese Han population. *Behav Brain Res*. 2017;322 Pt B:362–7.
- Sheng-Nan W, Yi Z, Gardner CO, Qi C, Yan L, Gu-Liang W, et al. Evidence for association of polymorphisms in *CYP2J2* and susceptibility to essential hypertension. *Ann Hum Genet*. 2012;71:519–25.
- Akhmadishina LZ, Korytina GF, Victorova TV. *CYP1B1* (4326C>G), *CYP2F1* (c.14.15insC), *CYP2J2* (-76G>T), and *CYP2S1* (13106C>T and 13255A>G) polymorphisms and genetic predisposition to chronic respiratory diseases induced by smoking and occupational factors. *Russ J Genet*. 2011;47:1248–55.
- Satoshi K, Naoya F, Mitsuhiro Y, Ken G, Satoshi S, Chiharu O, et al. Expression of cytochrome P450 mRNAs in Type II alveolar cells from subjects with chronic obstructive pulmonary disease. *Pharmacol Res Perspect*. 2018;6:e00405.
- Hukkanen J, Pelkonen O, Hakkola J, Raunio H. Expression and regulation of xenobiotic-metabolizing cytochrome P450 (*CYP*) enzymes in human lung. *Crit Rev Toxicol*. 2002;32:391–411.
- Vestbo J, Hurd SS, Rodriguez-Roisin R. [An overview of Global Strategy for the Diagnosis, Management and Prevention of Chronic Obstructive Pulmonary Disease (GOLD) (revised 2011)]. *Zhonghua Yi Xue Za Zhi*. 2012;92:937.
- Lei G, Asmitananda T, Yiqian L, Shuo Z, Ting W, Tianjun C, et al. Polymorphisms in the *TERT* gene are associated with lung cancer risk in the Chinese Han population. *Eur J Cancer Prev*. 2014;23:497–501.
- Ding Y, Yang D, Zhou L, Xu J, Chen Y, He P, et al. Variants in multiple genes polymorphism association analysis of COPD in the Chinese Li population. *Int J Chronic Obstr Pulm Dis*. 2015;10:1455–63.
- Zhang S, Thakur A, Liang Y, Wang T, Gao L, Yang T, et al. Polymorphisms in C-reactive protein and Glypican-5 are associated with lung cancer risk and Gartrokin-1 influences Cisplatin-based chemotherapy response in a Chinese Han population. *Dis Markers*. 2015;2015:824304.
- Zeldin DC, Foley J, Boyle JE, Moomaw CR, Tomer KB, Parker C, et al. Predominant expression of an arachidonate epoxygenase in islets of Langerhans cells in human and rat pancreas. *Endocrinology*. 1997;138:1338.
- Chen W, Yang S, Ping W, Fu X, Xu Q, Wang J. *CYP2J2* and *EETs* protect against lung ischemia/reperfusion injury via anti-inflammatory effects in vivo and in vitro. *Cell Physiol Biochem*. 2015;35:2043–54.
- Li R, Xu X, Chen C, Wang Y, Gruzdev A, Zeldin DC, et al. *CYP2J2* attenuates metabolic dysfunction in diabetic mice by reducing hepatic inflammation via the *PPARγ*. *Am J Physiol Endocrinol Metab*. 2015;308:270–82.
- Kamata S, Fujino N, Yamada M, Grime K, Suzuki S, Ota C, et al. Expression of cytochrome P450 mRNAs in Type II alveolar cells from subjects with chronic obstructive pulmonary disease. *Pharmacol Res Perspect*. 2018;6:e00405.
- Akhmadishina LA, Korytina GF, Victorova TV. [Polymorphic markers of the *CYP1B1* (4326C>G), *CYP2F1* (c.14.15insC), *CYP2J2* (-76G>T), and *CYP2S1* (13106C>T and 13255A>G) genes and genetic predisposition to chronic respiratory diseases induced by smoking and occupational factors]. *Russ J Genet*. 2011;47:1248–55.