

7. Kapralik J, Wayne C, Chan E, Nasr A. Surgical versus conservative management of congenital pulmonary airway malformation in children: A systematic review and meta-analysis. *J Pediatr Surg.* 2016;51:508–12.
8. David M, Lamas-Pinheiro R, Henriques-Coelho T. Prenatal and Postnatal Management of congenital Pulmonary Airway Malformation. *Neonatology.* 2016;110:101–15.
9. Bolde S, Pudale S, Pandit G, Ruikar K, Ingle SB. Congenital pulmonary airway malformation: A report of two cases. *World J Clin Cases.* 2015;3:470–3.
10. Taqvi I, Grikasitis M, Eastham K. Not all infantile respiratory distress in writer is bronchiolitis: Congenital lobar emphysema. *BMJ Case Rep.* 2011. <http://dx.doi.org/10.1136/bcr.08.2011.4618>. 2011;1-bcr0820114618.
11. Mattioli G, Pini A, Razole B, Leonelli L, Pio L, Avanzini S, et al. Congenital Lung Malformations: Shifting from open to Thoracoscopic Surgery. *Pediatr Neonatol.* 2016;57:463–6.
12. Adams S, Jobson M, Sangnawakij P, Heetun A, Thaventhiran A, Johal N, et al. Dos thoracoscopy have advantages over open surgery for asymptomatic congenital lung malformations? An analysis of 1626 resections. *J Pediatr Surg.* 2017;52:247–51.



The Effect of a Basic Basket on Tuberculosis Treatment Outcome in the Huambo Province, Angola

Efecto de una cesta básica alimentaria en el resultado del tratamiento de la tuberculosis en la provincia de Huambo, Angola

Dear Director:

Tuberculosis (TB) continues to be a worldwide health problem and is the greatest infectious cause of death among young people and adults.¹

Although TB treatment is free, patients will often stop treatment due to the inability of the patient and his/her relatives to provide the resources necessary to survive.² Studies have shown that poverty and TB are closely linked, since the conditions under which poor populations live make them prone to the onset of the disease.^{1,3} The economic impact of TB arises from the fact that the majority of those affected are part of the labour force.¹ Therefore, incentives have been employed by national TB control programs as a means of encouraging patients to adhere to treatment, helping to minimize the costs associated with the disease, as well as providing support to their families during the course of treatment.⁴

The consequences of the armed conflict that has occurred in Angola for about three decades has affected mainly the migration of the rural population to other regions, and the destruction of social infrastructures and traditional systems of economic activity.⁵ In addition, the destruction of roads has led to the shortage of basic and other essential goods, giving rise to dramatic humanitarian situations.⁶ In this sense, the government has focused its attention on the affected populations, with the support of international institutions, mainly the United Nations Organizations (UN), through the World Food Program (WFP).⁶ Throughout the 1990s, WFP provided Angola with logistical support for intervention among populations in critical areas.

Since 2002, after the signing of the peace agreement in Angola,⁷ WFP has developed a humanitarian aid plan for TB patients hospitalized at the Huambo Sanatorium hospital and those who were being treated at the Anti-Tuberculosis Dispensary (ATD) outpatient clinic. This aid lasted 6 years and consisted in the distribution of a basic food basket that patients took home once a month during the outpatient treatment period. Inpatients received the prepared food at the hospital.

The program's implementation had as its purpose increasing the adherence of TB patients to the Directly Observed Treatment (DOT) and improving the success rates of anti-tuberculosis treatment.⁸

Carmen Thais Lovera de Ferreira^{a,*}, Laura Serra Azuara^b, Oscar Asensio de la Cruz^a y María Montserrat Bosque García^a

^a *Unidad de Neurología y Alergia Pediátrica, Hospital Universitario Parc Tauli, Barcelona, España*
^b *Unidad de Ginecología y Obstetricia, Hospital Universitario Parc Tauli, Barcelona, España*

* Autor para correspondencia.
 Correo electrónico: thaislovera@hotmail.com
 (C.T. Lovera de Ferreira).

<https://doi.org/10.1016/j.arbres.2017.08.009>
 0300-2896/
 © 2017 SEPAR. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

Due to lack of funding, WFP was forced to end humanitarian aid in Angola by the end of 2006, ahead of schedule. The government then assumed responsibility for providing the basic food basket to patients through the Ministry of Planning and the Ministry of Social Assistance and Reintegration (MINARS) until 2008.⁹ Since 2009 there has been no further incentive for TB patients.

The objective of this study was to evaluate the effect of incentives (basic food basket) on the treatment outcome of patients with TB in the ATD, Huambo.

A retrospective study is carried out. The study included 19 739 cases of TB that were treated in the period between 2002 and 2014, in the ATD, Huambo. The study time was divided into two periods, the first one referring to the years 2002–2008, the period in which the basic baskets were distributed, and the second period referring to the years 2009–2014, during which there were no basic baskets. Data on the total TB cases and treatment outcomes were collected through the annual ATD reports, Huambo.

Treatment results were defined according to the WHO guidelines.¹⁰

The data available was analyzed through descriptive statistics. Mann–Whitney U test was used for group comparison.

The results of the treatment of 19 739 cases of TB were evaluated. In both periods, the median of therapeutic success (88% versus 60.5%, $P=0.01$), the median of failure (0.3% versus 0.85%, $P=0.02$) and the median of dropouts (7.1% versus 34.25%, $P=0.01$) were significantly different. No difference was found in the deceased (3.1% versus 3.7%, $P=0.73$). The results of the patients' treatment are shown in Table 1.

Our study showed that between 2002 and 2008, when basic food baskets were distributed to TB patients, the success rate of treatment was higher than in the period from 2009 to 2014, when there were no food baskets. With the removal of such incentives to patients, there was a considerable increase in treatment withdrawal. Similar to our study, a study conducted in Afghanistan found that the treatment success increased from the moment that the WFP introduced food assistance to TB patients.⁴ In Brazil, the provision of basic food baskets significantly increased adherence to TB treatment.¹¹ Other studies have also shown the positive impact of incentives on patients' adherence to TB treatment,^{12,13} taking into account the socio-economic difficulties experienced by these vulnerable groups.³

We assume that the incentive impact assessment in the outcome of treatment in our study was limited to the analysis of grouped data and that the sociodemographic and socioeconomic data of each patient would need to be assessed for a more complete analysis. However, our analysis strongly suggests the importance

Table 1
Results of the Patients' Treatment, Diagnosed With TB in the DAT, Huambo, During the Period From 2002 to 2014.

Year	Number of cases	Success (%)	Failure (%)	Deceased (%)	Abandonment (%)	Transferred (%)
<i>With basic basket</i>						
2002	1591	1421 (89.3)	4 (0.3)	53 (3.3)	110 (6.9)	3 (0.2)
2003	1694	1551 (91.6)	5 (0.3)	32 (1.9)	98 (5.8)	8 (0.5)
2004	1563	1395 (89.3)	7 (0.4)	41 (2.6)	111 (7.1)	9 (0.6)
2005	1664	1465 (88.0)	9 (0.5)	59 (3.5)	118 (7.1)	13 (0.8)
2006	1143	988 (86.4)	4 (0.3)	40 (3.5)	105 (9.2)	6 (0.5)
2007	1590	1328 (83.5)	3 (0.2)	50 (3.1)	200 (12.6)	5 (0.3)
2008	1567	1369 (87.4)	6 (0.4)	37 (2.4)	144 (9.2)	11 (0.7)
<i>Without basic basket</i>						
2009	1122	569 (50.7)	15 (1.3)	51 (4.5)	462 (41.2)	25 (2.2)
2010	1780	1042 (58.5)	24 (1.3)	76 (4.3)	617 (34.7)	21 (1.2)
2011	1871	1325 (70.8)	18 (1.0)	67 (3.6)	447 (23.9)	19 (1.0)
2012	1505	914 (60.7)	10 (0.7)	57 (3.8)	508 (33.8)	16 (1.1)
2013	1425	859 (60.3)	8 (0.6)	38 (2.7)	502 (35.2)	18 (1.3)
2014	1224	760 (62.1)	9 (0.7)	34 (2.8)	401 (32.8)	20 (1.6)

of incentives in patients' adherence to treatment, and therapeutic success against TB.

References

- World Health Organization. The economic impact of tuberculosis: the stop TB initiative. Geneva-Switzerland: WHO; 2000, 2000.
- Porto A. Social representations of tuberculosis: stigma and prejudice. *Rev Saude Publica*. 2007;41 Suppl. 1:43–9.
- Barter DM, Agboola SO, Murray MB, Barnighausen T. Tuberculosis and poverty: the contribution of patient costs in sub-Saharan Africa – a systematic review. *BMC Public Health*. 2012;12:980.
- Pedrazzoli D, Houben RM, Grede N, de Pee S, Boccia D. Food assistance to tuberculosis patients: lessons from Afghanistan. *Public Health Action*. 2016;6:147–53.
- Ministério do Planeamento. Estratégia de Combate à Pobreza - Reinserção Social, Reabilitação e Reconstrução e Estabilização Económica In: Direcção de Estudos e Planeamento editor. Luanda, Angola: 2004; 2003. 41.
- Governo de Angola. Estratégia Nacional de Segurança Alimentar e Nutricional (ENSAN) Luanda. 2009:87.
- Ferreira ME. Angola: conflict and development, 1961–2002. *Econ Peace Security J*. 2006;1:25–9.
- WHO. Guidelines Approved by the Guidelines Review Committee. Implementing the WHO Stop TB Strategy: A Handbook for National Tuberculosis Control Programmes. Geneva: World Health Organization Copyright (c) World Health Organization; 2008, 2008.
- World Food Program. After 30 years World Food Program ends food aid to Angola. United Nations. 2006.
- World Health Organization. Treatment of tuberculosis: guidelines. 4th ed. Geneva, Switzerland: WHO; 2010. p. 160.
- Cantalice Filho JP. Efeito do incentivo alimentício sobre o desfecho do tratamento de pacientes com tuberculose em uma unidade primária de saúde no município de Duque de Caxias Rio de Janeiro. *J Brasileiro Pneumologia*. 2009;35: 992–7.
- Ciobanu A, Domete L, Soltan V, Bivol S, Severin L, Plesca V, et al. Do incentives improve tuberculosis treatment outcomes in the Republic of Moldova? *Public Health Action*. 2014;4 Suppl. 2:S59–63.
- Garden B, Samarina A, Stavchanskaya I, Alsterlund R, Ovregaard A, Taganova O, et al. Food incentives improve adherence to tuberculosis drug treatment among homeless patients in Russia. *Scand J Caring Sci*. 2013;27:117–22.

Emília Santos^{a,b,*}, Óscar Felgueiras^{c,d}, Olena Oliveira^a, Raquel Duarte^{a,e,f}

^a EpiUnit – Instituto de Saúde Pública, Universidade do Porto, Porto, Portugal

^b Instituto Superior Politécnico da Universidade José Eduardo dos Santos, Huambo, Angola

^c Mathematics Department, Faculty of Sciences of the University of Porto, Porto, Portugal

^d Centre of Mathematics of the University of Porto, Porto, Portugal

^e Departamento de Pneumologia, Centro Hospitalar de Vila Nova de Gaia/Espinho, EPE, Vila Nova de Gaia, Portugal

^f Departamento de Ciências da Saúde Pública e Forenses e Educação Médica, Universidade do Porto, Porto, Portugal

* Corresponding author.

E-mail address: milalijanga@yahoo.com.br (E. Santos).

<https://doi.org/10.1016/j.arbres.2017.08.013>
0300-2896/

© 2017 SEPAR. Published by Elsevier España, S.L.U. All rights reserved.

Neumonía por *S. aureus* y artritis séptica esternoclavicular, una complicación insólita



S. aureus pneumonia and sternoclavicular septic arthritis: An unusual complication

Estimado Director:

Entre las complicaciones de la neumonía adquirida en la comunidad (NAC) raramente se producen artritis sépticas (AS). Suelen ser excepcionales, no simultáneas al proceso agudo^{1,2}, y con mayor frecuencia por *Streptococcus pneumoniae*^{1,3}. En estos casos las articulaciones más afectadas son la rodilla, la cadera y el hombro^{1,3}. La mayoría de las AS están producidas por *Staphylococcus aureus* y *Streptococcus* en pacientes con factores de riesgo predisponente¹⁻⁶. Presentamos un caso de neumonía por *S. aureus*, acompañada simultáneamente de una artritis séptica esternoclavicular (ASE) por

bacteriemia, en una mujer joven sin factores de riesgo conocidos, asociación no descrita previamente en la literatura.

Mujer de 60 años, no fumadora, sin antecedentes de inmunosupresión ni factores de riesgo neumológicos. Acudió a urgencias por dolor pleurítico y en hombro izquierdo, fiebre de 38 °C, tosin expectoración e intenso dolor en la región medioclavicular y el hombro izquierdo. La radiografía de tórax mostró tenue infiltrado en el lóbulo inferior izquierdo (LII). A la exploración destacaba semiología de condensación y egofonía en campo inferior izquierdo. Presentaba una zona abultada muy dolorosa no fluctuante, sin signos inflamatorios en piel, en la región esternoclavicular izquierda alta. Se realizó fibrobroncoscopia, que mostró signos inflamatorios, y se remitieron muestras para anatomía patológica y microbiología. Los hemocultivos fueron negativos. El ecocardiograma no mostró lesiones valvulares ni vegetaciones sugestivas de endocarditis. Se inició tratamiento empírico con ceftriaxona y levofloxacino, con escasa respuesta clínica, precisando