



Original article

Intra- and Inter-Physician Agreement in Therapeutic Decision for Sleep Apnea Syndrome

Vera M. Lugo^{a,1}, Marta Torres^{a,b,c,1}, Onintza Garmendia^{a,b,c}, Monique Suarez-Giron^a, Concepción Ruiz^a, Carmen Carmona^d, Eusebi Chiner^e, Nuria Tarraubella^f, Mireia Dalmases^{c,g}, Anna M. Pedro^h, Carlos J. Egea^{c,i}, Mireia Abellana^j, Mercè Mayos^{c,k}, Carmen Monasterio^{c,l,m}, Juan Fernando Masaⁿ, Ramón Farré^{c,o}, Josep M. Montserrat^{a,b,c,p,*}

^a Sleep Unit, Hospital Clínic, Barcelona, Spain

^b IDIBAPS, Barcelona, Spain

^c CIBER Enfermedades Respiratorias, Madrid, Spain

^d Medical-Surgical Unit of Respiratory Diseases, Virgen del Rocío University Hospital, Sevilla, Spain

^e Unidad multidisciplinar de Sueño, Servicio de Neumología, H.U. Sant Joan d'Alacant, Alacant, Spain

^f Atenció Primària Àmbit Lleida, Lleida, Spain

^g Hospital Universitari Arnau de Vilanova and Santa Maria, Group of Translational Research in Respiratory Medicine, IRB Lleida, Spain

^h ABS Gaudí, CAP Sagrada Família, Consorci Sanitari Integrat (CSI), Barcelona, Spain

ⁱ Sleep Disorders Unit, Hospital Txagorritxu, Vitoria, Spain

^j CAP Guissona, Àrea Cervera, Lleida, Spain

^k Sleep Unit, Department of Respiratory Medicine, Hospital de la Santa Creu i Sant Pau, Barcelona, Spain

^l Sleep Unit, Department of Respiratory Medicine, Hospital Universitari de Bellvitge, L'Hospitalet de Llobregat, Spain

^m IDIBELL, L'Hospitalet de Llobregat, Spain

ⁿ Respiratory Unit, Hospital San Pedro de Alcántara, Cáceres, Spain

^o Unitat de Biofísica i Bioenginyeria, Facultat de Medicina, Universitat de Barcelona, IDIBAPS, Barcelona, Spain

^p Universitat de Barcelona, Barcelona, Spain

ARTICLE INFO

Article history:

Received 21 December 2018

Accepted 21 February 2019

Available online 5 April 2019

Keywords:

Sleep apnea

CPAP treatment

Therapy agreement

ABSTRACT

Background: Large variation in diagnostic procedures and treatment recommendations may hinder the management of obstructive sleep apnea (OSA) and also compromise correct interpretation of the results of multicenter clinical trials, especially in subjects with non-severe OSA. The aim of this study was to analyze the therapeutic decision-making between different sleep physicians in patients with AHI < 40 events/h.

Methods: Six experienced senior sleep specialists from different sleep centers of Spain were asked to make a therapeutic decision (CPAP treatment) based on anonymized recordings of patients with suspected OSA that has previously performed a sleep study. The clinical data was shown in an online database and included anthropometric features, clinical questionnaires, comorbidities, physical examination and sleep study results. Intra- and inter-observer decision-making were analyzed by the Fleiss' Kappa statistics (Kappa).

Results: A total of 720 medical decisions were taken to analyze the agreement between sleep professionals. Overall intra-observer evaluation reliability was almost perfect (Kappa = 0.83, 95% CI, 0.75–0.90, $p < 0.001$). However, overall inter-observer concordance decreased to moderate agreement (Kappa = 0.46, 95% CI, 0.42–0.51, $p < 0.001$). Nevertheless, it was especially low when considering AHI < 15 events/h.

Conclusions: This study demonstrates a good intra-observer concordance in the therapeutic decision-making of different sleep physicians treating patients with low/moderate OSA. However, when analyzing inter-observer agreement the results were considerably worse. These findings underline the importance of developing improved consensus management protocols.

© 2019 Published by Elsevier España, S.L.U. on behalf of SEPAR.

* Corresponding author.

E-mail address: jmmontserrat@ub.edu (J.M. Montserrat).

¹ Equally contributed.

Concordancia intra e inter-médico en la decisión terapéutica para el síndrome de apnea del sueño

R E S U M E N

Palabras clave:
Apnea del sueño
Tratamiento CPAP
Terapia consensuada

Introducción: La gran variedad de procedimientos diagnósticos y recomendaciones de tratamiento puede dificultar el manejo del síndrome de apnea obstructiva del sueño (SAHS), y del mismo modo comprometer la correcta interpretación de los resultados de ensayos clínicos multicéntricos, especialmente en pacientes con SAHS no grave. El objetivo de este estudio fue analizar la decisión terapéutica de distintos médicos expertos en sueño en pacientes con el índice de apnea hipopnea < 40 eventos/h.

Métodos: Se pidió a seis especialistas con amplia experiencia en sueño de diferentes centros de España que tomaran una decisión terapéutica (terapia de presión positiva continua en las vías respiratorias o CPAP) basada en datos anónimos de los pacientes con sospecha de SAHS en los que previamente se había llevado a cabo un estudio del sueño. Los datos clínicos procedían de una base de datos online e incluían características antropométricas, cuestionarios clínicos, comorbilidades, examen físico y resultados del estudio del sueño. La concordancia intra- e interobservador de la toma de decisiones se analizó mediante el estadístico Fleiss' Kappa (Kappa).

Resultados: Se analizaron un total de 720 decisiones médicas para evaluar el consenso entre profesionales del sueño. De manera global, la fiabilidad de la evaluación intraobservador fue casi perfecta (Kappa = 0,83; 95% CI; 0,75 a 0,90, $p < 0,001$). Sin embargo, la concordancia global interobservador disminuyó hasta alcanzar un grado moderado de consenso (Kappa = 0,46; 95% CI; 0,42 a 0,51, $p < 0,001$), que fue especialmente bajo cuando se tuvo en cuenta un índice de apnea hipopnea < 15 eventos/h.

Conclusiones: Este estudio demuestra una buena concordancia intraobservador en la toma de decisiones terapéuticas de distintos médicos expertos en sueño que tratan a pacientes con SAHS leve o moderado. Sin embargo, los resultados relativos al acuerdo interobservador fueron notablemente peores. Estos hallazgos señalan la importancia de desarrollar mejores protocolos consensuados de manejo.

© 2019 Publicado por Elsevier España, S.L.U. en nombre de SEPAR.

Introduction

Obstructive sleep apnea (OSA) is one of the most common sleep disorders. It is a chronic condition secondary to complete or partial upper airway obstruction that results in daytime sleepiness and fatigue that negatively affects the quality of life of patients. OSA is also considered an important risk factor for cardiovascular, metabolic and neurological comorbidities.^{1,2}

OSA is diagnosed by using different sleep tests, from conventional polysomnography (PSG), which is actually the gold standard in diagnostic procedures, to most simplified single-channel home devices.³ Despite the proven efficacy and agreement of simplified devices with PSG, sometimes the clinical management of OSA can be hampered by different interpretations of the data from sleep studies. This fact is critical when dealing with patients with non-severe OSA, in whom the diagnosis and the subsequent therapeutic decision could be especially difficult.

There are several guidelines, consensus and research articles with various indications for CPAP treatment in OSA based on AHI severity,⁴ excessive daytime sleepiness,^{5,6} or improved driving performance.⁷ Other recommendations also include hypertension and cardiovascular comorbidities, regardless of OSA symptoms,⁸ impaired cognition, insomnia or mood disorders.⁹ In Spain the criteria for recommending CPAP are, as stated in the Spanish Sleep Network guidelines,¹⁰ an AHI between 5 and 30 events/h with significant symptoms or OSA associated pathologies, or AHI greater than 30 events/h, giving less importance to symptoms or pathologies.

Therefore, since different professionals may vary in interpreting the results of sleep studies, CPAP indication could be particularly complex when dealing with non-severe patients and different diagnostic methodologies. Thus, clinical decision-making in this group of OSA patients may be discordant and affect its management. This fact is also especially relevant since interpretation of results can be compromised in the context of clinical trials, especially those

involving several centers and/or countries. Due to the high number of multicenter clinical trials that are carried out at present, it is worth to analyze the consistency of decision-making concerning recommendation of CPAP among different sleep physicians, especially in case of patients with AHI < 30 events/h, when using PSG or home respiratory polygraphy (RP), as usually occurs in a sleep unit.

The aim of this study was to compare the intra- and inter-observer variability in decision-making on treatment recommendation in patients with AHI < 40 events/h among different Spanish sleep physicians.

Methods

To evaluate intra- and inter-observer agreement in therapeutic decision-making 6 sleep professionals, were asked to carry out an online therapeutic decision to recommend or not CPAP treatment based on anonymized records from 40 patients that were displayed on an online-encrypted database. These records included anthropometric data, clinical questionnaires, comorbidities, physical examination and the sleep study results, from either PSG or RP. The 6 professionals involved were experienced sleep physicians, working in different Sleep Units of Spanish University Hospitals, aged 36–60 years-old. All were recommended to follow the SEPAR guidelines for therapeutic decision making.

The study was carried out with the approval of the Hospitals' Ethics Committee. All data were obtained from patients recruited from May 2015 to May 2017 in the Sleep Unit of Hospital Clinic (Barcelona, Spain) and included in a clinical trial (trial number NCT02779894 in <http://www.clinicaltrials.gov/>). Forty patients were randomly selected from those in the study database presenting AHI < 40 events/h. All subjects were aged 18–75 years old and, before their sleep test, presented suspicion of OSA (heavy snoring with breathing pauses during the night, non-restful sleep and daytime somnolence or fatigue not explained by other pathologies) and/or refractory hypertension. None of them presented

invalidating somnolence (medical criteria), any unstable diseases, previous use of CPAP, uvulopalatopharyngoplasty, or risk profession.

The sleep studies corresponded to in-hospital PSG, in-hospital RP or home RP. In in-hospital PSG (Grael/Somt  PSG, Compumedics Limited 2006, Abbotsford, Victoria, Australia) the signals of electroencephalogram (EEG- leads F4-M1, C4-M1 and O2-M1), electrooculogram (EOG) chin and leg electromyogram (EMG), and electrocardiogram (ECG) were recorded, as well as nasal and oronasal flow (cannula and thermistor), respiratory effort, and oxygen saturation (SpO₂), body position, snoring (tracheal microphone). All patients were also video monitored. In-hospital PR (Somt  PSG, Compumedics Limited 2006, Abbotsford, Victoria, Australia) consisted in recording nasal flow (cannula), snoring, respiratory effort, SpO₂ and body position, as well as video monitoring. In home RP (Portable type 3 ApneaLink air, ResMed, Australia) the registered variables were nasal flow (cannula), snoring, respiratory effort, SpO₂, pulse frequency, and body position. Sleep scoring was performed using the standardized AASM criteria¹¹: apnea was defined as an absence of flow for more than 10 s. Hypopnea was defined as a discernible reduction in the amplitude of the airflow signal from the pre-event baseline for at least 10 s, associated with an oxygen desaturation $\geq 3\%$ in both PSG and PR and also associated with arousal in PSG.

Online procedure

To carry out the therapeutic decision, all professionals logged-in into a especially designed website with individual username and password. Professionals could see a list of patients identified by an anonymous alphanumeric code. To evaluate the intra-individual agreement, each of the patients was visualized 3 independent times, with a different code each, at random order and without knowledge of the evaluators. Therefore, each professional evaluated 120 cases. In order to avoid fatigue, each evaluator was allowed to analyze a maximum of 15 cases per day along a 3-month period.

When entering into each case evaluation, the data provided for therapeutic decision-making were displayed. They included anthropometric features (age, gender, body mass index (BMI), and systolic and diastolic blood pressure), clinical questionnaires (Epworth sleepiness scale (ESS), EuroQol-5D and EuroQol-VAS), comorbidities (high blood pressure, cardiac disease, neurological or respiratory diseases, diabetes, dyslipidemia, depression, anxiety or neoplasms), and physical examination data (micrognathia, retrognathia, amygdala and palatine Friedman class, nasal obstruction or ORL surgery). To help in analyzing the sleep history of the case, the following data were also displayed: ASDA sleepiness scale, daily amount of sleep hours, presence of snoring, choking attacks, nicturia, witnessed apneas, morning headache, restless sleep, daytime sleepiness, restless legs syndrome, aggressive behavior during sleep, and muscle weakness associated to intense emotions or sleepwalking. The results of the sleep studies included recorded time, AHI, central apneas, total respiratory events, Cheyne–Stokes breathing, postural predominance, basal and mean SpO₂, ODI 3%, and CT90 for all tests. In case of PSG sleep efficiency, sleep staging, arousal index, and postural ODI were included. Based on all the information provided, sleep professionals were asked to choose a therapeutic decision (CPAP/non-CPAP) if an OSA diagnosis was attributed to the case.

Statistical analysis

To assess observer reliability on the therapeutic decision-making (CPAP treatment or not) between the same and the different physicians (intra- and inter-individual agreement), the Fleiss' Kappa (Kappa) statistics for categorical variables¹² was




Fig. 1. CPAP indication (%) of the three therapeutic decisions for each actual patient made by each evaluator.

calculated. Data are presented as mean \pm standard deviation for the variables measured in numerical scale and in percentage for the measurement in nominal scale.

Percentage of CPAP indication of the three therapeutic decisions for each actual patient made by each evaluator (Fig. 1) was also analyzed by 2-way ANOVA, taking “time of therapeutic decision” and “sleep physician” as factors.

Results

Table 1 summarizes the patients' demographics and comorbidities of the sleep studies analyzed. Out of 40 sleep studies, 57.5% were from male patients, mean aged 51.07 ± 11.70 years and 28.13 ± 8.04 kg/m². Mean AHI was 18.42 ± 11.32 events/h, ranging 0–40. Concerning diurnal somnolence, patients presented a mean ESS of 8.02 ± 4.56 , ranging 0–22. Quality of life questionnaires showed mean EuroQol-5D of 0.84 ± 0.21 and EuroQol-VAS of 71.37 ± 19.18 , ranging 0.17–1 and 10–100, respectively.

Table 1
General patients characteristics (n=40).

Male gender	23 (57.35)
Mean age (years)	51.07 ± 11.70
Neck circumference (cm)	38.87 ± 3.79
BMI (kg/m ²)	28.13 ± 8.04
Nasal obstruction	15 (37.5)
ORL surgery	7 (17.5)
Smokers	10 (25.0)
Alcohol intake	21 (52.5)
Comorbidities	
Hypertension	10 (25.0)
Diabetes mellitus	7 (17.5)
Dislipidemia	23 (57.5)
Cardiovascular disease	3 (7.5)
Neurological disease	6 (15)
Respiratory disease	7 (17.5)
Depression	9 (22.5)
Anxiety	8 (20.0)
Cancer	4 (10.0)
AHI (events/h)	18.42 ± 11.32
CT90 (%)	5.05 ± 8.43
ODI3%	18.90 ± 14.27
EuroQol-5D	0.84 ± 0.21
EuroQol-VAS	71.37 ± 19.18
ESS	8.02 ± 4.56

Data are expressed as mean \pm SD or number of patients (%). BMI: body mass index. AHI: apnea hypopnoea index. ODI: oxygen desaturation index. QoL: quality of life. VAS: visual analog scale. ESS: Epworth sleepiness scale.

Table 2
On-line diagnosis and CPAP indication.

Total medical decisions	720
Diagnosis	
OSA	70.9%
Snoring	21.5%
Other*	7.6%
CPAP indication	36.0%

* Absence of pathology or non-OSA sleep pathologies.

Table 3
Fleiss' Kappa correlation for intra-physician therapeutic decision-making.

	Kappa	SE	95% CI	p Value
All AHI	0.83	0.04	0.75–0.90	>0.001
AHI < 15	0.81	0.06	0.70–0.93	>0.001
AHI > 15	0.78	0.05	0.68–0.87	>0.001

Kappa: Fleiss' Kappa statistic. SE: standard error. CI: confidence interval.

Table 4
Fleiss' Kappa correlation for inter-physician therapeutic decision-making.

	Kappa	SE	95% CI	p Value
All AHI	0.46	0.02	0.42–0.51	>0.001
AHI < 15	0.06	0.04	–0.01 to 0.14	0.095
AHI > 15	0.37	0.03	0.31–0.43	>0.001

Kappa: Fleiss' Kappa statistic. SE: standard error. CI: confidence interval.

Therefore, the sleep studies analyzed corresponded to a population of patients with wide ranges of sleepiness and quality of life.

As every actual sleep study was blindly analyzed 3 times by the 6 physicians, each case was analyzed 18 times, and a total of 720 medical decisions were made for assessing the concordance among health professionals. Table 2 shows the diagnosis and medical decisions chosen by all the health professionals that participated in the study.

Fleiss' Kappa agreement for intra-observer decision-making is represented in Table 3. Overall intra-observer evaluation reliability of sleep professionals was found to be good, showing almost perfect agreement (Kappa = 0.83, 95% CI: 0.75–0.90, $p < 0.001$). Concordance was also assessed considering two different categories of AHI. For AHI < 15 events/h sleep physicians showed again an almost perfect agreement (Kappa = 0.81, 95% CI: 0.70–0.93, $p < 0.001$) while for AHI > 15 events/h the agreement was found to be substantial (Kappa = 0.78, 95% CI: 0.68–0.87, $p < 0.001$).

As indicated in Table 4, Fleiss' Kappa agreement clearly decreased when assessing inter-observer agreement. Considering all AHI, inter-observer evaluation reliability was found to be moderate (Kappa = 0.46, 95% CI: 0.42–0.51, $p < 0.001$). For AHI < 15 events/h the worst values of concordance were found: agreement could be not demonstrated between sleep physicians (Kappa = 0.06, 95% CI: –0.01 to 0.14, $p = 0.095$). When analyzing AHI > 15, inter-observer agreement slightly increased, and was found to be fair (Kappa = 0.37, 95% CI: 0.31–0.43, $p < 0.001$).

Fig. 1 shows the percentage of CPAP indication of the 3 therapeutic decisions for each actual patient made by each evaluator. The results of 2-way ANOVA were consequent with intra- and inter-observer agreement analysis. Whereas the time of analysis factor was not significant ($p = 0.346$), a significant effect of physician was observed ($p < 0.001$).

Discussion

This study analyzed intra- and inter-observer agreement in the therapeutic decision-making among Spanish sleep physicians treating patients with AHI < 40 events/h. The results, obtained from more than seven hundred evaluations ($n = 720$) demonstrated a

good intra-observer concordance. However, regarding to inter-individual agreement this study revealed lower concordance: overall concordance was found to be moderate but in case of less severe patients (AHI < 15 events/h) the inter-subject concordance considerably worsened.

Despite the existence of a number of clinical guidelines and consensus for the management of OSA, these results show a level of concordance below it is desirable. The majority of available reports have not analyzed the concordance of therapeutic decision-making between different sleep physicians, but between different diagnostic procedures, usually comparing simplified devices versus the gold standard in-hospital PSG. Indeed, Masa et al.¹³ analyzed the therapeutic decision-making agreement between home RP and in-hospital PSG among 348 patients, describing that home RP was adequate for high AHI, but insufficient for mild or moderate AHI and concluded that this diagnostic method was effective only in patients with high pretest of OSA. Similar results have been obtained in a clinical study carried out in a pediatric population using in-lab RP versus PSG,¹⁴ and the authors concluded that clinical decision-making in children with mild and moderate OSA may be difficult. Studying an adult population, Guerrero et al.¹⁵ tried to go a step further and analyzed the agreement of therapeutic decision making after three consecutive nights with home RP versus PSG in patients with mild to moderate pretest of OSA or with associated comorbidities that could mask OSA symptoms. In this study, the authors also assessed the concordance between different specialists, comparing the therapeutic decisions made by sleep physicians and respiratory physicians. The authors demonstrated that this particular diagnostic method is useful to manage patients without high pretest probability of OSA or with comorbidities only when evaluated by a qualified sleep specialist. It is important to bear in mind that these and other clinical studies have also shown that these diagnostic approaches, in addition to effectively enabling a correct decision making to prescribe therapy for OSA in certain patient population, also reduced costs.¹⁶ A different study¹⁷ assessed both diagnosis and therapeutic decision agreement between personnel from sleep reference centers with PSG and from non-reference centers employing a simplified device. The authors found a substantial level of concordance between different professionals, but each group of professionals employed a different diagnostic method.

Our study is unique since, as far as we know, there are no published studies addressed to analyze the performance of CPAP therapeutic decision-making in OSA among sleep specialists in case that all of them analyze the same patient cases, using different real-life diagnostic methods. In fact, the available data are focused on comparing how respiratory events or sleep stage are scored.^{18–21} The causes that could explain the lack of concordance we found are not well known. However, we could speculate that, although it was recommended to follow the SEPAR regulations, it is possible that each evaluator has also used their own criteria by taking into account other circumstances (gender, quality of life, different pathologies, or only sleepiness). In addition, the lack of a clinical face-to-face interview could have conditioned the therapeutic decision proposed by the 6 professionals involved in the study. Moreover, it is remarkable that current consensus are global and have little impact on defining the management of mild/moderate cases. In order to simplify the study and to focus on the most relevant therapeutic decision (CPAP prescription) other aspects usual in clinical practice, such as decision to refer patients to other specialty doctors (e.g. otorhinolaryngologist or neurologist), have not been analyzed.

By showing the striking and worrying results we have found, especially taking into account that our 6 sleep physicians had solid experience in the field, we intend to draw the attention of the sleep community. There are several possible consequences of

this relatively poor agreement between sleep professionals. On the one hand, OSA management could differ especially of those patients with low AHI. On the other hand, the interpretation of the many multicentre studies that are currently on-going may be hampered by the lack of agreement we report here. The results found in different decision-making in other medical specialties regarding the agreement between specialist physicians have also revealed the need to seek strategies to improve inter-observer reliability.^{22–25} Thus, it seems advisable that clinical guidelines for the management of OSA need revision, in particular concerning recommendations to manage OSA patients with low AHI.

Financial

The Spanish Ministry of Economy and Competitiveness (PI14/00416 and PI17/01068).

Conflict of interest

The authors declare no conflict of interest.

Acknowledgements

The authors would thank Mr A. Gabarrús for his statistical assistance, Mr R. Pereira for implementing the designed on-line database and Ms G. Guerrero for her participation in the preparation of the patients' database.

References

1. Peppard PE, Young T, Barnet JH, Palta M, Hagen EW, Hla KM. Increased prevalence of sleep-disordered breathing in adults. *Am J Epidemiol*. 2013;177:1006–14.
2. Dempsey JA, Veasey SC, Morgan BJ, O'Donnell CP. Pathophysiology of sleep apnea. *Physiol Rev*. 2010;90:47–112.
3. Masa JF, Duran-Cantolla J, Capote F, Cabello M, Abad J, Garcia-Rio F, et al. Spanish sleep network efficacy of home single-channel nasal pressure for recommending continuous positive airway pressure treatment in sleep apnea. *Sleep*. 2015;38:13–21.
4. Kushida CA, Morgenthaler TI, Littner MR, Alessi CA, Bailey D, Coleman J Jr, et al. American Academy of Sleep Practice parameters for the treatment of snoring and obstructive sleep apnea with oral appliances: an update for 2005. *Sleep*. 2006;29:240–3.
5. McDaid C, Durée KH, Griffin SC, Weatherly HL, Stradling JR, Davies RJ, et al. A systematic review of continuous positive airway pressure for obstructive sleep apnoea-hypopnoea syndrome. *Sleep Med Rev*. 2009;13:427–36.
6. Kribbs NB, Pack AI, Kline LR, Getsy JE, Schuett JS, Henry JN, et al. Effects of one night without nasal CPAP treatment on sleep and sleepiness in patients with obstructive sleep apnea. *Am Rev Respir Dis*. 1993;147:1162–8.
7. Strohl KP, Brown DB, Collop N, George C, Grunstein R, Han F, et al. ATS Ad Hoc Committee on Sleep Apnea Sleepiness, and Driving Risk in Noncommercial Drivers. An official American Thoracic Society clinical practice guideline: sleep apnea, sleepiness, and driving risk in noncommercial drivers An update of a 1994 statement. *Am J Resp Crit Care Med*. 2013;187:1259–66.
8. Young T, Peppard P, Palta M, Hla KM, Finn L, Morgan B, et al. Population-based study of sleep-disordered breathing as a risk factor for hypertension. *Arch Intern Med*. 1997;157:1746–52.
9. Naëgelé B, Thouvard V, Pépin JL, Lévy P, Bonnet C, Perret JE, et al. Deficits of cognitive executive functions in patients with sleep apnea syndrome. *Sleep*. 1995;18:43–52.
10. Grupo Español de Sueño (GES). Consenso Nacional sobre el síndrome de apneas-hipopneas del sueño. *Arch Bronconeumol*. 2005;41:3–110.
11. Berry RB, Brooks R, Gamaldo CE, Harding SM, Marcus CL, Vaughn BV, for the American Academy of Sleep Medicine. Darien, IL: American Academy of Sleep Medicine. The AASM Manual for the Scoring of Sleep and Associated Events: Rules, Terminology and Technical Specifications; 2012. Version 2.0. www.aasmnet.org.
12. Landis JR, Koch GG. The measurement of observer agreement for categorical data. *Biometrics*. 1977;33:159–74.
13. Masa JF, Corral J, Pereira R, Duran-Cantolla J, Cabello M, Hernández-Blasco L, et al. Spanish Sleep Network Therapeutic decision-making for sleep apnea and hypopnea syndrome using home respiratory polygraphy: a large multicentric study. *Am J Respir Crit Care Med*. 2011;184:964–71.
14. Tan HL, Gozal D, Ramirez HM, Bandla HP, Kheirandish-Gozal L. Overnight polysomnography versus respiratory polygraphy in the diagnosis of pediatric obstructive sleep apnea. *Sleep*. 2014;37:255–60.
15. Guerrero A, Embid C, Isetta V, Farre R, Duran-Cantolla J, Parra O, et al. Management of sleep apnea without high pretest probability or with comorbidities by three nights of portable sleep monitoring. *Sleep*. 2014;37:1363–73.
16. Parra O, García-Escasans N, Montserrat JM, García Eroles L, Ruíz J, López JA, et al. Should patients with sleep apnoea/hypopnoea syndrome be diagnosed and managed on the basis of home sleep studies? *Eur Respir J*. 1997;10:1720–4.
17. Hernández L, Torrella M, Roger N, Lluell A, Ballester E, Quinto L, et al. Management of sleep apnea: concordance between nonreference and reference centers. *Chest*. 2007;132:1853–7.
18. Rosenberg RS, Van Hout S. The American Academy of Sleep Medicine inter-scoring reliability program: sleep stage scoring. *J Clin Sleep Med*. 2013;9:81–7.
19. Arnardottir ES, Verbraecken J, Gonçalves M, Gjerstad MD, Grote L, Puertas FJ, et al. The National Representatives as Study Collaborators Variability in recording and scoring of respiratory events during sleep in Europe: a need for uniform standards. *J Sleep Res*. 2016;25:144–57.
20. Collop NA. Scoring variability between polysomnography technologists in different sleep laboratories. *Sleep Med*. 2002;3:43–7.
21. Manser RL, Rochford P, Naughton MT, Pierce RJ, Sasse A, Teichtahl H, et al. Measurement variability in sleep disorders medicine: the Victorian experience. *Int Med J*. 2002;32:386–93.
22. van den Boom AL, de Wijckerslooth EML, Mauff KAL, Dawson I, van Rossem CC, Toorenvliet BR, et al. Interobserver variability in the classification of appendicitis during laparoscopy. *Br J Surg*. 2018;105:1014–9.
23. Piili RP, Lehto JT, Luukkaala T, Hinkka H, Kellokumpu-Lehtinen PI. Does special education in palliative medicine make a difference in end-of-life decision-making? *BMC Palliat Care*. 2018;17:94.
24. Duignan S, Ryan A, O'Keefe D, Kenny D, McMahon CJ. Prospective analysis of decision making during Joint Cardiology Cardiothoracic Conference in treatment of 107 consecutive children with congenital heart disease. *Pediatr Cardiol*. 2018;12 [Epub ahead of print].
25. Andrade R, Espinoza M, Gómez EM, Espinoza JR, Cruz E. Intra- and inter-examiner reproducibility of manual probing depth. *Braz Oral Res*. 2012;26:57–63.