



Editorial

Segmentectomy in Lung Cancer as a New Standard of Treatment: Can Less Be More?



Non-small cell lung cancer (NSCLC) remains the leading cause of cancer death worldwide,¹ especially in patients with chronic respiratory diseases such as chronic obstructive pulmonary disease (COPD). In recent decades, however, efforts to prevent, diagnose, and treat NSCLC have led to a decline in age-adjusted mortality rates (especially in men)² and the detection of smaller tumors at the time of diagnosis.

Today, the treatment of choice for small NSCLC tumors is surgical resection by lobectomy combined with lymph node dissection,³ preferably by video-assisted thoracic surgery (VATS). NSCLC treatment practice guidelines recommend the use of sublobar resections such as segmentectomy only in patients who would not tolerate lobectomy due to poor lung function or comorbidities.

However, the results of the JCOG0802 study, a multicenter randomized clinical trial in a Japanese population of patients with stage IA NSCLC ≤ 2 cm and a consolidation-to-tumor ratio > 0.5 cm have recently been published.⁴ In this phase III trial, the authors randomized more than 1000 participants to a segmentectomy arm or a lobectomy arm, with the primary objective of demonstrating the non-inferiority of overall survival after segmentectomy versus lobectomy. After a mean follow-up of more than 7 years, the authors found that segmentectomy was superior to lobectomy in overall 5-year survival (94.3% vs 91.1, HR 0.663, $p < 0.005$), and showed a similar 5-year recurrence-free survival rate (88.0% vs 87.9%) and a higher rate of local recurrences (10.5% vs 5.4%). Surprisingly, improved survival in the segmentectomy group appeared to be due to a decrease in deaths from a second cancer and, to a lesser extent, from non-cancer diseases.

The results of this study have generated great interest among the scientific community since it was first reported in 2021, but doubts have also emerged surrounding the resulting paradigm shift in the surgical treatment of NSCLC. Is this study sufficiently robust to prompt the medical community and stakeholders to change their guidelines and recommend segmentectomy as the procedure of choice for the surgical treatment of NSCLC? Several previously published studies support the benefit of segmentectomy versus lobectomy reported by Saji and colleagues.

A retrospective study⁵ of 312 patients with clinical stage IA NSCLC published in 2016 compared outcomes of patients undergoing segmentectomy or pulmonary lobectomy, and found similar 5-year and 10-year survival in both groups. Specifically, oncologic outcomes after segmentectomy were slightly better than after lobectomy, with the first group obtaining 5- and 10-year survival rates of 97.5% and 83.5%, respectively, compared with 87.75% and 75.0% in the lobectomy group.

In 2017, a systematic review based on retrospective data from 24,542 patients found that survival in patients with stage I NSCLC who underwent therapeutic segmentectomies did not vary from patients who underwent lobectomies, with a hazard ratio (relative risk) of 1.04 (95% confidence interval 0.92–1.18; $p = 0.50$).⁶

A more recent systematic review and meta-analysis that included 16 studies concluded that lobectomy has an overall survival similar to segmentectomy and superior to wedge resection. These results led the authors to suggest that segmentectomy should be positioned as a valuable alternative for patients with NSCLC smaller than 2 cm, and that wedge resection should be avoided whenever possible.⁷

Finally, a recently published retrospective analysis of data from a reference center in the USA concluded that there were no differences in clinical outcomes (5-year mortality, recurrence, or recurrence-free survival) between anatomic segmentectomy and lobectomy.⁸

Based on the evidence from the above studies and from the JCOG0802 study group, it seems clear that segmentectomy should be positioned as the surgical procedure of choice for patients with early-stage NSCLC, as it offers better survival than lobectomy and greater preservation of pulmonary parenchyma, despite a greater proportion of local relapses in the follow-up. Systematic nodal dissection can be performed with segmentectomy in a manner similar to that of lobectomy.^{9–11}

The findings of this study are only applicable to peripheral NSCLC measuring less than 2 cm in diameter, but this is precisely the area where diagnostic rates are increasing, especially as a result of lung cancer screening programs in at-risk populations.¹² Furthermore, as the target population for lung cancer screening consists predominantly of patients with COPD and previous emphysema, procedures of this type will be performed more frequently because patients who otherwise would not be eligible for lobectomy can be included.

However, it is also important to bear in mind that segmentectomy is a technically more complex procedure than lobectomy and requires more prolonged surgery.¹³ This means that more operating rooms will be needed to allow hospitals to continue offering patients the shortest possible surgery waiting time.

In conclusion, the results of the JCOG0802 study denote a clear change of mindset in the treatment of early-stage NSCLC (stage IA), reinforcing the idea that at least in this type of tumor, resection of a smaller amount of tissue may improve life expectancy. The details of this study still have to be thoroughly analyzed, but we

hope that this new evidence will soon be incorporated into the clinical practice guidelines. Perhaps a new day has dawned.

Conflict of interests

JJDV declares no conflicts of interest.

GRR declares no conflicts of interest.

BAN has received payments for scientific consultancy and/or activities in the last 3 years from AstraZeneca, Laboratories Bial, Boehringer Ingelheim, Chiesi, Gilead, GSK, Laboratories Menarini, MSD, and Novartis AG.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at doi:10.1016/j.arbres.2022.05.015.

References

- Dyba T, Randi G, Bray F, Martos C, Giusti F, Nicholson N, et al. The European cancer burden in 2020: incidence and mortality estimates for 40 countries and 25 major cancers. *Eur J Cancer*. 2021;157:308–47.
- Cayuela L, López-Campos JL, Otero R, Rodríguez Portal JA, Rodríguez-Domínguez S, Cayuela A. The beginning of the trend change in lung cancer mortality trends in Spain, 1980–2018. *Arch Bronconeumol*. 2021;57:115–21.
- Howington JA, Blum MG, Chang AC, Balekian AA, Murthy SC. Treatment of stage I and II non-small cell lung cancer: Diagnosis and management of lung cancer, 3rd ed: American College of Chest Physicians evidence-based clinical practice guidelines. *Chest*. 2013;143.
- Saji H, Okada M, Tsuboi M, Nakajima R, Suzuki K, Aokage K, et al. Segmentectomy versus lobectomy in small-sized peripheral non-small-cell lung cancer (JCOG0802/WJOG4607L): a multicentre, open-label, phase 3, randomised, controlled, non-inferiority trial. *Lancet*. 2022;399:1607–17.
- Kodama K, Higashiyama M, Okami J, Tokunaga T, Imamura F, Nakayama T, et al. Oncologic outcomes of segmentectomy versus lobectomy for clinical T1a NO MO non-small cell lung cancer. *Ann Thorac Surg*. 2016;101:504–11.
- Bedetti B, Bertolaccini L, Rocco R, Schmidt J, Solli P, Scarci M. Segmentectomy versus lobectomy for stage I non-small cell lung cancer: a systematic review and meta-analysis. *J Thorac Dis*. 2017;9:1615–23.
- Shi Y, Wu S, Ma S, Lyu Y, Xu H, Deng L, et al. Comparison between wedge resection and lobectomy/segmentectomy for early-stage non-small cell lung cancer: a Bayesian meta-analysis and systematic review. *Ann Surg Oncol*. 2022;29:1868–79.
- Chan EG, Chan PG, Mazur SN, Normolle DP, Luketich JD, Landreneau RJ, et al. Outcomes with segmentectomy versus lobectomy in patients with clinical T1cN0M0 non-small cell lung cancer. *J Thorac Cardiovasc Surg*. 2021;161, 1639–48.e2.
- Okada M, Yoshikawa K, Hatta T, Tsubota N. Is segmentectomy with lymph node assessment an alternative to lobectomy for non-small cell lung cancer of 2 cm or smaller? *Ann Thorac Surg*. 2001;71:956–60.
- Mattioli S, Ruffato A, Puma F, Daddi N, Aramini B, D'ovidio F. Does anatomical segmentectomy allow an adequate lymph node staging for cT1a non-small cell lung cancer? *J Thorac Oncol*. 2011;6:1537–41.
- Xiao F, Yu Q, Zhang Z, Liu D, Guo Y, Liang C, et al. Novel perspective to evaluate the safety of segmentectomy: clinical significance of lobar and segmental lymph node metastasis in cT1N0M0 lung adenocarcinoma. *Eur J Cardio-Thorac Surg*. 2018;53:228–34.
- Seijo LM, Trujillo JC, Zulueta JJ. Screening in lung cancer: the latest evidence. *Arch Bronconeumol*. 2020;56:7–8.
- Mesa-Guzmán M, González J, Alcaide AB, Bertó J, de-Torres JP, Campo A, et al. Surgical outcomes in a lung cancer-screening program using low dose computed tomography. *Arch Bronconeumol*. 2021;57:101–6.

Jesus Isea de la Viña^a, Gema Ramírez-Rodríguez^{b,*},
Bernardino Alcázar-Navarrete^{b,c,d}

^a Thoracic Surgery Unit, University Hospitals Coventry and
Warwickshire, Coventry, United Kingdom

^b Servicio de Neumología, Hospital Universitario Virgen de las Nieves,
Granada, Spain

^c Departamento de Medicina, Universidad de Granada, Spain

^d IBS-Granada, Spain

Corresponding author.

E-mail address: gemaramirezrodriguez@hotmail.com

(G. Ramírez-Rodríguez).