



Editorial

Smoking in COPD Patients: A New Clinical Phenotype?☆

Tabaquismo en pacientes con EPOC, ¿un nuevo fenotipo clínico?

Francisco Carrión Valero,^{a,*} Susana Paulos Dos Santos,^a Bartolomé R. Celli^b^a Servicio de Neumología, Hospital Clínic Universitari, Departamento de Medicina, Universitat de València, Valencia, Spain^b Pulmonary and Critical Care Division, Brigham and Women's Hospital, Harvard Medical School, Boston, MA, United States

Chronic obstructive pulmonary disease (COPD) is a common, preventable, and treatable disease. The major cause of COPD is smoking, which is responsible for 85% of cases, and 20% of smokers can expect to develop COPD at some time during their life. Exposure to tobacco smoke before birth, in childhood and in adolescence can affect pulmonary growth.¹ In adults, passive smoking also contributes to the symptoms and development of COPD.

Although most smokers want to give up, very few receive dedicated medical help to quit. Recently, the European Commission Eurobarometer² documented that in Spain only 1 in 10 smokers who stopped smoking did so with the help of their doctor, a figure well below that of other European countries, such as the United Kingdom.

In 2016, 2.9 million people died worldwide as a result of COPD, and this disease was the sixth cause of death. The World Health Organization has predicted that in 2030 it will be responsible for 8.3% of all deaths, and will be the third cause of death.³

The prevalence of COPD in individuals between 40 and 80 years of age in Spain is 10.2%.⁴ The EPI-SCAN study showed that in Spain, 2 million individuals (25% women) have COPD. Up to 73% of patients are unaware that they have the disease, do not receive treatment for it, and often continue to smoke.⁵

Helping patients to stop smoking is the most important intervention in COPD. Until now, however, this has not been a priority among European pulmonologists,⁶ and as a result, more than a third of COPD patients continue their habit. Treatment of smoking is a very cost-effective intervention that helps reduce the risk of developing COPD and limits the progress of already established disease.^{6–8} In an international multicenter study, 40% of participants with moderate to severe COPD were smokers.⁹ Similarly, according to data from the AUDIPOC study, almost 25% of patients admitted for a COPD exacerbation are active smokers.¹⁰

The benefits of stopping smoking on FEV₁ decline and on COPD prognosis are well established in the literature,^{7,8} and similar

benefits have also been documented recently with respect to hospitalization rates. A Danish study that included more than 19 000 participants with 1260 hospitalizations in a follow-up period of 14.4 years showed that stopping smoking was associated with a reduction in the risk of hospital admission (HR: 0.57; 95% CI: 0.33–0.99).¹¹

Treatment of smoking is the most important and cost-effective intervention in COPD. Smoking cessation is the only effective preventive measure for delaying the progressive development of this disease,^{7,8} and has a decisive impact on the reduction of mortality. It is surprising then to note the high rate of smoking among COPD patients, and the lack of care routinely offered by specialists.

The recent update of the GOLD document (Global Strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease 2017 report) represents a considerable advance in the prevention and treatment of COPD. The most important contribution is the summary of therapeutic options for severity groups A–D. Yet, although the full document mentions the implementation of anti-smoking strategies in all scenarios, the topic occupies only 19 lines of the executive summary that was published in all the major respiratory medicine journals.¹²

However, if effective resources and time are devoted to smoking cessation, outcomes are encouraging.¹³ Nicotine replacement therapy, bupropion, varenicline, and in some patients nortriptyline, increase long-term abstinence rates, but must not be used as a single intervention without follow-up. Surprisingly, these medications have been well studied in numerous controlled clinical trials, but very few have been conducted in COPD patients. Smokers with COPD have specific characteristics associated with their smoking habit which confer them particular difficulties for quitting, but very few studies have attempted to identify the reasons for this phenomenon.

The Spanish COPD Guidelines (GesEPOC) were the first to propose phenotype-guided management, an approach that has been adopted by the national COPD guidelines of several European countries.¹⁴ GesEPOC describes the clinical phenotype as those features of the disease which, alone or in combination, define differences between individuals with COPD in terms of clinically significant parameters. The guidelines recommend identifying the phenotype and prescribing a targeted, specific treatment for each case. The 2017 edition of the GesEPOC recognizes 4 phenotypes:

☆ Please cite this article as: Carrión Valero F, Paulos Dos Santos S, Celli BR. Tabaquismo en pacientes con EPOC, ¿un nuevo fenotipo clínico? Arch Bronconeumol. 2018;54:249–250.

* Corresponding author.

E-mail address: carrión.fra@gva.es (F. Carrión Valero).

(1) non-exacerbator, (2) asthma-COPD overlap (ACO), (3) exacerbator with emphysema, and (4) exacerbator with chronic bronchitis.

In short, while smokers with COPD do not strictly meet the established criteria for a disease phenotype, as smoking is not a feature of the disease as such, evidence supports the notion that the identification of “COPD+smoking” in the initial classification may contribute to improving treatment, since it is useful for assigning patients to a subgroup with a certain prognosis that can help guide the physician towards the most appropriate therapy, which must include the systematic, routine treatment of smoking. Thus, in order to comply with the first of the 10 commandments of COPD,¹⁵ we urge that smoking is included in the initial classification.

References

1. Carrión Valero F, Jiménez Ruiz CA. El tabaquismo pasivo en la infancia. Arch Bronconeumol. 1999;35:39–47.
2. Special Eurobarometer 458. Attitudes of Europeans towards tobacco and electronic cigarettes. European Commission; 2017.
3. Murray CJL. Global, regional, and national age-sex specific mortality for 264 causes of death, 1980–2016: a systematic analysis for the Global Burden of Disease Study 2016 GBD 2016. Lancet. 2017;390:1151–210.
4. Miravittles M, Soriano JB, García Río F, Muñoz L, Durán Tauleria E, Sánchez G, et al. Prevalence of COPD in Spain: impact of undiagnosed COPD on quality of life and daily life activities. Thorax. 2009;64:863–8.
5. Soriano JB, Miravittles M, Borderías L, Durán-Tauleria E, García Río F, Martínez J, et al. Diferencias geográficas en la prevalencia de EPOC en España: relación con hábito tabáquico, tasas de mortalidad y otros determinantes. Arch Bronconeumol. 2010;46:522–30.
6. Jiménez Ruiz CA, Andreas S, Lewis KE, Tonnesen P, van Schayck, Hajek P, et al. Statement on smoking cessation in COPD and other pulmonary diseases and in smokers with comorbidities who find it difficult to quit. Eur Respir J. 2015;46:61–79.
7. Anthonisen NR, Connett JE, Kiley JP, Altose MD, Bailey WC, Buist S, et al. Effects of smoking intervention and the use of an inhalers anticholinergic bronchodilator on the rate of decline of FEV₁: the Lung Health Study. JAMA. 1994;272:2721497–505.
8. Kohansal R, Martínez-Cambor P, Agustí A, Buist S, Soriano JB. The natural history of chronic airflow obstruction revisited: an analysis of the Framingham offspring cohort. Am J Respir Crit Care Med. 2009;180:3–10.
9. Vogelmeier C, Hederer B, Glaab T, Schmidt H, Rutten-van Mölken MP, Beeh KM, et al., POET-COPD Investigators. Tiotropium versus salmeterol for the prevention of exacerbations of COPD. NEJM. 2011;364:1093–103.
10. Pozo Rodríguez F, López Campos JL, Álvarez Martínez C, Castro Acosta A, Agüero R, Hueto J, et al., AUDIPOC Study Group. Clinical Audit of COPD patients requiring hospital admissions in Spain. AUDIPOC study. PLoS ONE. 2012;7:e42156.
11. Godtfredsen NS, Vestbo J, Osler M, Prescott E. Risk of hospital admission for COPD following smoking cessation and reduction: a Danish population study. Thorax. 2002;57:967–72.
12. Vogelmeier CF, Criner GJ, Martínez FJ, Anzueto A, Barnes PJ, Borbeau J, et al. Informe 2017 de la Iniciativa Global para el Diagnóstico, Tratamiento y Prevención de la Enfermedad Pulmonar Obstructiva Crónica: Resumen Ejecutivo de GOLD. Arch Bronconeumol. 2017;53:128–49.
13. Van Eerd EA, van der Meer RM, van Schayck OC, Kotz D. Smoking cessation for people with chronic obstructive pulmonary disease. Cochrane Database Syst Rev. 2016. CD010744.
14. Guía de Práctica clínica para el diagnóstico y tratamiento de pacientes con EPOC. Guía Española de la EPOC (GesEPOC). Versión 2017. Arch Bronconeumol. 2017;53(Ext 1):4–64.
15. Marín JM, Cote C, Casanova C, Pinto-Plata V, de Oca MM, Divo MJ, et al. Simplificando las guías: los diez mandamientos de la EPOC. Arch Bronconeumol. 2016;52:179–80.