



Original article

Patterns of Physical Activity Progression in Patients With COPD

Maria Koreny^{a,b,c}, Heleen Demeyer^{d,e,f}, Marta Benet^{a,b,c}, Ane Arbilla-Etxarri^g,
Eva Balcells^{b,h,i}, Anael Barberan-García^j, Elena Gimeno-Santos^j, Nicholas S. Hopkinson^k,
Corina De Jong^{l,m}, Niklas Karlssonⁿ, Zafeiris Louvaris^{d,o}, Michael I. Polkey^k, Milo A. Puhan^p,
Roberto A. Rabinovich^q, Robert Rodríguez-Roisin^j, Pere Vall-Casas^r, Ioannis Vogiatzis^{o,s},
Thierry Troosters^{d,e}, Judith Garcia-Aymerich^{a,b,c,*}, The Urban Training Study Group and PROactive
Consortium members[◇]

^a ISGlobal, Barcelona, Spain^b Pompeu Fabra University (UPF), Barcelona, Spain^c CIBER Epidemiología y Salud Pública (CIBERESP), Barcelona, Spain^d Department of Rehabilitation Sciences, KU Leuven – University of Leuven, Leuven, Belgium^e Department of Respiratory Diseases, University Hospitals KU Leuven, Leuven, Belgium^f Department of Rehabilitation Sciences, Ghent University, Ghent, Belgium^g Physical Activity and Sports Sciences, Faculty of Psychology and Education, University of Deusto, Donostia-San Sebastián, Spain^h CIBER Enfermedades Respiratorias (CIBERES), Madrid, Spainⁱ Pneumology Department, Institut Hospital del Mar d'Investigacions Mèdiques (IMIM), Hospital del Mar, Barcelona, Spain^j University of Barcelona, Institut d'Investigació Biomèdica August Pi i Sunyer (IDIBAPS), CIBER Enfermedades Respiratorias (CIBERES), Hospital Clínic, Barcelona, Spain^k National Heart and Lung Institute, Imperial College, Royal Brompton Hospital Campus, London UK^l Department of General Practice, University Medical Center Groningen, University of Groningen, Groningen, The Netherlands^m Groningen Research Institute for Asthma and COPD (GRIAC), University Medical Center Groningen, University of Groningen, Groningen, The Netherlandsⁿ Patient Centered Science, BioPharmaceuticals, AstraZeneca R&D, Gothenburg, Sweden^o First Department of Respiratory Medicine, National and Kapodistrian University of Athens, Athens, Greece^p Epidemiology, Biostatistics and Prevention Institute, University of Zurich, Zurich, Switzerland^q ELEGY Colt Laboratory, Centre for Inflammation Research, The Queen's Medical Research Institute, University of Edinburgh, Edinburgh, UK^r Universitat Internacional de Catalunya (UIC), Barcelona, Spain^s Department of Sport, Exercise and Rehabilitation, Northumbria University Newcastle, Newcastle UK

ARTICLE INFO

Article history:

Received 17 June 2020

Accepted 6 August 2020

Keywords:

COPD

Physical activity

Patterns of progression

Cluster analysis

Determinants

ABSTRACT

Introduction: Although mean physical activity in COPD patients declines by 400–500 steps/day annually, it is unknown whether the natural progression is the same for all patients. We aimed to identify distinct physical activity progression patterns using a hypothesis-free approach and to assess their determinants.

Methods: We pooled data from two cohorts (usual care arm of Urban Training [NCT01897298] and PROactive initial validation [NCT01388218] studies) measuring physical activity at baseline and 12 months (Dynaport MoveMonitor). We identified clusters (patterns) of physical activity progression (based on levels and changes of steps/day) using k-means, and compared baseline sociodemographic, interpersonal, environmental, clinical and psychological characteristics across patterns.

Results: In 291 COPD patients (mean \pm SD 68 \pm 8 years, 81% male, FEV₁ 59 \pm 19%_{pred}) we identified three distinct physical activity progression patterns: *Inactive* (n=173 [59%], baseline: 4621 \pm 1757 steps/day, 12-month change (Δ): -487 ± 1201 steps/day), *Active Improvers* (n=49 [17%], baseline: 7727 \pm 3275 steps/day, Δ : $+3378 \pm 2203$ steps/day) and *Active Decliners* (n=69 [24%], baseline: 11 267 \pm 3009 steps/day, Δ : -2217 ± 2085 steps/day). After adjustment in a mixed multinomial logistic regression model using *Active Decliners* as reference pattern, a lower 6-min walking distance (RRR [95% CI] 0.94 [0.90–0.98] per 10 m, P=.001) and a higher mMRC dyspnea score (1.71 [1.12–2.60] per 1 point, P=.012) were independently related with being *Inactive*. No baseline variable was independently associated with being an *Active Improver*.

Abbreviations: BMI, body mass index; CAT, COPD assessment test; CCQ, clinical COPD questionnaire; CI, confidence interval; COPD, chronic obstructive pulmonary disease; C-PPAC, clinical visit—PROactive physical activity in COPD; FEV₁, forced expiratory volume in 1 second; FFMI, fat free mass index; FVC, forced vital capacity; GOLD, global initiative for chronic obstructive lung disease; HAD-A, hospital anxiety and depression scale – anxiety; HAD-D, hospital anxiety and depression scale – depression; LABA, long-acting beta₂-agonists; LAMA, long-acting anti-muscarinics; MET, metabolic equivalent of task; mMRC, modified medical research council dyspnea score; 6MWD, 6-min walking distance; MVPA, moderate to vigorous physical activity; RRR, relative risk ratio; SD, standard deviation.

* Corresponding author.

E-mail address: judith.garcia@isglobal.org (J. Garcia-Aymerich).◇ Please see a list of the members of the Urban Training Study Group and PROactive Consortium in [Appendices B and C](#).

Conclusions: The natural progression in physical activity over time in COPD patients is heterogeneous. While *Inactive* patients relate to worse scores for clinical COPD characteristics, *Active Improvers* and *Decliners* cannot be predicted at baseline.

© 2020 SEPAR. Published by Elsevier España, S.L.U. All rights reserved.

Palabras clave:

EPOC
Actividad física
Patrones de progresión
Análisis de conglomerados
Determinantes

Patrones de progresión de la actividad física en pacientes con EPOC

R E S U M E N

Introducción: Aunque la actividad física en pacientes con EPOC declina una media anual de 400-500 pasos/día, se desconoce si esta progresión es igual en todos los pacientes. Este estudio pretendió identificar los patrones de progresión de la actividad física mediante métodos libres de hipótesis y evaluar sus determinantes.

Métodos: Se estudiaron 291 pacientes con EPOC estable (media \pm DE: 68 \pm 8 años, 81% hombres, VEMS 59 \pm 19%_{pred}) de dos cohortes europeas con actividad física basal y a 12 meses (acelerómetro Dynaport MoveMonitor). Se identificaron conglomerados (patrones) de progresión de actividad física basados en los niveles y cambios de pasos/día usando *k-means*, y se compararon entre patrones las características sociodemográficas, interpersonales, ambientales, clínicas y psicosociales basales.

Resultados: Se identificaron tres patrones: *inactivo* (n = 173 [59%], basal: 4.621 \pm 1.757 pasos/día, cambio en 12 meses (Δ): -487 \pm 1.201 pasos/día), *activo que aumenta* (n = 49 [17%], basal: 7.727 \pm 3.275 pasos/día, Δ : +3.378 \pm 2.203 pasos/día) y *activo que reduce* (n = 69 [24%], basal: 11.267 \pm 3.009 pasos/día, Δ : -2.217 \pm 2.085 pasos/día). La distancia en la prueba de la marcha de 6 minutos (6MWD) y la disnea se asociaron independientemente con ser *inactivo*: RRR [IC 95%] 0,94 [0,90-0,98] por cada 10 m de 6MWD ($p = 0,001$) y 1,71 [1,12-2,60] por cada punto en la escala mMRC ($p = 0,012$), respectivamente, en comparación con el patrón *activo que reduce*. No se encontraron variables basales independientemente asociadas con ser *activo que aumenta*.

Conclusiones: La progresión natural de la actividad física en pacientes con EPOC es heterogénea. Mientras que el patrón de pacientes *inactivo* se relaciona con peores características clínicas de EPOC, no se pudo predecir la evolución de los *activos a aumentar o reducir*.

© 2020 SEPAR. Publicado por Elsevier España, S.L.U. Todos los derechos reservados.

Introduction

Physical activity is a key prognostic factor in chronic obstructive pulmonary disease (COPD), yet poorly understood. COPD patients are less active than healthy controls from the early stages of disease onwards¹⁻³ and this reduced activity has been associated with impaired prognosis and accelerated progression of COPD.^{4,5} For this reason, several national and international COPD guidelines recommend encouraging patients to maintain a good physical activity level.^{6,7}

Despite patients' and health professionals' efforts, physical activity has been shown to exhibit a spontaneous decline of an average of 400–500 steps/day per year in COPD patients.⁸⁻¹⁴ Such observed decline has been related, although not consistently, to lower lung function levels,^{10,11} the presence of exacerbations¹⁵⁻¹⁷ or the seasonality of testing (e.g., decline observed in patients going from summer to winter).^{12,18} Given the heterogeneous nature and progression of COPD,¹⁹ it can be hypothesized that physical activity progression also displays different patterns, not captured by the mean physical activity values. Two previous studies support this hypothesis by showing distinct physical activity trajectories over 9 months after a pulmonary rehabilitation program.^{20,21} However, the reported progression in physical activity after rehabilitation programs probably does not reflect how physical activity evolves in the wider COPD population nor in an observational setting, where patients receive a variable combination of pharmacological and non-pharmacological treatments.

We aimed (1) to identify, using a hypothesis-free approach, distinct patterns of natural physical activity progression in COPD patients recruited from diverse settings (primary care, hospital and rehabilitation services) and followed during 12 months; and (2) to assess the baseline sociodemographic, interpersonal, environmental, clinical and psychological determinants for the identified patterns. Better understanding of the natural

progression of physical activity, of potential distinct patterns and of their determinants could help to individualize strategies to increase (or prevent a decline in) physical activity.

Methods

Study Design and Patient Population

This was an observational (no intervention) cohort study of 12-month follow-up including patients from: (1) the usual care arm (n = 205) from the Urban Training study,²² that recruited patients from primary care and tertiary hospitals in five Catalan seaside municipalities (Badalona, Barcelona [center and shore areas], Mataró, Viladecans and Gavà); and (2) the clinically stable patients (n = 207) from the PROactive validation study,²³ that recruited patients from primary care settings, rehabilitation centers and tertiary hospitals in five European cities (Athens/Greece, Edinburgh and London/United Kingdom, Groningen/Netherlands, and Leuven/Belgium). Both studies defined COPD according to ATS/ERS (post-bronchodilator forced expiratory volume in 1 second (FEV₁) to forced vital capacity (FVC) ratio <0.70).²⁴ Patients were included in the present analyses if they had a valid physical activity measure (see below) at baseline and 12-month follow-up.

Both studies were approved by all local institutional review boards and written informed consent, including re-use of data for COPD-related research, was obtained from all patients.

Physical Activity Measurements

Physical activity was objectively measured using the Dynaport MoveMonitor (McRoberts BV, The Hague, The Netherlands)²⁵ for one week at baseline and follow-up. In Urban Training, patients wore the monitor for 24 h and data during waking hours (from

07:00 h to 22:00 h) were retrieved. In PROactive, patients wore the device during waking hours. A valid physical activity measurement was defined as a minimum of three days with at least 8 h of wearing time within waking hours for both studies²⁶; details have been previously published.^{22,23} A physical activity report was provided to patients if requested.

We used step count as the primary outcome to define physical activity progression patterns, and time spent in physical activity of moderate to vigorous intensity (MVPA, ≥ 3 METs [metabolic equivalents of tasks] min/day), movement intensity (m/s^2) during walking, and sedentary time (sum of lying and sitting time, hours/day) as secondary physical activity outcomes to describe patterns. Physical activity experience was assessed by the amount, difficulty and total scores of the Clinical visit-PROactive Physical Activity in COPD (C-PPAC) tool.²³

Other Measurements

We used variables available from both studies (i.e. exactly the same or equivalent standardized questions and procedures had been used) or variables that were available from one study only but had been related to physical activity or its evolution in the literature: (i) *sociodemographic*: age, sex, smoking history and education; (ii) *interpersonal*: marital status, working status, grandparenting and dog walking; (iii) *environmental*: season of recruitment, average yearly rainfall and urban vulnerability index (a measure of socioeconomic status at the census tract level); (iv) *clinical*: post-bronchodilator FEV₁ and FVC, the 6-min walking distance (6MWD) test, the COPD Assessment test (CAT), the Clinical COPD Questionnaire (CCQ), the modified Medical Research Council dyspnea scale (mMRC), the number of acute COPD exacerbations requiring a hospital admission in the previous 12 months and during follow-up, body mass index (BMI) and fat free mass index (FFMI) by physical examination and bioelectrical impedance, comorbidities from medical records, pharmacological treatment for COPD, pulmonary rehabilitation at baseline and follow-up, incident diseases during follow-up, and knowledge of baseline physical activity (i.e. report on request); and (v) *psychological*: the Hospital Anxiety (HAD-A) and Depression (HAD-D) scores. Full details on study procedures and quality control have been reported previously.^{22,23,27}

Statistical Analysis

Sample size calculations, missing data strategy and full statistical analyses are provided in the supplement.

We identified cluster groups (physical activity patterns) using k-means,²⁸ a hypothesis-free method that allowed grouping patients based on the baseline level, the final level and the change in daily step count. To characterize the patterns, we described

physical activity and physical activity experience variables according to the cluster groups and compared baseline to follow-up values by paired *t*-tests.

To assess determinants of physical activity progression patterns, we first compared subjects' characteristics by physical activity patterns and obtained *P*-values from mixed logistic regression models with random intercepts for study and city area to account for possible heterogeneity in unmeasured characteristics related to study and city area. Then we built a multivariable multinomial regression model using the generalized linear latent and mixed model, with also random intercepts for study and city.²⁹ Model building combined step-forward and backward algorithms and we tested goodness of fit of the final model.

As sensitivity analyses, we (1) repeated cluster analysis separately for Urban Training and PROactive; (2) tested the association between the change in daily step count and the change in wearing time overall and per pattern; and (3) repeated the clustering after excluding patients included in pulmonary rehabilitation programs at baseline and/or during follow-up.

All analyses were conducted using Stata/SE 14.2 (StataCorp, College Station, TX, USA).

Results

From 412 patients at baseline, 291 (71%) completed the follow-up visit and were included in the current analyses (Fig. S1). These patients had a higher proportion of males, better functional status and were more active at baseline than those lost-to follow-up (Table S1). Included patients were 81% male and had a mean age of 68 years, FEV₁ of 59% predicted, 6MWD of 477 m, mMRC dyspnea score of 1.3, and walked 6720 steps/day (Table 1). Compliance with the activity monitor during waking hours was excellent: at baseline median (range) valid days of 7 (3–7) and mean \pm SD wearing hours of 14.6 ± 0.5 in Urban Training, and 6 (3–7) days and 14.8 ± 2.2 wearing hours in PROactive; and at follow-up, 7 (4–7) days and 14.6 ± 0.6 wearing hours in Urban Training, and 6 (3–7) days and 14.4 ± 2.2 wearing hours in PROactive.

At the group level, the step count did not change over 12 months. In the hypothesis-free approach, we identified three cluster groups (three distinct physical activity patterns) (Fig. 1, Table S2). A first cluster ($n = 173$ [59%]), labeled *Inactive* pattern due to the low step count, walked at baseline mean \pm SD 4621 ± 1757 steps/day and decreased their physical activity by 487 ± 1201 steps/day over 12 months. A second cluster ($n = 49$ [17%]), labeled *Active Improvers*, walked 7727 ± 3275 steps/day at baseline and increased by 3378 ± 2203 steps/day. The third cluster ($n = 69$ [24%]), labeled *Active Decliners*, walked $11,267 \pm 3009$ steps/day at baseline and decreased by 2217 ± 2085 steps/day. Distribution of

Table 1
Patient Characteristics at Baseline and at 12-month Follow-up for All Patients ($n = 291$) and by Study Group (Urban Training and PROactive study).

	All Patients $n = 291$ (100%)	Urban Training Study $n = 148^a$ (51%)	PROactive Study $n = 143^a$ (49%)
Sociodemographic			
Age (years)	68 ± 8	69 ± 8	67 ± 8
Sex (men)	237 (81)	130 (88)	107 (75)
Current smoker	52 (18)	30 (20)	22 (15)
Pack-years	58 ± 41	60 ± 45	56 ± 37
Education, high school or higher	168 (58)	49 (33)	119 (83)
Interpersonal			
Living with a partner ^b	216 (74)	124 (84)	92 (65)
Active worker ^c	36 (12)	16 (11)	20 (14)
Grandparenting ^d	67 (45)	67 (45)	–
Dog walking ^d	20 (14)	20 (14)	–

Table 1 (Continued)

	All Patients n = 291 (100%)	Urban Training Study n = 148 ^a (51%)	PROactive Study n = 143 ^a (49%)
Environmental			
Recruitment season			
Spring	35 (12)	35 (24)	0 (0)
Summer	58 (20)	15 (10)	43 (30)
Fall	154 (53)	54 (36)	100 (70)
Winter	44 (15)	44 (30)	0 (0)
Average rainfall (h/day) ^{e,f}	0.62 (0.30–1.13)	–	0.62 (0.30–1.13)
Urban vulnerability index (from 0 -lowest to 1 - highest) ^{d,g}	0.637 ± 0.175	0.637 ± 0.175	–
Clinical			
FEV ₁ (% predicted)	58.6 ± 19.3	58.2 ± 17.6	59.0 ± 21.0
FEV ₁ /FVC ratio	0.51 ± 0.13	0.55 ± 0.12	0.48 ± 0.13
Airflow limitation severity (post-bronchodilator FEV ₁)			
GOLD 1: Mild (FEV ₁ ≥ 80% predicted)	39 (13)	15 (10)	24 (17)
GOLD 2: Moderate (50% ≤ FEV ₁ < 80% predicted)	147 (51)	80 (54)	67 (47)
GOLD 3: Severe (30% ≤ FEV ₁ < 50% predicted)	88 (30)	45 (30)	43 (30)
GOLD 4: Very severe (FEV ₁ < 30% predicted)	17 (6)	8 (6)	9 (6)
6MWD (meters)	477 ± 103	501 ± 83	452 ± 116
CAT score (0–40)	12.9 ± 7.6	12.2 ± 7.6	13.6 ± 7.5
CCQ score (0–6)	1.55 ± 0.98	1.40 ± 0.95	1.70 ± 0.98
C-PPAC amount score (0–100)	69.0 ± 15.8	74.7 ± 14.9	63.8 ± 14.9
C-PPAC difficulty score (0–100)	78.4 ± 14.5	82.7 ± 13.4	74.5 ± 14.5
C-PPAC total score (0–100)	73.7 ± 12.8	78.7 ± 11.5	69.2 ± 12.3
mMRC score (0–4)	1.3 ± 0.9	1.1 ± 0.8	1.5 ± 1.0
Any COPD exacerbation with hospital admission in previous 12 months	34 (12)	12 (8)	22 (15)
BMI (kg/m ²)	27.6 ± 4.6	28.3 ± 4.6	26.8 ± 4.6
FFMI (kg/m ²)	19.0 ± 3.0	19.6 ± 3.2	18.4 ± 2.8
Cardiovascular disease ^h	176 (60)	90 (61)	86 (60)
Ischemic heart disease ^h	29 (10)	13 (9)	16 (11)
Diabetes mellitus ^h	51 (18)	38 (26)	13 (9)
LABA or LAMA, alone	41 (14)	23 (16)	18 (13)
Inhaled corticosteroid with LABA and/or LAMA	179 (62)	80 (54)	99 (71)
Pulmonary rehabilitation at baseline	15 (5)	6 (4)	9 (6)
Knowledge of baseline PA	19 (7)	19 (13)	0 (0)
Psychological			
Anxiety (HAD-A, 0–21)	5 ± 4	5 ± 4	5 ± 4
Depression (HAD-D, 0–21)	4 ± 3	3 ± 3	5 ± 3
Physical activity			
Step count (steps/day)	6720 ± 3667	7783 ± 3847	5619 ± 3121
Time in moderate-to-vigorous physical activity (≥3 METs; min/day)	99.4 ± 45.3	109.1 ± 45.7	89.4 ± 42.8
Intensity during walking (m/s ²)	1.86 ± 0.31	1.88 ± 0.32	1.84 ± 0.29
Sedentary time (h/day)	10.53 ± 1.94	10.43 ± 1.48	10.64 ± 2.31
Wearing time (h/day)	14.73 ± 1.56	14.64 ± 0.54	14.81 ± 2.16
Follow-up data			
Any COPD exacerbation with hospital admission during follow-up	28 (10)	10 (7)	18 (13)
Any incident comorbidity during follow-up ^{d,i}	34 (23)	34 (23)	–
Pulmonary rehabilitation during follow-up	16 (6)	6 (4)	10 (7)
Wearing time at follow-up (h/day)	14.52 ± 1.63	14.60 ± 0.61	14.43 ± 2.24

Notes: Data are presented as n (%), mean ± SD or median (interquartile range).

Abbreviations: FEV₁: forced expiratory volume in 1 s; FVC: forced vital capacity; GOLD: Global Initiative for Chronic Obstructive Lung Disease; 6MWD: 6-min walking distance; CAT: COPD Assessment Test; CCQ: Clinical COPD Questionnaire; C-PPAC: Clinical visit–PROactive Physical Activity in COPD (higher numbers indicate a better score); mMRC: modified Medical Research Council; BMI: body mass index; FFMI: fat free mass index; LABA: long-acting beta₂-agonists; LAMA: long-acting anti-muscarinics; HAD-A: Hospital Anxiety and Depression scale – Anxiety; HAD-D: Hospital Anxiety and Depression scale – Depression; MET: metabolic equivalent of task.

^a Some variables have missing values, as follows. Urban Training: 1 in education, 25 in C-PPAC scores, 1 in any COPD exacerbation with hospital admission in previous 12 months, 18 in FFMI, 2 in HAD anxiety and depression, 5 in any COPD exacerbation with hospital admission during follow-up, 2 in pulmonary rehabilitation during follow-up; PROactive: 1 in living with a partner, 21 in average rainfall, 1 in CAT score, 1 in CCQ score, 6 in C-PPAC scores, 8 in FFMI, 3 in LABA or LAMA, alone, 3 in inhaled corticosteroid with LABA and/or LAMA, 1 in HAD anxiety and depression, 3 in pulmonary rehabilitation during follow-up.

^b Marital status: living with a partner vs single, widowed or divorced.

^c Working status: active worker (working full-time or part-time) vs. unemployed, housework or retired.

^d Only available for Urban Training.

^e Only available for PROactive.

^f Average rainfall was calculated as the mean of the measurements at baseline, 6 and 12 months.

^g The urban vulnerability index is a measure of socioeconomic status at the census tract level that combines demographic, economic, residential and subjective indicators, and ranges from lowest [0] to highest [1] level of neighborhood vulnerability.

^h ICD10 codes: I00–I99 for cardiovascular diseases; I20–I25 for ischemic heart disease, E14 for diabetes mellitus.

ⁱ Incident comorbidities included ICD10 codes C00–N99.

MVPA and walking intensity by physical activity pattern followed the same sequence as steps/day, except for walking intensity in *Active Improvers* that did not change. Sedentary time did not change for *Inactive*, decreased for *Active Improvers* and increased

moderately for *Active Decliners*. The physical activity experience as expressed by C-PPAC scores did not change for the *Inactive* pattern; the *Active Improvers* increased the C-PPAC scores (i.e., increased amount and reduced difficulty); the *Active Decliners* decreased the




Fig. 1. Physical activity variables at baseline and at 12-month follow-up, overall and by PA progression pattern (*Inactive*, *Active Improvers* and *Active Decliners*). Notes: Data are presented as mean \pm SE (specific numbers are presented in Table S2). * P -value ≤ 0.05 . Abbreviations: MVPA: moderate-to-vigorous physical activity; MET: metabolic equivalent of task.

C-PPAC amount and total scores while the C-PPAC difficulty score did not change (Fig. 2, Table S2).

Patients in the *Inactive* physical activity pattern had a higher degree of education, a smaller proportion was living with a partner or grandparenting, and they presented with a worse general health status, lower lung function, poorer exercise capacity, worse quality of life and higher dyspnea and depression scores than those in the *Active Improvers* or *Decliners* patterns (Table 2). *Active Improvers* and

Decliners were very similar in their baseline characteristics, except for their daily step count.

In the multivariable multinomial logistic regression model we used *Active Decliners* as the reference pattern to capture both the determinants of being *Inactive* vs *Active* and the determinants of being an *Active Improver* vs *Decliner*. A lower exercise capacity and a higher mMRC dyspnea score were independently related with being *Inactive* whereas no variable was identified as

Table 2
Patient Characteristics by Physical Activity Progression Pattern (*Inactive, Active Improvers and Active Decliners*) in 291 COPD Patients.

	<i>Inactive</i>	<i>Active Improvers</i>	<i>Active Decliners</i>	<i>P-Value for Inactive vs Active Improvers and Decliners^b</i>	<i>P-Value for Active Improvers vs Decliners^b</i>
	<i>n</i> = 173 ^a (59%)	<i>n</i> = 49 ^a (17%)	<i>n</i> = 69 ^a (24%)		
<i>Urban training study</i>	59 (34)	39 (80)	50 (72)		
<i>PROactive study</i>	114 (66)	10 (20)	19 (28)		
<i>Sociodemographic</i>					
Age (years)	68 ± 8	69 ± 9	67 ± 7	.282	.079
Sex (men)	137 (79)	41 (84)	59 (86)	.931	.789
Current smoker	33 (19)	5 (10)	14 (20)	.152	.152
Pack-years	60 ± 38	63 ± 52	49 ± 40	.187	.082
Education, high school or higher	124 (72)	20 (42)	24 (35)	.006	.452
<i>Interpersonal</i>					
Living with a partner ^c	115 (66)	42 (88)	59 (86)	.017	.714
Active worker ^d	18 (10)	6 (12)	12 (17)	.088	.454
Grandparenting ^e	20 (34)	21 (54)	26 (52)	.039	.863
<i>Environmental</i>					
Average rainfall (h/day) ^{f,g}	0.63 (0.33–1.13)	0.90 (0.57–1.47)	0.33 (0.23–1.00)	.877	.329
Urban vulnerability index (from 0 – lowest to 1 – highest) ^{e,h}	0.646 ± 0.176	0.613 ± 0.200	0.646 ± 0.153	.312	.369
<i>Clinical</i>					
FEV ₁ (% predicted)	55.9 ± 19.8	62.9 ± 15.8	62.4 ± 19.5	.001	.875
FEV ₁ /FVC ratio	0.48 ± 0.14	0.55 ± 0.11	0.55 ± 0.11	.004	.904
6MWD (meters)	446 ± 105	521 ± 90	524 ± 78	<.001	.861
CAT score (0–40)	14.2 ± 7.7	11.5 ± 7.3	10.5 ± 6.6	.002	.435
CCQ score (0–6)	1.74 ± 0.97	1.23 ± 0.91	1.29 ± 0.93	.001	.780
C-PPAC difficulty score (0–100) ⁱ	74.9 ± 14.7	82.6 ± 13.6	84.8 ± 11.6	<.001	.380
mMRC score (0–4)	1.5 ± 1.0	1.0 ± 0.8	0.9 ± 0.7	<.001	.329
Any COPD exacerbation with hospital admission in previous 12 months	24 (14)	4 (8)	6 (9)	.517	.918
BMI (kg/m ²)	27.6 ± 5.0	27.5 ± 3.9	27.5 ± 4.2	.139	.999
FFMI (kg/m ²)	18.9 ± 3.0	19.3 ± 2.9	19.0 ± 3.1	.650	.591
Cardiovascular disease ^j	109 (63)	28 (57)	39 (57)	.221	.930
Ischemic heart disease ^l	18 (10)	5 (10)	6 (9)	.898	.807
Diabetes mellitus ^j	23 (13)	11 (22)	17 (25)	.412	.786
LABA or LAMA, alone	24 (14)	7 (14)	10 (15)	.796	.949
Inhaled corticosteroid with LABA and/or LAMA	115 (67)	28 (57)	36 (53)	.311	.658
<i>Psychological</i>					
Anxiety (HAD-A, 0–21)	5 ± 4	5 ± 3	5 ± 4	.755	.774
Depression (HAD-D, 0–21)	5 ± 3	3 ± 3	3 ± 3	.009	.992
<i>Follow-up data</i>					
Any COPD exacerbation with hospital admission during follow-up	19 (11)	4 (8)	5 (7)	.759	.846
Any incident comorbidity during follow-up ^{e,k}	10 (17)	10 (26)	14 (28)	.191	.804

Notes: Data are presented as *n* (%), mean ± SD or median (interquartile range).

Abbreviations: FEV₁: forced expiratory volume in 1 second; FVC: forced vital capacity; 6MWD: 6-min walking distance; CAT: COPD Assessment Test; CCQ: Clinical COPD Questionnaire; C-PPAC: Clinical visit–PROactive Physical Activity in COPD (higher numbers indicate a better score); mMRC: modified Medical Research Council; BMI: body mass index; FFMI: fat free mass index; LABA: long-acting beta₂-agonists; LAMA: long-acting anti-muscarinics; HAD-A: Hospital Anxiety and Depression scale - Anxiety; HAD-D: Hospital Anxiety and Depression scale - Depression.

^a Some variables have missing values, as follows. *Inactive*: 15 in average rainfall, 1 in CAT total, 1 in CCQ score, 14 in C-PPAC difficulty score, 17 in FFMI, 2 in LABA or LAMA, alone, 2 in inhaled corticosteroid with LABA and/or LAMA, 1 in HAD anxiety and depression, 3 in any COPD exacerbation with hospital admission during follow-up; *Active Improvers*: 1 in education, 1 in living with a partner, 2 in average rainfall, 5 in C-PPAC difficulty score, 4 in FFMI, 1 in HAD anxiety and depression, 1 in any COPD exacerbation with hospital admission during follow-up; *Active Decliners*: 4 in average rainfall, 12 in C-PPAC difficulty score, 1 in any COPD exacerbation with hospital admission in previous 12 months, 5 in FFMI, 1 in LABA or LAMA, alone, 1 in inhaled corticosteroid with LABA and/or LAMA, 1 in HAD anxiety and depression, 1 in any COPD exacerbation with hospital admission during follow-up.

^b *P*-Value from mixed logistic regression models with random effects for study (UT and PROactive) and city area (Badalona, Barcelona-center, Barcelona-shore, Mataró, Viladecans/Gavà, Athens, Edinburgh, Groningen, Leuven, London).

^c Marital status: living with a partner vs single, widowed or divorced.

^d Working status: active worker (working full-time or part-time) vs. unemployed, housework or retired.

^e Only available for Urban Training.

^f Only available for PROactive.

^g Average rainfall was calculated as the mean of the measurements at baseline, 6 and 12 months.

^h The urban vulnerability index is a measure of socioeconomic status at the census tract level that combines demographic, economic, residential and subjective indicators, and ranges from lowest [0] to highest [1] level of neighborhood vulnerability.

ⁱ Only C-PPAC difficulty is provided as C-PPAC amount and total score include steps/day which were used for the generation of the PA patterns and therefore cannot be assessed as predictors.

^j ICD10 codes: I00 to I99 for cardiovascular diseases; I20–I25 for ischemic heart disease, E14 for diabetes mellitus.

^k Incident comorbidities included ICD10 codes C00–N99.




Fig. 2. Physical activity experience variables at baseline and at 12-month follow-up, overall and by PA progression pattern (*Inactive*, *Active Improvers* and *Active Decliners*). Notes: Data are presented as mean \pm SE (specific numbers are presented in Table S2). * P -value $\leq .05$. Abbreviations: C-PPAC: Clinical visit–PROactive Physical Activity in COPD (higher numbers indicate a better score). C-PPAC variables have 87 missing values: 38 in *Inactive*, 21 in *Active Improvers*, and 28 in *Active Decliners*.

Table 3
Adjusted Predictive Factors for *Inactive* and *Active Improvers* vs *Active Decliners* in 291 COPD Patients.

	<i>Active Decliners</i>	<i>Inactive</i>		<i>Active Improvers</i>	
	RRR (95% CI)	RRR (95% CI)	P -Value ^a	RRR (95% CI)	P -Value ^a
6MWD (per 10 m)	1.00 (ref)	0.94 (0.90–0.98)	.001	1.00 (0.96–1.05)	.868
mMRC score (per 1 point)	1.00 (ref)	1.71 (1.12–2.60)	.012	1.23 (0.73–2.07)	.437

Abbreviations: RRR: relative risk ratio; CI: confidence interval; 6MWD: 6-min walking distance; mMRC: modified Medical Research Council.

^a P -Value from multinomial regression model with random effects for study (UT and PROactive) and city area (Badalona, Barcelona-center, Barcelona-shore, Mataró, Viladecans/Gavà, Athens, Edinburgh, Groningen, Leuven, London).

independently associated with being an *Active Improver* (Table 3). The final model showed good fit. Sensitivity analyses confirmed the results (Tables S3–S5).

Discussion

This study identified, for the first time to our knowledge, the natural progression of physical activity in COPD patients. We used a hypothesis-free approach that allowed the identification of patterns without a priori assumptions about the physical activity

changes over time. We found that (1) the natural change in physical activity over time was indeed heterogeneous; (2) the majority of patients (59%) was inactive at baseline and decreased their physical activity level subsequently; (3) among active individuals some increased and some decreased their physical activity level; and (4) although clinical COPD characteristics were related to the physical activity level at baseline they could not predict subsequent physical activity changes.

A first important finding is that physical activity progression in COPD is heterogeneous. In our 12-month study, mean changes in

the full group were virtually zero; however when using hypothesis-free clustering methods, we identified one *Inactive* pattern which decreased and two *Active* patterns which increased or decreased physical activity. This observation is in line with previous reports of heterogeneous physical activity progression in patients with rheumatoid arthritis.³⁰ The average lack of 12-month change in step count differs from previous studies that showed overall a decrease in physical activity.^{8,10,11,13} A potential explanation is that most of these studies recruited patients from outpatient or pulmonary clinics, who may have slightly more advanced disease and reduced variability in physical activity and COPD characteristics as compared to our sample including also primary care. Supporting this, the group of patients who started with a lower physical activity (59%) was similar in their baseline characteristics to previous studies and also had a comparable mean decrease of around 500 steps/day.^{10,11} Notably, the low overall dyspnea score may have positively influenced the physical activity level of our study population. A second explanation could lie in the high proportion of male subjects and regional differences in physical activity practice (a cohort of patients included in the Mediterranean region⁵ had a baseline physical activity comparable to the Urban Training sample). These characteristics of our sample could justify the two patterns with relatively high baseline physical activity and an average small physical activity change.

The second important finding is that there seems to exist a group of COPD patients (our *Active Improvers*) that spontaneously increase their physical activity over time. Of note, such observed increase of >3000 steps/day is remarkably high given that the minimal important difference has been proposed between 600 and 1100 steps/day.³¹ There are several possibilities that would explain this observed increase. First, some patients could have been inactive at baseline by chance; however, we tested this option against study records by screening for atypical events and it did not hold true. Second, regression to the mean could account for part of the increase, but in our data regression to the mean was estimated to account for maximal 25% of the effect. Third, changes in daily steps could be due to changes in wearing time, but this was not the case in our study (Table S4). Fourth, patients could have increased their physical activity after participation in rehabilitation programs, but this was dismissed in our analysis (Table S5). Finally, we considered that some patients in the usual care arm of Urban Training could have increased their physical activity due to being enrolled in a physical activity study. However, the proportion of patients from Urban Training was similar between *Active Improvers* and *Active Decliners*. Thus, we suggest that some patients do actually increase their physical activity.

The evolution of other physical activity variables provided complementary information. Time in MVPA and sedentary time (opposite direction) paralleled the progression of step count in all three patterns, supporting previous research that suggested that in COPD patients, physical activity and sedentary time provide information about the same concept.⁵ We also investigated the progression of physical activity from the perspective of patients. As expected, C-PPAC amount and total scores followed a trend similar to the objectively measured physical activity, as they include steps/day in their calculation. However, C-PPAC difficulty score remained unchanged in *Inactive* and *Active Decliners* and increased (i.e., less difficulty) in *Active Improvers*, suggesting that the observed increase in amount could be related to experiencing fewer difficulties (less dyspnea for instance³²) while being active.

Our third main finding is the impossibility to predict the physical activity progression patterns, despite having included sociodemographic, interpersonal, environmental and psychological characteristics in addition to the typical clinical COPD variables. We found a large set of COPD-related, functional characteristics associated with the *Inactive* pattern, in accordance with previous,

mainly cross-sectional, literature about the determinants of physical activity levels in COPD.^{2,4} Also higher education levels, lower social support (living alone, not taking care of grandchildren) and higher depression scores related to being in the *Inactive* pattern, although none of these factors remained in the multivariable model suggesting they were subject to confounding.

Most tellingly, we did not identify any factors that could predict among *Active* patients, the evolution to *Improvers* or *Decliners*. Surprisingly, the presence of severe exacerbations during follow-up did not play any role. It could be speculated that our harmonized exacerbation data was not detailed enough to distinguish the severity of exacerbations, the length of hospital stay or the time from the last exacerbation to physical activity assessment at follow-up. We also considered the role of incident comorbidities during follow-up, which could have influenced behavior, but they were not significantly different for the three patterns. Pharmacological treatment for COPD was not different across progression patterns discarding any potential role for treatment inappropriateness. Moreover, we did not find an association between the recruitment season and physical activity progression. This is in line with the hypothesis that the recruitment season, although possibly affecting the baseline levels of physical activity,^{12,18,33} would not affect the progression pattern during a follow-up of 12 months. Finally, we did not find an effect of accumulated rainfall on physical activity progression, as recently described cross-sectionally in the same PROactive population.¹⁴

Our study has several implications. It adds to the current knowledge that contrary to the general belief not all patients decline but some patients considerably improve their physical activity, which should be confirmed in future research and shows the importance of including a usual care group in intervention studies. The limitation of traditional clinical COPD characteristics to predict physical activity progression suggests that further research should broaden the view and give more attention to interpersonal and environmental factors potentially related to the individual's motivation. As the optimal timing and use of physical activity interventions to improve physical activity (especially in the long term) is still unclear,³⁴ understanding the different COPD progression patterns may help to overcome a one-size-fits-all approach and customize physical activity promotion to reflect different physical activity practices and different treatment needs.³⁵ Finally, our results highlight the limitation of using mean population values in phenomena that are heterogeneous in nature.

A major strength of our study is the inclusion of patients across a broad spectrum of disease severities and physical activity in several European cities. This makes our results applicable (i.e., more representative) to a larger COPD population than a single recruitment setting or severity group. In addition, the inclusion of patients from diverse geographic locations allowed us to indirectly control for residual confounding. Moreover, we included some variables beyond the conventional clinical COPD characteristics.³⁶ The use of the hypothesis-free clustering approach allowed us to identify patterns of physical activity progression based on the distribution of the data without prior assumptions.

However, we acknowledge some shortcomings. We had a small sample size for some of the hypothesized determinants of physical activity progression patterns, such as dog walking, current pulmonary rehabilitation and knowledge of baseline physical activity, which precluded our ability to test their role. Similarly, we did not collect information on some physical activity barriers (e.g. costs or transportation difficulties), which precluded testing their role on physical activity progression. The drop-out was 29% which is comparable to previous studies^{37,38} but the excluded patients had worse functional parameters, and we cannot rule out that they would have presented with a fourth, potentially declining pattern. The two measurement points available for both studies allowed

to investigate only linear patterns over time. Having more data points could provide more detailed information on the trajectories. In addition, a longer follow-up would have been desirable, but the 12-month span appears reasonably long to provide this first novel insight into physical activity patterns. Finally, one might argue that pooling of the two studies was not appropriate, although our sensitivity analyses showed similar cluster results and characteristics and it resulted in a broad spectrum of physical activity and COPD severity.

In conclusion, the natural change in physical activity over time in COPD patients is heterogeneous and three distinct patterns of physical activity progression have been identified: a predominant *Inactive* pattern, related to worse scores for clinical COPD characteristics, and two *Active* patterns, *Improvers* and *Decliners*, which cannot be predicted at baseline.

Authors' Contribution

JGA had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis. MK, HD and JGA drafted the first version of the manuscript; MK, HD, MB and JGA had full access to the data and were responsible for the statistical analysis; MK, HD, MB, AAE, EB, ABG, EGS, NH, CdJ, NK, ZL, MPo, MPu, RRa, RRo, PVC, IV, TT, JGA contributed to data collection and coordination. All authors (1) provided substantial contributions to the conception or design of the work; or the acquisition, analysis or interpretation of data for the work; (2) drafted or revised the manuscript for important intellectual content; (3) approved the final version; and (4) agreed to be accountable for all aspects of the work.

Ethics Approval

This study was conducted in accordance with the amended Declaration of Helsinki. Both studies were approved by all local institutional review boards and written informed consent, including re-use of data for COPD-related research, was obtained from all patients.

The Urban Training trial was approved by the ethics committees of all participating institutions (Comitè Ètic d'Investigació Clínica Parc de Salut MAR 2011/4291/I, Comitè Ètic d'Investigació Clínica de l'IDIAP Jordi Gol i Gurina P11/116, Comitè Ètic d'Investigació Clínica de l'Hospital Universitari de Bellvitge PR197/11, Comitè Ètic d'Investigació Clínica de l'Hospital Universitari Germans Trias i Pujol AC-12-004, Comitè Ètic d'Investigació Clínica de l'Hospital Clínic de Barcelona 2011/7061, Comitè Ètic d'Investigació Clínica de l'Hospital de Mataró November 23rd, 2011).

The PROactive study was advised and approved by the PROactive ethics and patient advisory boards, and approved by the local ethics committee at each center (Commissie medische ethiek van de universitaire ziekenhuizen KU Leuven (Leuven, S-55919); Medische ethische toetsingscommissie universitair medisch centrum Groningen (Groningen, Metc 2013.362); RES Committee London–South East (London and Edinburgh, 13/LO/1660); Scientific Council of the 'Sotiria' General Hospital for Chest Diseases (Athens, 27852/7-10-13); Kantonale Ethikkommission Zürich and Ethikkommission Nordwest- und Zentralschweiz (Zurich, KEK-ZH-Nr. 2013-0469).

Prior abstract presentation/publication: This work was presented at the European Respiratory Society Congress 2019 in Madrid and the abstract was published in the European Respiratory Journal 2019; 54: Suppl. 63, OA5361.

Funding Information

The Urban Training study was funded by grants from Fondo de Investigación Sanitaria, Instituto de Salud Carlos III (ISCIII, PI11/01283 and PI14/0419), integrated into Plan Estatal I+D+I 2013–2016 and co-funded by ISCIII-Subdirección General de Evaluación y Fomento de la Investigación and Fondo Europeo de Desarrollo Regional (FEDER); Sociedad Española de Neumología y Cirugía Torácica (SEPAR, 147/2011 and 201/2011), Societat Catalana de Pneumologia (Ajuts al millor projecte en fisioteràpia respiratòria 2013). We acknowledge support from the Spanish Ministry of Science and Innovation through the "Centro de Excelencia Severo Ochoa 2019–2023" Program (CEX2018-000806-S), and support from the Generalitat de Catalunya through the CERCA Program.

The PROactive project was funded by the European Commission Innovative Medicines Initiative Joint Undertaking (IMI JU # 115011). HD is a post-doctoral research fellow of FWO Vlaanderen. ZL is a post-doctoral fellow of the FWO-Flanders (#12U5618N).

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Conflict of Interest

NK is employed by AstraZeneca. MP reports personal fees from Philips, during the conduct of the study. TT reports lecture fees to his institution from Boehringer Ingelheim, Chiesi Belgium and AstraZeneca outside the submitted work. JGA reports payments for consulting and lecture fees to her institution from AstraZeneca and lecture fees from Esteve, Chiesi and Menarini outside the submitted work. The authors report no other conflict of interest in this work.

Acknowledgments

The authors thank Anne-Elie Carsin for the statistical support.

Appendix A. Supplementary Data

Supplementary data associated with this article can be found, in the online version, at [doi:10.1016/j.arbres.2020.08.001](https://doi.org/10.1016/j.arbres.2020.08.001).

Appendix B. The Urban Training Study Group:

ISGlobal, Barcelona: Ane Arbillaga-Etxarri, Marta Benet, Anna Delgado, Judith Garcia-Aymerich, Elena Gimeno-Santos, Jaume Torrent-Pallicer; FCS Blanquerna, Universitat Ramon Llull, Barcelona: Jordi Vilaró; Servei de Pneumologia, Hospital Clínic de Barcelona, Barcelona: Anael Barberan-Garcia, Robert Rodriguez-Roisin; Hospital del Mar, Institut Hospital del Mar d'Investigacions Mèdiques (IMIM), Barcelona: Eva Balcells, Diego A Rodríguez Chiaradía; Hospital Universitari Germans Trias i Pujol, Badalona: Alicia Marín; Hospital de Mataró, Consorci Sanitari del Maresme, Mataró: Pilar Ortega; Hospital de Viladecans, Viladecans: Nuria Celorrio; Institut Universitari d'Investigació en Atenció Primària Jordi Gol (IDIAP Jordi Gol): Mónica Monteagudo, Nuria Montellà, Laura Muñoz, Pere Toran; Center d'Atenció Primària Viladecans 2, Institut Català de la Salut, Viladecans: Pere Simonet; Center d'Atenció Primària Passeig de Sant Joan, Institut Català de la Salut, Barcelona: Carme Jané, Carlos Martín-Cantera; Center d'Atenció Primària Sant Roc, Institut Català de la Salut, Badalona: Eulàlia Borrell; Universitat Internacional de Catalunya (UIC), Barcelona: Pere Vall-Casas.

Appendix C. The PROactive Consortium Members:

Almirall, Barcelona, Spain: Nathalie Ivanoff; AstraZeneca AB, Mölndal, Sweden: Niklas Karlsson and Solange Corriol-Rohou; British Lung Foundation, London, UK: Ian Jarrod; Boehringer Ingelheim, Nieder-Ingelheim, Germany: Damijen Erzen; Chiesi Farmaceutici S.A. Parma, Italy: Caterina Brindicci, Tim Higenbottam and Mario Scuri; Choice Healthcare Solutions, Hitchin, UK: Paul McBride; European Respiratory Society, Lausanne, Switzerland: Nadia Kamel; GlaxoSmithKline, Uxbridge, UK: Margaret Tabberer; Katholieke Universiteit Leuven, Leuven, Belgium: Thierry Troosters and Fabienne Dobbels; Municipal Institute of Medical Research, Barcelona, Spain: Judith Garcia-Aymerich; Netherlands Asthma Foundation, Amersfoort, The Netherlands: Pim de Boer; Novartis, Basel, Switzerland: Karoly Kulich and Alastair Glendenning; Pfizer Walton Oaks, UK: Katja Rudell and Frederick J. Wilson; Royal Brompton and Harefield NHS Foundation Trust, London, UK: Michael I. Polkey and Nick S. Hopkinson; Thorax Research Foundation, Athens, Greece: Ioannis Vogiatzis; UCB, Brussels, Belgium: Enkeleida Nikai; University Medical Center, Groningen, The Netherlands: Thys van der Molen and Corina De Jong; University of Edinburgh, Edinburgh, UK: Roberto A. Rabinovich and Bill MacNee; University of Zurich, Zurich, Switzerland: Milo A. Puhan and Anja Frei.

References

1. Van Remoortel H, Hornikx M, Demeyer H, Langer D, Burtin C, Decramer M, et al. Daily physical activity in subjects with newly diagnosed COPD. *Thorax*. 2013;68:962–3.
2. Pitta F, Troosters T, Spruit MA, Probst VS, Decramer M, Gosselink R. Characteristics of physical activities in daily life in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med*. 2005;171:972–7.
3. Shrikishna D, Patel M, Tanner RJ, Seymour JM, Connolly BA, Puthuchery ZA, et al. Quadriceps wasting and physical inactivity in patients with COPD. *Eur Respir J*. 2012;40:1115–22.
4. Gimeno-Santos E, Frei A, Steurer-Stey C, De Batlle J, Rabinovich RA, Raste Y, et al. Determinants and outcomes of physical activity in patients with COPD: a systematic review on behalf of PROactive consortium. *Thorax*. 2014;69:731–9.
5. Demeyer H, Donaire-Gonzalez D, Gimeno-Santos E, Ramon MA, De Batlle J, Benet M, et al. Physical activity is associated with attenuated disease progression in chronic obstructive pulmonary disease. *Med Sci Sports Exerc*. 2019;51:833–40.
6. Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease: 2020 REPORT. www.goldcopd.org [accessed 12.11.19].
7. Pleguezuelos E, Gimeno-Santos E, Hernández C, Mata M, del C, Palacios L, Piñera P, et al. Recommendations on non-pharmacological treatment in chronic obstructive pulmonary disease from the Spanish COPD Guidelines (GesEPOC 2017). *Arch Bronconeumol*. 2018;54:568–75.
8. Waschki B, Kirsten AM, Holz O, Mueller K-C, Schaper M, Sack A-L, et al. Disease progression and changes in physical activity in patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med*. 2015;192:295–306.
9. Agarwal V, Tetenta S, Bautista J, ZuWallack R, Lahiri B. Longitudinal changes in directly measured physical activity in patients with chronic obstructive pulmonary disease. *J Cardiopulm Rehabil Prev*. 2012;32:292–5.
10. Clarenbach CF, Sievi NA, Haile SR, Brack T, Brutsche MH, Frey M, et al. Determinants of annual change in physical activity in COPD. *Respirology*. 2017;22:1133–9.
11. Sievi NA, Brack T, Brutsche MH, Frey M, Irani S, Leuppi JD, et al. Physical activity declines in COPD while exercise capacity remains stable: a longitudinal study over 5 years. *Respir Med*. 2018;141:1–6.
12. Moy ML, Danilack VA, Weston NA, Garshick E. Daily step counts in a US cohort with COPD. *Respir Med*. 2012;106:962–9.
13. Troosters T, Blondeel A, Rodrigues FM, Janssens W, Demeyer H. Strategies to increase physical activity in chronic respiratory diseases. *Clin Chest Med*. 2019;40:397–404.
14. Boutou AK, Raste Y, Demeyer H, Troosters T, Polkey MI, Vogiatzis I, et al. Progression of physical inactivity in COPD patients: the effect of time and climate conditions – a multicenter prospective cohort study. *Int J Chron Obstruct Pulmon Dis*. 2019;14:1979–92.
15. Alahmari AD, Patel AR, Kowlessar BS, Mackay AJ, Singh R, Wedzicha JA, et al. Daily activity during stability and exacerbation of chronic obstructive pulmonary disease. *BMC Pulm Med*. 2014;14:1–8.
16. Demeyer H, Costilla-frias M, Louvaris Z, Gimeno-Santos E, Tabberer M, Rabinovich RA, et al. Both moderate and severe exacerbations accelerate physical activity decline in COPD patients. *Eur Respir J*. 2018;51:1702110.
17. Sievi NA, Kohler M, Thurnheer R, Leuppi JD, Irani S, Frey M, et al. No impact of exacerbation frequency and severity on the physical activity decline in COPD: a long-term observation. *Int J Chron Obstruct Pulmon Dis*. 2019;14:431–7.
18. Couto Furlanetto K, Demeyer H, Sant'anna T, Aparecida Hernandez N, Augusto Camillo C, Serra Pons I, et al. Physical activity of patients with COPD from regions with different climatic variations physical activity of patients with COPD from regions with different climatic variations. *COPD J Chronic Obstr Pulm Dis*. 2017;14:276–83.
19. Agustí A, Hogg JC. Update on the pathogenesis of Chronic Obstructive Pulmonary Disease. *N Engl J Med*. 2019;381:1248–56.
20. Saunders TJ, Dechman G, Hernandez P, Spence JC, Rhodes RE, McGannon K, et al. Distinct trajectories of physical activity among patients with COPD during and after pulmonary rehabilitation. *COPD J Chronic Obstr Pulm Dis*. 2015;12:539–45.
21. Soicher JE, Mayo NE, Gauvin L, Hanley JA, Bernard S, Maltais F, et al. Trajectories of endurance activity following pulmonary rehabilitation in COPD patients. *Eur Respir J*. 2012;39:272–8.
22. Arbillaga-Etxarri A, Gimeno-Santos E, Barberan-Garcia A, Balcells E, Benet M, Borrell E, et al. Long-term efficacy and effectiveness of a behavioural and community-based exercise intervention (Urban Training) to increase physical activity in patients with COPD: a randomised controlled trial. *Eur Respir J*. 2018;52:1800063.
23. Gimeno-Santos E, Raste Y, Demeyer H, Louvaris Z, De Jong C, Rabinovich RA, et al. The PROactive instruments to measure physical activity in patients with chronic obstructive pulmonary disease. *Eur Respir J*. 2015;46:988–1000.
24. Celli BR, Macnee W, Agusti A, Anzueto A, Berg B, Buist AS, et al. Standards for the diagnosis and treatment of patients with COPD: a summary of the ATS/ERS position paper. *Eur Respir J*. 2004;23:932–46.
25. Rabinovich RA, Louvaris Z, Raste Y, Langer D, Van Remoortel H, Giavedoni S, et al. Validity of physical activity monitors during daily life in patients with COPD. *Eur Respir J*. 2013;42:1205–15.
26. Demeyer H, Burtin C, Van Remoortel H, Hornikx M, Langer D, Decramer M, et al. Standardizing the analysis of physical activity in patients with COPD following a pulmonary rehabilitation program. *Chest*. 2014;146:318–27.
27. Arbillaga-Etxarri A, Gimeno-Santos E, Barberan-Garcia A, Benet M, Borrell E, Dadvand P, et al. Socio-environmental correlates of physical activity in patients with chronic obstructive pulmonary disease (COPD). *Thorax*. 2017;0:1–7.
28. Genolini, Christophe Falissard B. KML: k-means for longitudinal data. *Comput Stat*. 2010;25:317–28.
29. Rabe-Hesketh S, Skrondal A, Pickles A. Maximum likelihood estimation of limited and discrete dependent variable models with nested random effects. *J Econom*. 2005;128:301–23.
30. Demmelmaier I, Dufour AB, Nordgren B, Opava CH. Trajectories of physical activity over two years in persons with rheumatoid arthritis. *Arthritis Care Res*. 2016;68:1069–77.
31. Demeyer H, Burtin C, Hornikx M, Camillo CA, Van Remoortel H, Langer D, et al. The minimal important difference in physical activity in patients with COPD. *PLOS ONE*. 2016;11:e0154587.
32. Dubé BP, Vermeulen F, Laveneziana P. Exertional dyspnoea in chronic respiratory diseases: from physiology to clinical application. *Arch Bronconeumol*. 2017;53:62–70.
33. Hoaas H, Zanaboni P, Hjalmarsen A, Morseth B, Dinesen B, Burge AT, et al. Seasonal variations in objectively assessed physical activity among people with COPD in two Nordic countries and Australia: a cross-sectional study. *Int J Chron Obstruct Pulmon Dis*. 2019;14:1219–28.
34. Burge AT, Cox NS, Abramson MJ, Holland AE. Interventions for promoting physical activity in people with chronic obstructive pulmonary disease (COPD). *Cochrane Database Syst Rev*. 2020. Art. No.: CD012626.
35. Miravittles M, Troosters T, Janssens W, Ancochea J. Multidisciplinary perspectives on the importance of physical activity in COPD. *Arch Bronconeumol*. 2019;55:551–2.
36. Bauman AE, Reis RS, Sallis JF, Wells JC, Loos F, Martin RJBW, et al. Correlates of physical activity: why are some people physically active and others not? *Lancet*. 2012;380:258–71.
37. Coultas DB, Jackson BE, Russo R, Peoples J, Sloan J, Singh KP, et al. A lifestyle physical activity intervention for patients with chronic obstructive pulmonary disease a randomized controlled trial. *Ann Am Thorac Soc*. 2016;13:617–26.
38. Altenburg WA, Ten Hacken NHT, Bossenbroek L, Kerstjens HAM, De Greef MHG, Wempe JB. Short- and long-term effects of a physical activity counselling programme in COPD: A randomized controlled trial. *Respir Med*. 2015;109:112–21.