

with numerous flagellae, the longest in the center, in an irregular arrangement. No terminal bar is observed, but a small nucleus can be seen in the apical pole just below the flagellar insertion.

Because this protozoan is easily confused with ciliated epithelial cells on direct microscopy of fresh samples, we recommend using specific staining techniques (Wheatley trichrome, Giemsa or Papanicolaou) to assist in identification. The current lack of an appropriate culture method for this parasite could be supplemented by molecular biology techniques.

Funding

This study has not received specific grants from public sector agencies, the commercial sector, or non-profit organizations.

Conflict of interests

The authors state that they have no conflict of interests.

References

1. Pinos Vélez N, Ordoñez Vintimilla R, Agreda Orellana S. Infección pulmonar por *Lophomonas blattarum*. Arch Bronconeumol. 2020; <http://dx.doi.org/10.1016/j.arbres.2020.12.005>.
2. Li R, Gao ZC. *Lophomonas blattarum* infection or just the movement of ciliated epithelial cells? Chin Med J (Engl). 2016;129:739–42. <http://dx.doi.org/10.4103/0366-6999.178025>.
3. Alam-Eldin YH, Abdulaziz AM. Identification criteria of the rare multi-flagellate *Lophomonas blattarum*: comparison of different staining techniques. Parasitol Res. 2015;114:3309–14. <http://dx.doi.org/10.1007/s00436-015-4554-4>.
4. Fakhar M, Nakhaei M, Sharifpour A, Kalani H, Banimostafavi ES, Abedi S, et al. First molecular diagnosis of Lophomoniasis: the end of a controversial history. Acta Parasitol. 2019;64:390–3. <http://dx.doi.org/10.2478/s11686-019-00084-2>.
5. Martínez-Girón R, Martínez-Torre C, van Woerden HC. The prevalence of protozoa in the gut of German cockroaches (*Blattella germanica*) with special reference to *Lophomonas blattarum*. Parasitol Res. 2017;116:3205–10. <http://dx.doi.org/10.1007/s00436-017-5640-6>.
6. van Woerden HC, Martínez-Girón R, Martínez-Torre C. Protozoan cysts in faecal pellets of German cockroaches (*Blattella germanica*), with particular emphasis on *Lophomonas blattarum*. Acta Parasitol. 2020;65:831–6. <http://dx.doi.org/10.2478/s11686-020-00213-2>.

Rafael Martínez-Girón,^{a,*} Cristina Martínez-Torre^b

^a Fundación INCLÍNICA, Oviedo, Spain

^b Facultad de Farmacia, Universidad de Salamanca, Salamanca, Spain

* Corresponding author.

E-mail address: rmartinezgiron@hotmail.com (R. Martínez-Girón).

<https://doi.org/10.1016/j.arbr.2021.09.008>

1579-2129/ © 2021 SEPAR. Published by Elsevier España, S.L.U. All rights reserved.

Reply to “Lophomonas or Ciliated Epithelial Cells?”[☆]



Respuesta a «¿Lophomonas o células epiteliales ciliadas?»

To the Editor:

We have read in detail the letter published by Martínez-Girón et al. entitled “Lophomonas or ciliated epithelial cells?”¹, in which they question the evidence of the findings of flagellated parasites in patient samples from bronchoscopy samples.¹

It is true that it is unusual to find this type of protozoan in bronchial lavage samples and it is also true to say that they may go unnoticed due to the lack of expertise of the microscopist, as they can be easily confused with ciliated cells originating in the bronchial tree. No specific culture media have been identified to date, but molecular techniques for confirming lophomoniasis were published by Fakhar et al. in 2019².

We reported the presence of *Lophomona* sp. in patient specimens using the wet-mount technique with 400× magnification post-centrifugation and subsequent Giemsa staining. This was used to distinguish between ciliated respiratory cells and the protozoan and also to analyze various morphological characteristics of the parasite, including its rounded or ovoid form (20–60 mm long × 12–20 mm wide); the double strand of flagellae at the anterior end; the absence of a terminal bar; and a certain cytoplasmic plasticity. It also helped detect the presence of thick granules and some vacuoles along with a difficult-to-visualize nucleus and the main characteristic of these protozoa, namely, asynchronous movements that generate vibratory, rotational and revolving movements in the protozoan cytoplasm (Fig. 1).

These characteristics distinguish them from ciliated cells originating in the tracheobronchial tree, which are characterized by a basophilic cytoplasm, a basal end with a sharp conical insertion, and an apical end with a reinforced edge corresponding to the terminal bar from which the cilia arise. The nucleus is central, round, or oval with fine chromatin and a discrete nucleolus can be seen^{3,4}.

Martínez-Girón et al. also mention that the protozoa were perhaps confused with the phenomenon called ciliocytophthoria, which is defined as a degenerative process of the ciliated cells as a consequence of viral infections and characterized by typical morphological changes. As we stated in our article, our lophomoniasis cases did not present any viral process at the time of diagnosis nor did we detect any evidence of carcinoma, another typical cause of these phenomena⁵.




Fig. 1. 400× magnification of slide-mounted bronchial brushing sample.

[☆] Please cite this article as: Agreda Orellana S, Pinos Vélez N. Respuesta a «¿Lophomonas o células epiteliales ciliadas?». Archivos de Bronconeumología. 2021;57:727–728.

It is true that the species of *Lophomonas* sp. cannot be distinguished in fresh mounts, but they can be distinguished from ciliated

cells using stains such as Giemsa and by observing the internal characteristics of the protozoan.

References

- Martínez-Girón R, Martínez-Torre C. ¿Lophomonas o células epiteliales cilidas? Arch Bronconeumol. 2021. <http://dx.doi.org/10.1016/j.arbres.2021.03.006>.
- Fakhar M, Nakhaei M, Sharifpour A, Kalani H, Banimostafavi ES, Abedi S, et al. First molecular diagnosis of lophomoniasis: the end of a controversial story. Acta Parasit. 2019;64:390–3. <http://dx.doi.org/10.2478/s11686-019-00084-2>.
- Van Woerden HC, Martínez-Girón R. *Lophomonas blattarum*: is it only its morphology that prevents its recognition? Chin Med J (Engl). 2017;130:117. <http://dx.doi.org/10.4103/0366-6999.196579>.
- Fichas de citología SA. Ciliocitoforia. Sociedad Argentina de Citología [Accessed 30 March 2021]. Available from: <http://sociedaddecitologia.org.ar/sac/ciliocitoforia/>.

- Matteo G, Giorgio C. Ciliocytophthoria of nasal epithelial cells after viral infection: a sign of suffering cell. Acta Biomed. 2019;90:7–9. <http://dx.doi.org/10.23750/abm.v90i2-5.8103>.

Solmayra Agreda Orellana,^b Nathalie Pinos Vélez^{a,b,*}

^a Hospital José Carrasco Arteaga, Cuenca, Ecuador

^b Hospital Universitario del Río, Cuenca, Ecuador

* Corresponding author.

E-mail address: nathaliepinos@hotmail.com (N. Pinos Vélez).

<https://doi.org/10.1016/j.arbr.2021.09.014>

1579-2129/ © 2021 SEPAR. Published by Elsevier España, S.L.U. All rights reserved.

Prevalence and Mortality of Patients with Palliative Needs in an Acute Respiratory Setting[☆]



Prevalencia y mortalidad de pacientes con necesidades paliativas en una planta de Neumología

To the Editor:

We were pleased to read the article entitled “Prevalence and Mortality of Patients with Palliative Needs in an Acute Respiratory Setting”, published by Eva Taberner Huguet et al.¹.

It can be difficult to identify end-of-life situations in COPD patients, given the multiple exacerbations they overcome successfully during the course of their disease, but we believe that efforts must be made to identify patients beyond cancer sufferers who require palliative care. We agree with the authors on the need to extend knowledge of palliative medicine to professionals who treat chronic diseases, since the vast majority of our hospitals have limited access to palliative teams, and it is impossible to offer this care to all patients who require it.

The authors make the important point that mortality differs little between cancer and non-cancer patients, but it is also clear that the symptom burden of patients with advanced COPD is similar to that of cancer patients² and, as such, this population would benefit from being treated by medical specialists who are familiar with non-oncological palliation, an approach that would change the perception of death as failure. We applaud the authors' initiative to highlight the need for training in the field of palliation in a disease such as COPD, a true model of chronicity. This may lead to better patient care, better quality of care, and better communication with patients and their families in end-of-life situations.

The NECPAL instrument³ is a screening tool for patients with palliative needs, but in order to be able to respond adequately to the Surprise Question, we must improve our understanding of the prognostic factors. The profile of COPD patients that could die within 6–12 months includes older age, limited physical activity, high consumption of health resources⁴, and general status determined by comorbidities and a BODE score of ≥ 7 . We also know that dependency is a factor that can predict mortality more reliably than indices such as Charlson⁵.

The mean age of the series presented by the authors is 76 years, and their multimorbidity is considerable (76 patients had > 2 chronic diseases). It is therefore mandatory to calculate a Barthel index and perform a geriatric assessment, generating a diagnosis of the patient's status that includes geriatric syndromes, in order to help recognize the palliative needs of patients.

Finally, the basic criterion for initiating palliative care must be the refractoriness of symptoms to standard treatment, adjusted to the patient's preferences, leaving the survival estimates in the background. Our rapidly aging, pluripathological population demands a new view of patients with chronic diseases such as advanced COPD and a shift towards a medicine that focuses on the patient and their needs and not on their life expectancy.

References

- Taberner Huguet E, Ortiz de Urbina Antia B, González Quero B, Garay Llorente E, Andia Iturrate J, Pérez Fernández S, et al. Prevalence and Mortality of Patients with Palliative Needs in an Acute Respiratory Setting. Arch Bronconeumol. 2021;57:345–50. <http://dx.doi.org/10.1016/j.arbres.2020.08.009>.
- Wilson IM, Bunting JS, Curnow RN, Knock J. The need for inpatient palliative care facilities for noncancer patients in the Thames Valley. Palliat Med. 1995;9:13–8. <http://dx.doi.org/10.1177/026921639500900103>.
- Gómez-Batiste X, Martínez-Muñoz M, Blay C, Amblàs J, Vila L, Costa X, et al. Utility of the NECPAL CCOMS-ICO© tool and the Surprise Question as screening tools for early palliative care and to predict mortality in patients with advanced chronic conditions: A cohort study. Palliat Med. 2017;31:754–63. <http://dx.doi.org/10.1177/0269216316676647>.
- Patel K, Janssen DJ, Curtis JR. Advance care planning in COPD. Respirology. 2012;17:72–8. <http://dx.doi.org/10.1111/j.1440-1843.2011.02087.x>.
- Fernández-García S, Represas-Represas C, Ruano-Raviña A, Botana-Rial M, Martínez-Reglero C, Fernández Villar A. Dependence IN Performing Activities as a Predictor of Mortality Following Hospitalization for Chronic Obstructive Pulmonary Disease Exacerbation. Arch Bronconeumol. 2020;56:291–7. <http://dx.doi.org/10.1016/j.arbres.2019.10.005>.

Isabel Torrente Jiménez,^{a,b,*} Macarena Cabrera Pajarón,^c

Marc Moreno-Ariño,^a Mariona Palou Campmoll,^d

Ricard Comet Monte^a

^a Unidad de Geriátrica de Agudos y Paciente Crónico Complejo. Corporació Sanitària Parc Taulí, Sabadell, Spain

^b Instituto de Investigación e Innovación Parc Taulí (I3PT), Sabadell, Spain

^c Unidad de Convalecencia. Corporació Sanitària Parc Taulí, Sabadell, Spain

^d Servicio de Medicina Interna. Corporació Sanitària Parc Taulí, Sabadell, Spain

* Corresponding author.

E-mail address: itorrente@tauli.cat (I. Torrente Jiménez).

<https://doi.org/10.1016/j.arbr.2021.09.010>

1579-2129/ © 2021 SEPAR. Published by Elsevier España, S.L.U. All rights reserved.

[☆] Please cite this article as: Torrente Jiménez I, Cabrera Pajarón M, Moreno-Ariño M, Palou Campmoll M, Comet Monte R. Prevalencia y mortalidad de pacientes con necesidades paliativas en una planta de Neumología. Archivos de Bronconeumología. 2021;57:728–728.